¹Clinical Psychology, Bambino Gesù Children Hospital, Rome, Italy and ²Neurological Sciences, Bambino Gesù Children Hospital, Rome, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1780

Introduction: Anxious-depressive disorders are common among children with epilepsy. A recent Systematic Review and Meta-Analysis (Scott et al., 2020) reported that the overall pooled prevalence of anxiety disorders is 18.9% while of depressive disorders is 13.5%. COVID-19 pandemic has centralized the attention of governors and careers on the health emergency. As a result, the trajectory of the psychological care needs of this at risk population may have been neglected.

Objectives: The aim of the study was to assess the prevalence rate of depressive and anxiety symptoms among children with epilepsy during COVID-19 pandemic. Children were hospitalized in- and out-patients under a neurological and psychological follow up program in an Italian Children Hospital.

Methods: We conducted a cross-sectional study among 38 hospitalized children and adolescents with epilepsy (21F; 17M, mean age: 14.5; range: 11-18) during COVID-19 pandemic. We performed face-to face interviews and assessed depressive and anxiety symptoms with the Patient Health Questionnaire (PHQ-9) and the Generalized Anxiety Disorders (GAD-7) questionnaire during scheduled follow up checks.

Results: Preliminary results showed a rate of mild-to-severe anxious depressive symptoms by 49.9% and 60.5% respectively. In detail: 21.1% mild, 15.7% moderate and 13.1% of severe anxiety, meanwhile 23.7% mild, 26.3% moderate and 10.5% of severe depression. The prevalence of comorbid depressive and anxiety symptoms was 39.5% among the entire sample.

Conclusions: Depressive and anxiety rates among hospitalized children with epilepsy during COVID-19 outbreak are very high. Pediatric services should deserve special attention to those patients' mental health. Regular screening protocols and empowerment interventions in Hospital should be promoted.

Disclosure: No significant relationships.

Keywords: COVID-19; Epilepsy; Anxiety; Depression

EPV0196

A study on patterns of use of mobile phone and nomophobia in medical undergraduate students during a COVID-19 pandemic lockdown

P. Bhatnagar*, S. Tarachandra and S. Undaru

Psychiatry, K.S. Hegde Medical Academy, Mangalore, India

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1781

Introduction: The governments of various countries enforced a lockdown to contain the COVID -19 pandemic. As the colleges remain closed, the academic teachings for students was conducted online. The mobile phone remained the main source for academics and entertainment during this period.

Objectives: To assess patterns of use of mobile phone by Medical Undergraduate students prior to and during the COVID-19 lockdown. To assess Nomophobia among same participants.

Methods: This study was done by an online survey method after obtaining approval from the Institutional Ethics Committee. A validated questionnaire on patterns of mobile phone use and the Nomophobia Questionnaire(NMP-Q) was completed by the medical students (n=187) who consented to participate in the study

Results: Prior to the pandemic lockdown, 52.9% of the participants used the mobile phones for 2-4 hours per day with 78% of the usage in social media. During lockdown, 89.3% of the participants reported an increase in the usage of mobile phones. 35.65% reported an increase in use by 2-4 hours everyday. About 30.5 % used the mobile phone for 6-8 hours per day. 80.2 % reported a maximum usage for social media. 59.45% reported a maximum usage for online academics. 33.7% frequently checked their phones once in 15 minutes. About 60.43% of the participants were in the moderate and 21.4% in the severe category of nomophobia.

Conclusions: There is an increase in mobile phone usage during the lockdown with a significant proportion of students in the moderate and severe category of nomophobia.

Disclosure: No significant relationships.

Keywords: nomophobia; COVID-19 lockdown; Medical Students; mobile phone

EPV0197

Psychosis and COVID-19: About a case

T. Ponte*, S. Castela-Almodóvar, B. Sanz-Aranguez, E. Gil-Benito and J. Herranz

Psiquiatría, Hospital Universitario Puerta de Hierro, Madrid, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1782

Introduction: Cases of psychosis are being reported in people infected by the SARS-CoV-2 virus. The relationship between psychosis and corticosteroids treatment is well known. However, there are relatively limited data so far correlating psychosis and SARS-CoV-2.

Objectives: To describe a case of manic psychosis in a 55-year-old woman treated with corticosteroids for COVID-19 infection. Discuss the etiopathogenic factors involved in psychosis in patients infected by COVID-19.

Methods: We present the case of a 55-year-old woman, without previous psychiatric history, who was admitted to psychiatry due to a psychotic episode with maniac symptoms. Three weeks earlier, the patient had been admitted to Internal Medicine for bilateral SARS-CoV2 pneumonia, under treatment with high doses of corticosteroids. The patient presents a verbose and salty speech, euphoric mood with hyperergia, subjective increase of capacities, insomnia and delusional ideation with mystical-spiritual content with delusional interpretations and auditory hallucinations. The patient comes from Ukraine and she has been living in Spain for 20 years. She works as a household assistant. The patient relates various psychosocial stressors throughout her life.

Results: Complementary diagnostic tests were without alterations. Low-dose antipsychotic treatment is prescribed, with a rapid recovery within a week. Finally, the patient showed complete insight of the episode and was discharged from the hospital being asymptomatic.

Conclusions: It would be interesting to publish the reported cases of psychosis and infection by COVID-19 as well as to investigate the