

On the origin of solar and stellar flares

Subhon Ibadov^{1,2} and Firuz S. Ibodov¹

¹Moscow State University, Sternberg Astronomical Institute
119991 Moscow, Russia
email: mshtf@sai.msu.ru

²Institute of Astrophysics, Tajik Academy of Sciences,
734042 Dushanbe, Tajikistan
email: ibadovsu@yandex.ru

Abstract. Solar and stellar flares due to impacts of comet nuclei and falling evaporating bodies, FEBs, with the Sun/stars are analytically considered. It is shown that impacts of sun/stargrazing comets will be accompanied by essential aerodynamic effects: nuclei crushing and expansion/flattening of crushed mass within the chromosphere. These processes lead to impulse generation of a hot plasma, strong shock wave in the thin layer near photosphere, eruption of the hot ionized clump to the lower corona, i.e., impact-induced solar/stellar flares.

Keywords. Sun/stargrazing comets, falling evaporating bodies/FEBs, impacts, stellar flares

1. Introduction

Data from solar space missions (SOLWIND, SMM, SOHO, STEREO, SDO) and ground-based observations as well as spectral observations of young stars like Beta Pic-toris together with celestial mechanics calculations indicate the presence of sun/stargrazing comet/FEB fluxes passing close to the solar/stellar surface, at distances less than 0.01 AU (e.g., Marsden 1989, Bailey *et al.* 1992, Beust *et al.* 1996, <http://sungrazer.nrl.navy.mil/>). Besides, calculations show that solar photospheric thermal radiation leads to a very limited decrease in the cometary nuclei radii, less than 10-20 m, in the 'classical'/vacuum approach (e.g. Weissman 1983, MacQueen & St.Cyr 1991).

Meantime, parabolic velocity of comets near the solar surface is $V_o = 617$ km/s, while the density of the solar atmosphere within the chromosphere considerably exceeds 10^{-15} g/cm³ being $10^{-8} - 5 \times 10^{-7}$ g/cm³ in the photosphere, so that aerodynamic pressure on the comet nuclei will considerably exceed the tension for mechanical dis-integration. At the same time the specific kinetic energy of comet nuclei in the inner heliosphere, especially near the surface of the Sun, more than thousand times exceeds the evaporation/sublimation energy of comet nuclei material, i.e., of the order of 10^{10} erg/g. Hence, high-temperature explosive phenomena due to comet/FEBs impacts with the Sun/stars are possible and should be considered as one of possible processes giving rise to solar/stellar flares: there are well-known planetary analogs as the 1908 Tunguska and 2013 Chelyabinsk air explosions (Grigorian 1980, Ibadov *et al.* 2008, Ibadov *et al.* 2009, Grigorian *et al.* 2013 and references therein).

We present completely analytic approach in investigating the passage of comet nuclei/FEBs through the stars atmosphere for revealing peculiarities of an impact mechanism of solar/stellar flares.

2. Aerodynamic heating and crushing of comet nuclei/FEBs in solar/stellar atmospheres

For a comet approaching the Sun the critical density of the solar atmosphere at which the energy flux on the surface of the nucleus due to collisions with atoms of the solar atmosphere becomes equal to that due to irradiation by the solar photosphere thermal emission will be $\rho_{cr} = L_o/(2\pi R_o^2 V^3 = 5 \times 10^{-13} \text{ g/cm}^3$ with L_o and R_o as the solar luminosity and radius, respectively. The numerical value obtained corresponds to the density of the solar atmosphere near the upper boundary of the chromosphere.

A criterion for the onset of a comet nucleus aerodynamic crushing $P_{a*} = \rho_{a*} V_*^2 = \sigma_*$ together with the equation for aerodynamic deceleration of a constant-mass comet nucleus, i.e., that having sufficiently large initial radius, $R_n > 100 \text{ m}$, written as

$$M \frac{dV}{dt} = \frac{1}{2} C_x S \rho_a V^2 \quad (2.1)$$

allow us to find the onset height for the nucleus aerodynamic destruction, namely

$$h_* = H \ln \left(\frac{\rho_o V_o^2}{\sigma} \right). \quad (2.2)$$

Here M , C_x and S are the mass, drag coefficient and cross-section of a comet nucleus; $\rho_a(h) = \rho_o \exp(-h/H)$ is the density distribution of atmosphere; h is the height for which the limb of the Sun's disk with the optical density $\tau_\lambda(\lambda = 5000 \text{ \AA}) = 0.005$ is taken as the reference point and $\rho_o = \rho_a(h = 0) = 10^{-8} \text{ g/cm}^3$; H is the local height scale: $H = 3 \times 10^3 \text{ km}$ and 200 km for the chromosphere and photosphere, respectively; V , V_o are the nucleus velocity in the atmosphere and its initial velocity; P_{a*} , ρ_{a*} , V_* are the frontal pressure of the incoming atmospheric gas, mass density of the atmosphere, comet nucleus velocity - all these three values correspond to the onset of the nucleus aerodynamic crushing, σ_* is the strength of the nucleus material.

Substituting into (2.2) $H = 3 \times 10^8 \text{ cm}$, $\rho_o = 10^{-8} \text{ g/cm}^3$, $\sigma_* = 10^4 \text{ dyn/cm}^2$ we obtain the crush-height in atmosphere $h_* = 10^9 \text{ cm}$. The corresponding characteristic atmosphere density will be $\rho_{a*} = \sigma_*/V_o^2 = 3 \times 10^{-12} \text{ g/cm}^3$. Accordingly, the characteristic intensity of energy flux onto the nucleus due to aerodynamic heating will be $(\rho_{a*} V_o^3)/2 = 3 \times 10^{11} \text{ erg/(cm}^2 \text{ s)}$: it is four times more than the maximum energy flux from the solar thermal radiation.

It should be noted that according to the solution of the deceleration Eq. (2.1) the velocity of large bodies above the solar photosphere will be practically constant, $V = V_o$.

So, the above analytic data indicate that within the solar chromosphere an intense aerodynamic crushing of comet nuclei occurs. Besides, the intensity of energy flux due to aerodynamic heating will be strongly more than by solar thermal radiation, i.e., the comet nuclei disintegration phenomenon near the Sun/stars acquires completely meteoric character (cf. Grigorian *et al.* 1997, Ibadov *et al.* 1999, Ibadov *et al.* 2007).

3. Aerodynamic explosion of FEBS in atmosphere: photospheric solar/stellar flares

The rapid rise of aerodynamic pressure at the entry of the nucleus into denser atmosphere layers leads to the situation when the destruction being started locally propagates through the nucleus body from the frontal surface to the rear one. At the same time the crushed mass loses its integrity, spreading in the lateral direction under the action of pressure gradient at the frontal surface. So, to take into account these aerodynamic effects we have to modify basic equations of the physical theory of meteors, like made

for the 1908 Tunguska explosion of the fireball/superbolide origin. Then the equation for aerodynamic deceleration, Eq. (1), for the completely fragmented transversally expanding nucleus/meteoroid in the constant-mass approximation, that is acceptable due to essential decrease of the coefficient for heat transfer at high temperatures for high-velocity large meteor bodies because of shielding effects (cf. Grigorian 1980), will be reduced to the following integral equation:

$$\int_{\tilde{V}}^V \frac{dV}{V} = -\frac{3C_x b \rho_o H}{8R_n^3 \nu \rho_n \sin \alpha} \int_{\tilde{r}}^r R^2 dr. \quad (3.1)$$

Here $R = R[r(h)]$ is the law for increase of the transverse radius of the flattening crushed mass:

$$\frac{R}{2R_n} = 1 + \frac{\sqrt{b/2}}{C} (\sqrt{1+r} - \sqrt{1+\tilde{r}}); \quad (3.2)$$

R_n , ρ_n are the initial radius and the density of the comet nucleus, respectively; α is the angle between the entry velocity of the nucleus into the atmosphere and the local horizon;

$$b = \nu \exp\left(-\frac{h_*}{H}\right); \quad \nu = \frac{3C_x \rho_o H}{4\rho_n R_n \sin \alpha}; \quad (3.3)$$

$$C = \left(\frac{3C_x R_n \sin \alpha}{8H}\right)^{1/2}; \quad r = \left[\exp\left(\frac{h_* - h}{H}\right)\right] - 1; \quad \tilde{r} \approx \frac{4C^2}{b}; \quad (3.4)$$

furthermore, ν is the parameter of the aerodynamic deceleration, it is very small for large bodies; \tilde{r} , \tilde{h} , \tilde{V} are the characteristic values of r , h and V , which correspond to the value of $R = 2R_n$, i.e., to the time instant when the nucleus is completely crushed and its transverse radius is equal to the doubled value of the initial radius.

An analytic solution of Eq. (3.1), using Eq. (3.2), for the region close to the endpoint of the deceleration trajectory in atmosphere, i.e., at small distances from the solar/stellar surface, $h \ll \tilde{h} \ll h_*$, $r \gg 1$, gives the law for velocity variation of completely crushed FEBs near the stellar photosphere in the form:

$$V = \tilde{V} \exp\left[-\frac{b^2}{2C^2}(r^2 - \tilde{r}^2)\right] = V_o \exp\left(-\frac{b^2}{2C^2}r^2\right). \quad (3.5)$$

From Eqs. (3.4), (3.5) we can find the height-range of the basic deceleration of the FEB, where the decrease of velocity from $V_1 = 0.9V_o$ to $V_2 = 0.1V_o$ occurs, namely

$$\Delta h_d = h_2 - h_1 = H \ln \frac{1+r_1}{1+r_2} = H \ln \frac{r_1}{r_2} \approx 0.7H. \quad (3.6)$$

Using Eqs. (2.1), (3.2), (3.4), (3.5) an equation for the rate of energy loose by the FEB in atmosphere may be obtained, in the form of dE/dr as a function of $r=r(h)$. On the basis of this equation explicit analytical expressions may be found for the parameters of the site where the maximum energy release due to aerodynamic deceleration of the crushed and flattening FEB, "explosion", in atmosphere takes place (cf. Grigorian *et al.* 2013):

$$r_m = r_{ex} = \frac{C}{b}, \quad (3.7)$$

$$h_m = h_{ex} = h_* - H \ln \left(1 + \frac{C}{b}\right) = H \ln \left[\frac{\rho_o V_o^2}{(1+C/b)\sigma_*}\right], \quad (3.8)$$

$$R_m = R_{ex} = 2R_n(1 + \frac{1}{\sqrt{2C}}), \tag{3.9}$$

$$V_m = V_{ex} = \frac{V_o}{\sqrt{e}}. \tag{3.10}$$

Accepting $C_x = 1$, $\rho_n = 0.5 \text{ g/cm}^3$, $R_n = 10^5 \text{ cm}$, $\sin \alpha = 0.5$, $\sigma_* = 10^6 \text{ dyn/cm}^2$ from (3.3), (3.4), (3.7), (3.8), (3.9) we have $\nu = 7 \times 10^{-5}$, $C = 7 \times 10^{-2}$, $b = 3.5 \times 10^{-6}$, $r_{ex} = 2 \times 10^4$, $h_{ex} = -7H = -1400 \text{ km}$, $R_{ex} = 7.4R_n = 7.4 \text{ km}$.

The explosive energy deposited in the zone of maximum deceleration, $E_m = E_{ex}$, is determined by the initial mass of the comet nucleus, M_o , coming into the atmosphere, and the mass of the atmosphere within the decelerating layer, ΔM_a . Usage of $\rho_a(h)$ and Eq.(3.6) gives $\Delta M_a \approx M_o$, so that with the help of Eq. (3.10) an analytic relation to estimate the energy of “explosion” may be found as:

$$E_{ex} = \frac{M_o \Delta M_a V_{ex}^2}{2(M_o + \Delta M_a)} = \frac{\rho_n R_n^3 V_o^2}{3e}. \tag{3.11}$$

On the basis of Eqs. (3.6) and (3.10) the characteristic time for thermalization of the kinetic energy of the fragmented nucleus may be estimated:

$$\tau_{ex} = \frac{\Delta h_d}{V_{ex}} = \frac{0.7\sqrt{e}H}{V_o} \approx \frac{H}{V_o} = 0.3 \text{ s}, \tag{3.12}$$

that explicitly indicates an impulsive and strongly explosive character of the energy release process in the near photosphere decelerating layer.

According to Eq. (3.11) the explosive energy will be around the energy of large solar flares, 10^{32} erg , for impacts of comets like comet Halley 1986 III while FEBs like comet Hale-Bopp 1995 OI can lead to superflares (cf. Grigorian *et al.* 2000, Ibadov *et al.* 2009, Eichler & Mordecai 2012).

The asymptotic velocity of expansion of a hot cylindrical plasma column in the near photosphere explosive layer, i.e., the initial velocity of a “blast” shock wave that will provide eruption of a hot plasma clump/plume from the solar/star surface may be determined using the analogy with the expansion of a short-living hot plasma clots, “compound particles”, produced by collision between two high-velocity dust particles (see, e.g., Ibadov 1990):

$$V_{sh} = \left[\frac{kT_o}{2\pi Am_p} + \frac{3k(1+z)T_o}{Am_p} \right]^{1/2}, \tag{3.13}$$

where T_o is the initial temperature of plasma in the explosive layer:

$$T_o = \frac{Am_p V_{ex}^2}{12k[(1+z+(2x_1)/3)]}, \tag{3.14}$$

A is the mean atomic number for FEB material and matter of photosphere, m_p is the proton mass, k is the Boltzmann constant, z is the mean multiplicity of ions charge, x_1 is the mean relative ionization potential (Ibadov 1986, 1996, 2011).

The maximum height of the photospheric mass ejections due to cometary impacts may be estimated as:

$$h_m = \frac{V_{sh}^2}{2g_o} = \frac{R_o^2 V_{sh}^2}{2GM_o}. \tag{3.15}$$

Here g_o is the gravity acceleration on the star’s surface, G is the gravity constant, M_o is the mass of the Sun/star.

It should be noted that “blast” wave generated in the near-photosphere decelerating layer can be considered as a “strong” shock wave. Indeed, the initial pressure in the front of shock wave generated due to explosive energy release in the layer, according to (3.6), (3.9), (3.11) will be

$$P_{sh} = \frac{E_{ex}}{\pi \Delta h_d R_{ex}^2} = 3 \times 10^5 \rho_n R_n \approx 10^{10} \text{ dyn/cm}^2 = 10^4 \text{ atm}, \quad (3.16)$$

that is greater than the ambient photospheric pressure more than 10^4 times, for $R_n \geq 10^5$ cm.

Using (3.13)–(3.15) with realistic values of $A = 20$, $z = 5$, $x_1 = 3$, $M_o = 2 \times 10^{33}$ g we get $T_o = 7 \times 10^6$ K, $V_{sh} = 1.7 \times 10^7$ cm/s, $h_m = 5 \times 10^9$ cm.

It is known that there are variety of solar prominences having maximum heights in the range 30–50 thousand kilometers (e.g. Filippov *et al.* 2006, Harrison *et al.* 2010). Hence, we can note that FEBs impact generated photospheric mass ejections can form a certain type of solar/stellar prominences having relatively high abundance of metal atoms (Ibadov & Ibadov 2011, Ibadov & Ibadov 2014).

The analytic approach presented had been tested explaining the HST data obtained during observations of ejecta from collision of comet SL 9 with Jupiter in 1994 (Hammel *et al.* 1995) and also studying the 2013 Chelyabinsk event (Grigorian *et al.* 2013 and references therein).

4. Conclusions

Impacts of comet nuclei/FEBs with the Sun/stars will lead to impulse generation of a hot plasma, “explosion”, strong “blast” wave, ejection of a hot plasma plume to the heights reaching the lower corona.

Impact mechanism of solar flares is capable to lead to solar/stellar superflares: it is reasonable coordinated observations/monitoring of short perihelion comets and nearby young stars abundant in FEBs.

The study of comet impact-generated high-temperature plasma phenomena in the solar photosphere/chromosphere by space telescopes like SDO, having high spatio-temporal resolutions, are of interest for the physics of solar/stellar flare activity as well as physics of comets.

5. Acknowledgements

The authors are grateful to SOC/LOC, 29th IAU General Assembly, for invitation (Honolulu, Hawaii, USA, 3-14 August 2015), SAI MSU for hospitality, Dr. G.M.Rudnitskij for useful discussions, Drs.E.Shimanovskaya and N.Gostev for technical support, as well as D. Talbott and S. Schirott for the support of the trip.

References

- Bailey, M. E. & Chambers, J. E., Hahn G. 1992, *A&A*, 257, 315
 Beust, H., Lagrange, A.-M., Plazy, F., & Mouillet, D. 1996, *A&A*, 310, 181
 COSPAR 1998, *COSPAR Inform. Bull.*, 142, 22
 Eichler, D. & Mordecai, D. 2012, *ApJL*, 761, L27
 Filippov, B. P., Zagnetko, A. M., & Ajabshirizadeh, A., Den O. G. 2006, *Sol. Sys. Res.*, 40, 319
 Grigorian, S. S. 1980, *Cosmic Res.*, 17, 724
 Grigorian, S. S., Ibadov, F. S., & Ibadov, S. 1997, *Phys.-Dokl.*, 42, 262

- Grigorian, S. S., Ibadov, S., & Ibodov, F. S. 2000, *Dokl.-Phys.*, 45, 463
- Grigorian, S. S., Ibodov, F. S., & Ibadov, S. 2013, *Sol. Sys. Res.*, 47, 268
- Hammel, H. B., Beebe, R. F., & Ingersoll, A. P., *et al.* 1995, *Science*, 267, 1288
- Harrison, R. A., Davis, C. J., & Bewsher, D., *et al.* 2010, *ASR*, 45, 1
- Ibadov, S. 1986, *ESA SP-250*, 1, 377
- Ibadov, S. 1990, *Icarus*, 86, 283
- Ibadov, S. 1996, *Physical Processes in Comets CI*, M., 181
- Ibadov, S. 2011, in: A. Manchado, L. Stanghellini, & D. Schonberner (eds.), *Proc. IAUS 283*, CUP, p.392
- Ibadov, S. & Ibodov, F. S. 2015, in: G. Meynet, C. Georgy, J. Groh & P. Stee (eds.), *Proc. IAUS 307*, p., 108
- Ibadov, S., Ibodov, F. S., & Grigorian, S. S. 2007, in: J. Bouvier & I. Appenzeller (eds.), *Proc. IAUS 243*, P.VI.4
- Ibadov, S., Ibodov, F. S., & Grigorian, S. S. 2008, in: V. Svetsov (ed.), *Proc. Intl. Conf. "100 Years Tunguska Phenomenon"*, RAS, M., p.45
- Ibadov, S., Ibodov, F. S. & Grigorian, S. S. 2009, in: N. Gopalswamy & D.F. Webb (eds.), *Proc. IAUS 257*, CUP, p.341
- Ibodov, F. S. & Ibadov, S. 2011, in: A. Bonanno & A. Kosovichev (eds.), *Proc. IAUS 274*, CUP, p.92
- Ibodov, F. S. & Ibadov, S. 2014, in: B. Schmieder, J.-M. Malherbe, & S.T. Wu, (eds.), *Proc. IAUS 300*, CUP, p. 509
- &MacQueen, R. M., St.Cyr, O. C. 1991, *Icarus*, 90, 96
- Marsden, B. G. 1989, *AJ*, 98, 2306
- Weissman, P. R. 1983, *Icarus*, 55, 448