# S22 Structure quality of mental health care systems in Europe Recent changes in the Hungarian mental health care system.

## Füredy, Janos (Budapest/Hungary)

Hungary is with ist ca 10 millions of inhabitants one of the smaller European country with a glorious past. Allthoug according to its image as a rural country more than the half of the inhabitants live in the five biggest towns. Before the political change during the last decade because of the communistic ideology the mental health area was widely neglected by the Hungarian Government. Inpatient care was provided by bigger state hospitals, whereas the outpatients care has been provided by governmental outpatients clinic. Since the political change and increasing number of psychiatrists begin to work in private practice and postgraduate training is improving. However, because of its famous psychoanalytical tradition for instance Ferency and Balint have been of Hungarian offspringpsychotherapeutic thinking has maintained in our county The orientation of Hungarian psychiatry has been directed more to Germany than to other countries. Now decentralization of hospital care as one form of reforms in Hungarian psychiatry is advancing, though the lack of finances is tremendous and the main obstacle to promote reforms quicker. The main part of psychiatrists is concentrated in the towns with the result of poor outpatient facility equipment in the rural areas A concept of improving the situation will be presented.

## S23 How to teach psychosocial skills to general practitioners? TEACHING PSYCHOSOMATIC MEDICINE - A TWO YEAR CURRICULUM FOR POST-GRADUATE CLINICIANS

#### Claus Buddeberg, Zurich, Switzerland.

25 to 30% of the patients seen in a general practitioner's office suffer from psychosomatic disorders. In Switzerland there is no structured teaching of a psychosomatic approach to illness, neither in medical school nor during postgraduate training. In this presentation experiences with a two year training program in psychosomatic medicine for post-graduate clinicians are reported. The training program has three parts: Part one consists of 12 training days over a two year period given in seminars with lectures and case presentations using videotapes, one way mirror as well as role plays. Part two is case supervision in small groups. In part three participants are taught and get experiences in a self relaxation technique. In our opinion the two year curriculum serves as a model and example for a certified training in "Psychomatic Medicine". The participants can benefit from their training in various ways: with regard to the organization and atmosphere of their practice, in respect of improved communication skills when treating patients and advanced small group interaction skills. We have just finished the second course and we received excellent feed-back from most of the participants.

#### S23 How to teach psychosocial skills to general practitioners? BEHAVIORAL SCIENCE CURRICULA FOR GENERAL INTERNAL MEDICINE: LEADING THE FEARFUL AND FREACHING TO THE CONVERTED

#### S. R. Hahn.

The goal of isaching psychiatry and behavioral sciences to primary care physician is uneque in both its importance and in the challenges it preserve. The goal between need and actual provide in the delivery of psychiatry treatment in the general medicine setting is well. Medical renderins' and factivity is still and knowledge and millingness to learn about psychiatry and behavior sciences are very heterogeneous menerging from "the factivit" in the conversed". However, the squalty of care the will be delivered in general medicine depends on enhancing the stills and knowledge of the majority of prestitionare, nor merely a handful of enhancements. The magnitude of the task discussion the behavior is carecoalise to the decise the two behavior is carecoalise to the stark of the stark of the task discussion of the task discussion of any stark of the stark of the task discussion of the task discussion of any stark of the task discussion of any stark of the task discussion of any stark of the stark of the task discussion of the task discussion of any stark of the task discussion of any stark of the stark of the task discussion of the task discussion of any stark of the stark discussion of the task discussion of the task discussion of the task discussion of the stark behavior is care tasks of most psychiatric problems that an inductate to task the father of these subject. Therefore we built a carriculum based on passes toscaris, exacted by professional action playing patients who have combination of psychiatric domiting using a mashed of least-stark of fathy starks and tasks approxing and the special problems. "Simulaid-patient' and anticide problem that a diverage of the semation. The simulaid-patient subspecialized facily, by contrast, the carriculum based on passes toscaris as exploremented by the cardinarce is assessment that a presense that an producting discussion process and psychiatry discussions are optimal three discussions of the stark of the stark diverse is astark and ta stark diverge of the sematers and t

S23 How to teach psychosocial skills to general practitioners? GETTING MEDICAL SPECIALISTS INVOLVED IN PSYCHOSOCIAL CARE: A ROLE FOR GENERAL INTERNISTS?

### <u>P. Cathebras</u> Service de Medicine Interne, Hôpitaux de St Etienne, 42055 St Etienne, France

Internal medicine holds a special position in France. Confined to tertiary care settings, it is considered as a speciality among other medical specialities. In the meantime, intensist favour a 'holistic' perspective very similar to the GP's approach, and opposed to most specialists' point of view. Since consultation-liaison psychiatry is poorly developed in many hospitals, internists sometimes have to act as consultants for clinical problems situated outside the organ specialities domain, including 'psychosomatic' problems. This means that internists have a special role in educating their fellow specialists on emotional factors in physical disease, attention to psychosocial context, patient-centred approach and emphasis on 'illness' vs. disease perspective in the management of symptoms. Our experience as internists as psychosocial specialists in the general hospital environment will be briefly discussed.