

Aims. There is still little information available on the negative impact of online activities on psychotic experiences. This limitation is further compounded for online gaming, where even a beneficial impact has been suggested via the evocation of positive emotions. We aimed to examine how problematic online gaming (POG) is associated with subsequent psychotic experiences in adolescents.

Methods. This birth cohort study employed randomly sampled adolescents born between September 2002 and August 2004. The eligibility criterion was those who did not have psychotic experiences at age 14. We analyzed the association between POG at age 14 and subsequent psychotic experiences at age 16. Adolescents were categorized into the no, low, and high POG groups based on the behaviors and emotions related to online gaming at age 14. Missing data were handled using random forest imputations.

Results. A total of 1722 adolescents without psychotic experiences at age 14 were analyzed. At age 16, 55 adolescents exhibited psychotic experiences, while 225 showed potential psychotic experiences. Compared with the no POG group, a higher risk of psychotic experiences was shown in both the low (RR 1.93, 95% CI 1.74–2.15) and high (RR 2.81, 95% CI 2.50–3.15) POG groups. Findings were consistent when analyzing potential psychotic experiences.

Conclusion. POG appears detrimental to the development of psychotic experiences in adolescents. Our findings provide public health implications in the context of policymaking.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Is Body Dissatisfaction a Risk Factor for Diabulimia and How Is It Assessed? A Rapid Systematic Review

Mr Navaneeth Natarajan* and Dr Shaymaa Obousy

King's College London, London, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.217

Aims. Diabulimia is an increasingly used term referring to “an eating disorder (ED) with type 1 diabetes”. It is difficult to detect and presents in multiple ways, which can potentially include feelings of body dissatisfaction (BD), which in itself is a complex symptom to quantify clinically.

This rapid systematic review aimed to identify whether feelings of BD are a risk factor for diabulimia by researching if and how BD is assessed in patients with the condition.

Methods. A rapid systematic review was undertaken. A literature review was performed on Ovid Medline (all) and Ovid Embase databases using search terms for Type 1 Diabetes, ED, and BD and looked at cross-sectional studies only. One reviewer performed the literature search and screened titles and abstracts. Out of 589 papers screened, four papers met the inclusion criteria. These papers then went through critical appraising using the Appraisal tool for Cross-Sectional Studies, with all papers showing mid-level quality clearing 16 to 17/20 questions. Therefore, data was extracted from all of them.

Results. All four papers came from different countries and used a wide range of sample sizes (43–477).

There was widespread heterogeneity between the data collected in each study due to the various tools used to identify BD, paired with differences in analysing extracted data.

To ensure transparency and quality of the results provided, the Synthesis Without Meta-analysis tool was used. Three studies looked at effects on adolescents and three had a higher proportion of

females. All papers used previously established and tested BD screening methods. Two papers found female diabetics were more likely to have BD symptoms, and one paper saw that males were more at risk. All four papers concluded that BD had some correlations with one or more aspects of diabetes and/or other ED symptoms related to diabulimia. Two commented on positive correlations between BD and HbA1c levels and one commented on BD symptoms and insulin restriction trending together. Two papers also saw BD symptoms and depressive symptoms correlating in patients as well.

Conclusion. All four studies showed that BD was related to diabulimia, both from a psychological and diabetic perspective, and most highlighted how BD manifested between the different sexes of diabetics. This review highlights the need for more standardised and comprehensive BD questionnaires to draw out key signs of EDs in diabetics that could improve screening, detection and management of diabulimia.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Cognitive Behavioural Therapy Versus Psychodynamic Therapy for Medically Unexplained Symptoms: A Retrospective Study of Healthcare Utilisation and Cost Analysis

Dr Rory Naylor*

University of Oxford, Oxford, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.218

Aims. Patients experiencing Medically Unexplained Symptoms (MUS) are some of the costliest in both primary and secondary care. Psychotherapy is one of the most efficacious ways of treating them although the most superior modality is unclear. Cognitive Behavioural Therapy (CBT) has the greatest evidence base, but a growing number of studies have investigated the role of Psychodynamic Psychotherapy (PPT). This is the first study to compare the two modalities concerning their impact on healthcare utilisation and cost to the NHS.

Methods. Patients referred to the Oxford Community Psychological Medicine Service in 2021 and who went on to complete a course of psychotherapy for MUS were included. 78 patients were referred, 66 patients were assessed, 16 patients began treatment and 9 patients completed treatment. 4 received CBT and 5 received PPT based on a ‘best fit’ assessment. Their healthcare utilisation (GP appointments, health investigations, A&E attendances, inpatient admissions and outpatient appointments) was assessed during the 6 months prior to their initial assessment and compared with the 6 months after therapy had ended using data from ‘Health Information Exchange’.

Results. Overall, psychotherapy reduced primary care use but our data was insufficiently powered for this to be statistically significant. There was a significant reduction in outpatient appointments after psychotherapy, mostly representing mental health consults.

Significant differences between pre-therapy and post-therapy were only observed for the number of health investigations in the PPT group which, surprisingly, increased with a large effect size ($d = 1.19$ 95% CI 1.12–2.88, $P = 0.03$). The same trend towards increased utilisation were observed for every outcome measure in PPT besides outpatient appointments. Conversely, all outcome measures showed an improvement after CBT apart from the number of health investigations which marginally increased.