

Sex steroid modulation of cortisol secretion in sheep

E. van Lier^{1,2†}, M. Carriquiry¹ and A. Meikle³

¹Dpto. de Producción Animal y Pasturas, Facultad de Agronomía, Universidad de la República, Avda. E. Garzón 780, 12900 Montevideo, Uruguay; ²Estación Experimental Facultad de Agronomía Salto, Ruta 31 Km 21, 50000 Salto, Uruguay; ³Laboratorio de Técnicas Nucleares, Facultad de Veterinaria, Universidad de la República, Lasplacas 1550, 11600 Montevideo, Uruguay

(Received 16 October 2013; Accepted 7 March 2014; First published online 4 April 2014)

There is strong evidence that the gonads modulate the hypothalamic–pituitary–adrenal axis. To investigate these sex differences at the adrenal glands of sheep we compared the cortisol response to ACTH (experiment 1) and measured the relative expression of oestrogen receptor alpha (ERS1), androgen receptor (AR), melanocortin 2 receptor (MC2R) and steroid acute regulatory protein (STAR) mRNA in adrenal glands (experiment 2) of gonadectomised rams and ewes either with or without sex steroid replacement. In experiment 1 six castrated adult rams and four ovariectomised adult ewes were used in two ACTH trials. On each trial blood samples were taken every 15 min for 4 h through an indwelling jugular catheter and each animal received 0.5 mg of an ACTH analogue i.v., immediately after the sample at 1 h from the beginning of the trial. Four days after the first trial the males received 100 mg of Testosterone Cyclopentilpropionate (TC) i.m. and the females received 2.5 mg of Oestradiol Benzoate (EB) i.m. At 72 h after TC or EB administration the second trial was performed. In experiment 2 the adrenal glands were obtained from gonadectomised adult rams (n = 8) and adult ewes (n = 8). Four rams received 100 mg of TC i.m. and four females received 0.5 mg of EB i.m. Blood samples were taken at 0, 12, 24, 48 and 72 h relative to steroid replacement and the animals were thereafter slaughtered. Cortisol, testosterone and 17 β -oestradiol were determined by radioimmunoanalysis. The transcripts of ERS1, AR, MC2R and STAR were determined by real-time reverse transcription PCR in adrenal tissue. Cortisol secretion was higher in female sheep than in male sheep, and higher in EB-treated than non-treated ewes. No difference in cortisol secretion was observed between TC-treated and non-treated rams. Gonadectomised rams treated with TC presented greater AR mRNA and MC2R mRNA expression than males without the steroid replacement. Gonadectomised ewes treated with EB tended to present lower AR mRNA than the ones without steroid replacement. Gonadectomised rams with TC also had greater AR mRNA, ERS1 mRNA and MC2R mRNA expression than ewes treated with EB. The relative amount of STAR transcript was not different among the different groups. The results confirm sex differences in ACTH-induced cortisol secretion in sheep, as well as in the expression of the receptor proteins for both 17 β -oestradiol and testosterone in the sheep adrenal gland. However, the underlying mechanisms for sex steroid modulation remain unresolved.

Keywords: sex steroids, gonadectomy, adrenal, receptors, sheep

Implications

Different stressors result in different responses from the hypothalamus–pituitary–adrenal (HPA) axis and stress can reduce reproductive performance in mammals. Considering the different roles of males and females in reproduction, it is reasonable to think that each sex has its own strategy for dealing with the effects of stress. There is abundant evidence that the gonads affect the way the HPA axis responds to stress. Understanding the regulatory pathways of sex steroid modulation of HPA activity will help to develop methods to reduce the impact of stress in animal husbandry and improve animal welfare.

Introduction

There is strong evidence that the gonads modulate the hypothalamus–pituitary–adrenal (HPA) axis. Sex steroids affect the HPA axis centrally, affecting the secretion of corticotropin-releasing hormone, arginine vasopressin and ACTH in rodents and primates (Burgess and Handa, 1992; Handa *et al.*, 1994; Young, 1995). The stress response of the HPA axis of female rats is greater than that of male rats and gonadal steroid levels are in part responsible for these differences (Handa *et al.*, 1994). This has been observed both under basal and stressful conditions in rats (Kitay, 1961; Viau and Meaney, 1991) and in primates (Uhart *et al.*, 2006). Oestrogens enhance ACTH secretion and testicular androgens suppress corticosterone secretion in the

[†] E-mail: evanlier@fagro.edu.uy

rat (Ogilvie and Rivier, 1997). However, most reports have focused on the central effect of steroids (central nervous system, Burgess and Handa, 1992; Young, 1995; Viau and Meaney, 1996). Direct effects of oestrogens on adrenal corticosterone secretion have been observed in ovariectomised female rats (Figueiredo *et al.*, 2007). In sheep, the response of the adrenal gland to ACTH administration – in terms of cortisol secretion – is also affected by sex: ewes secrete more cortisol than rams (Van Lier *et al.*, 2003a), which has also been found *in vitro* (Canny *et al.*, 1999). Gonadectomy seems to have an inverse effect on cortisol levels after ACTH treatment in both sexes; differences were observed in intact rams and ewes, while the cortisol response in gonadectomised rams and ewes was similar (Van Lier *et al.*, 2003a), although this is not always observed (Tilbrook *et al.*, 1999; Turner *et al.*, 2002). Therefore, sex differences in the stress response appear also to be expressed peripherally at the adrenal, and are probably partly due to the gonadal steroids. However, to our knowledge, no hormone replacement studies have been performed in sheep in order to test their effects on the adrenal gland.

Synthesis of cortisol in the adrenal cortex is induced by the binding of ACTH to its membrane receptor (melanocortin 2 receptor, MC2R), which activates adenylate cyclase, resulting in increased intracellular levels of cAMP, which in turn activates protein kinase. The activated protein kinase stimulates transcription of the steroid acute regulatory protein (STAR), which participates in the transport of cholesterol towards the inner mitochondrial membrane where the first enzymatic reaction of steroidogenesis occurs. This transfer of cholesterol from the outer to the inner mitochondrial membrane is the rate-limiting step in steroidogenesis (Manna *et al.*, 2009). Modulation of cortisol secretion by sex steroids would be exerted through their binding to a specific receptor in adrenocortical cells. Oestrogen receptors (ER) and androgen receptors (AR) have been found in the rat adrenal gland (ER: Cutler *et al.*, 1978; AR: Calandra *et al.*, 1978) and in the sheep adrenal gland (ER: Van Lier *et al.*, 2003b; AR: Gay *et al.*, 2007). Sex steroids do affect adrenocortical sensitivity to ACTH in rats (Atkinson and Waddell, 1997), but no reports in ruminants were found. We hypothesise that steroid treatment modulates adrenal response – in terms of cortisol secretion – to an exogenous ACTH treatment, and that the differences between cortisol secretion in male and female sheep are partly due to sex steroids (oestrogens and testosterone) acting directly on the gland. Furthermore, the sex steroids might affect the sensitivity of the adrenal gland to ACTH by modulating the concentration of MC2R, or they might affect precursor availability for cortisol production by modulating STAR abundance (Van Lier *et al.*, 2003b). Thus, in the first study we compared the cortisol response to ACTH in gonadectomised sheep with or without sex steroid replacement. In the second study we measured the relative expression of *ERS1*, *AR*, *MC2R* and *STAR* mRNAs in adrenal glands of gonadectomised male and female sheep either with or without sex steroid replacement (Testosterone Cyclopentilpropionate (TC) and Oestradiol Benzoate (EB), respectively).

Material and methods

Working conditions

All of the sheep were of the Corriedale breed and were accustomed to frequent handling. Before the experiments the animals were kept at pasture with free access to water. The experiments were carried out in autumn (May, breeding season, Southern Hemisphere). Both experimental protocols were approved by the Honorary Committee of Animal Experimentation of the University of Uruguay (Comisión Honoraria de Experimentación Animal, CHEA, protocols No. 7101/05 and 7101/07).

Experiment 1

Six adult rams (mean BW \pm s.e.m.: 70.6 \pm 1.52 kg) and four adult ewes (mean BW \pm s.e.m.: 42.5 \pm 1.53 kg) were used. Gonadectomy was performed 4 weeks before the experiment. On the day before each of two frequent blood-sampling periods, each animal's neck was shorn and fitted with an indwelling jugular catheter for the collection of blood samples into heparinised tubes. The samplings took place in a shed with the animals in small pens in pairs of the same sex. The animals were all in visual contact with each other and were fed with hay and had free access to water. During these periods, blood samples were taken every 15 min for 4 h.

On the day of the first frequent blood sampling each animal received 0.5 mg of an ACTH analogue (a polypeptide based on the first 24 amino acids of the 39 occurring in the natural molecule: Tetracosactid, Synacthen Depot[®], 1 mg/ml; Ciba Geigy, Basel, Switzerland) *i.v.*, immediately after the sample at 1 h from the beginning of the trial, which was considered time 0 (Van Lier *et al.*, 2003a). Four days later the males received 100 mg of TC *i.m.* (in oil vehicle, 100 mg/ml, Laboratorios Dispert S.A., Montevideo, Uruguay; recommended dose for androgenisation of wethers for oestrus detection) and the females received 2.5 mg of EB *i.m.* (in oil vehicle, 5 mg/ml, Laboratorios Dispert S.A.; half of the minimum therapeutic dose recommended). One week after the first frequent blood sampling a second identical sampling was performed (72 h after TC or EB administration) with ACTH administration at 0 h (1 h after the beginning of the trial). Sampling began at 1000 h on both occasions. In order to establish the effectiveness of the TC and EB treatment, samples for analysis of testosterone or 17 β -oestradiol were taken at 0, 24, 48 and 72 h relative to steroid replacement.

Experiment 2

Eight adult rams (mean BW \pm s.e.m.: 70.6 \pm 2.80 kg) and eight adult ewes (mean BW \pm s.e.m.: 43.1 \pm 2.70 kg) were used. Gonadectomy was performed at least 3 weeks before the experiment. Four rams received 100 mg of TC *i.m.* (in oil vehicle, 100 mg/ml) and four females received 0.5 mg of EB *i.m.* (in oil vehicle, 5 mg/ml; 1/10 of the minimum therapeutic dose recommended). Blood samples were taken at 0, 12, 24, 48 and 72 h relative to steroid replacement in order to establish the effectiveness of the TC and EB treatments and the animals were thereafter slaughtered in order to obtain

their adrenal glands. Slaughter was performed under barbiturate general anaesthesia (1 g of thiopental sodium) and the adrenal glands were sampled within 15 min. Adrenal glands were weighed and then cut in half on ice. The tissues were packed and immediately frozen in liquid nitrogen and then stored at -80°C .

Hormone assays

Plasma was obtained by centrifugation of the samples immediately after collection. Approximately 3 ml of plasma was taken from each sample and then stored at -20°C within 2 h after collection. All hormone concentrations were determined using previously validated radioimmunoassays (Coat-A-Count radioimmunoassay kits; Diagnostic Products Corporation, Los Angeles, CA, USA).

Cortisol was analysed in every half an hour sample for the first hour and then in every 15 min sample for 2 h and then each half hour until the end of experiment 1. All of the samples were analysed in the same assay. Quality control samples containing endogenous cortisol (low 36.9 nmol/l, medium 154.9 nmol/l and high 296.0 nmol/l) were assayed in duplicates at both the beginning and end of the assay. The intra-assay CV were 8.6%, 3.7% and 5.2%, respectively. The analytical detection limit of the assay was 10.8 nmol/l.

Testosterone was determined in the samples of the rams in order to evaluate the effectiveness of the replacement treatment. For experiment 1 the samples at 0, 24 and 48 h relative to steroid replacement, as well as one sample before and after ACTH administration were analysed, and for experiment 2 all of the samples of the rams were analysed. The intra-assay CV for two control samples (4.0 and 24.1 nmol/l), assayed in duplicates at the beginning and end of each assay, were 9.9% and 6.1%, respectively. The corresponding inter-assay CV were 12.9% and 5.7%, respectively. The analytical detection limit of the assay was 0.4 nmol/l.

In the samples of the ewes 17β -oestradiol was determined in order to evaluate the effectiveness of the replacement treatment. For experiment 1 the samples at 0, 24 and 48 h relative to steroid replacement, as well as one sample before and after ACTH administration were analysed, and for experiment 2 all of the samples of the ewes were analysed. The samples of each experiment were analysed in the same assay. The intra-assay CV for two control samples (low 14.4 pmol/l and medium 45.3 pmol/l) were 23.6% and 6.2%, respectively. The analytical detection limit of the assay was 2.6 pmol/l.

Real-time reverse transcription PCR (RT-PCR)

In order to quantify mRNA expression of *ERS1*, *AR*, *MC2R* and *STAR* using real-time RT-PCR, total RNA was extracted using TRIzol (Invitrogen, Life Technologies, Carlsbad, CA, USA), precipitated with lithium chloride to remove cDNA synthesis inhibitors, and treated with DNase to remove contaminating DNA (DNA-Free kit; Applied Biosystems/Ambion, Austin, TX, USA). The concentration of RNA was determined by measuring absorbance at 260 nm, the purity of all RNA isolates was assessed from 260/280 absorbance

ratio and the integrity by electrophoresis (1% agarose gel). All samples presented $A_{260/280}$ ratios between 1.8 and 2.0.

For each sample, cDNA was synthesised by reverse transcription using the SuperScript III First-Strand Synthesis System Kit (Invitrogen) with random primers and 1 μg of total RNA added as a template. Primers to specifically amplify cDNA of *MC2R*, *ERS1* and *STAR*, as well as the housekeeping genes *hypoxanthine phosphoribosyltransferase (HPRT)* and *ribosomal protein L19 (RPL19)* were obtained from the literature (Schams *et al.*, 2003; Myers *et al.*, 2005; Chen *et al.*, 2006; Jaquiere *et al.*, 2006; Carriquiry *et al.*, 2009) and further primer information is given in Supplementary Table S1. Primers to amplify cDNA of *AR* were designed based on ovine nucleotide sequences available from NCBI (<http://www.ncbi.nlm.nih.gov/>) using Primer Express 3.0 software from Applied Biosystems. Before use, primer product size (1% agarose gel separation) was determined to ensure that primers produced the desired amplification products (results not reported). Real-time PCR reactions were performed using 10 μl SYBR green (Quantimix; Biotools, Madrid, Spain) equal amounts (200 nM) of forward and reverse and 3 μl diluted cDNA (1 : 7.5 in RNase/DNase free water) in a final volume of 20 μl . Samples were analysed in duplicate in a 72-disk Rotor-GeneTM 6000 (Corbett Life Sciences, Sydney, Australia). Standard amplification conditions were 5 min at 95°C and 40 cycles of 10 s at 95°C , 45 s at 60°C and 20 s at 72°C . Each disk included a pool of total RNA from adrenal samples analysed in triplicate to be used as the basis for the comparative expression results (exogenous control) and duplicate tubes of water (non-template control). Gene expression was measured by relative quantification to the exogenous control and normalised to the mean expression (geometric mean) of the endogenous control genes (*HPRT* and *RPL19*). Dilution curves were developed for each primer from the quantification of 6.25, 12.5, 25, 100 and 200 ng of transcribed RNA and amplification efficiency was estimated by linear regression (Supplementary Table S1). The intra- and inter-assays CV were $<7.6\%$ and 13.1% , respectively.

Statistical analysis

Data were analysed using the SAS Systems programs (SAS 9.2 V; SAS Institute Inc., Cary, NC, USA). Cortisol data (experiment 1) were analysed as repeated measures, using the MIXED procedure. The model included the effect of sex (male or female) and steroid replacement (with or without) within sex, time (repeated measure within subject) and their interactions as fixed effects and animal (subject) as a random effect. The first-order autoregressive covariance structure (AR(1)) and the Kenward–Rogers procedure to adjust the degrees of freedom of denominator were specified. Gene expression data, animal BW, absolute and relative tissue weight (experiment 2) were analysed using the MIXED procedure with a model that included the fixed effects of sex (male or female) and steroid replacement (with or without) within sex. Mean separation was performed using the Tukey test. Correlation coefficients used to describe relationships

Table 1 The effect of steroid replacement (immediately after the 0 h sample) on mean concentrations of testosterone (nmol/l) in Testosterone Cyclopentilpropionate (TC)-treated castrated rams and concentrations of 17 β -oestradiol (pmol/l) in Oestradiol Benzoate (EB)-treated ovariectomised ewes either before ACTH administration (experiment 1) or before slaughter (experiment 2)

	0 h	12 h	24 h	48 h	72 h	SEp	P (time)
Testosterone¹							
Experiment 1	0.0	–	10.4	11.0	13.4	1.9	0.0013
Experiment 2	0.0	18.6	23.8	24.9	22.8	1.6	<0.0001
17β-oestradiol²							
Experiment 1	5	–	1692	49	237	185	0.0009
Experiment 2	13	275	288	252	118	40	0.0012

– = no sample; SEp = pooled standard error.

¹The same dose of TC (100 mg/ram) was used in both experiments. The analytical detection limit of the assay was 0.4 nmol/l.

²Different doses of EB per ewe were used in experiments 1 and 2 (2.5 and 0.5 mg of EB, respectively). The analytical detection limit of the assay was 2.6 pmol/l.

between gene expressions were estimated using the CORR procedure. The level of significance was considered to be $P \leq 0.05$, and tendency as $0.5 < P < 0.10$.

Results

Testosterone and 17 β -oestradiol

Testosterone was not detected in any of the samples of the non-treated castrated rams. Steroid replacement with TC effectively raised testosterone levels in all of the castrated rams (Table 1). Testosterone concentrations in the samples after ACTH administration were not different from the sample taken just before ACTH administration (data not shown). In all of the ovariectomised ewes 17 β -oestradiol was low, but detectable. EB administration increased 17 β -oestradiol levels in plasma of all of the ewes (Table 1). The concentrations of 17 β -oestradiol in the samples after ACTH administration were not different from the sample taken just before ACTH administration (data not shown).

Cortisol in ACTH trials

The main effects *sex* ($P < 0.0001$), *steroid replacement*(*sex*) ($P = 0.024$) and *time* (minutes relative to ACTH treatment) ($P < 0.0001$), were significant, as were the interactions *sex* \times *time* ($P < 0.0001$) and *time* \times *steroid replacement*(*sex*) ($P = 0.0073$). Before ACTH administration cortisol levels were low, while after ACTH administration they were significantly higher in all of the animals and did not reach pre-treatment levels before the end of the experiment (Figure 1). The rams had lower cortisol levels than the ewes at time 0 ($P = 0.0321$) and after ACTH administration ($P < 0.0002$), but TC did not affect these cortisol levels. In the EB-treated ewes, however, the cortisol levels after ACTH administration were higher as compared with non-treated ewes ($P = 0.0097$) (Figure 1).




Figure 1 The effect of sex and steroid replacement on mean (\pm s.e.m.) cortisol levels (nmol/l) before and after administration (i.v.) of an ACTH analogue at time 0 in gonadectomised sheep (dots are ewes and squares are rams; open symbols are non-steroid-treated sheep and solid symbols are steroid-treated sheep). The cortisol response was significantly different between rams and ewes and stars indicate significant differences between Oestradiol Benzoate treated and control ewes ($P < 0.05$).

BW, absolute and relative adrenal weight

BW was different between sexes ($P < 0.0001$), but *steroid replacement*(*sex*) had no effect ($P = 0.6777$) (Table 2). The absolute adrenal weight was not different between sexes ($P = 0.3811$) and *steroid replacement*(*sex*) had no effect ($P = 0.8304$) (Table 2). The relative adrenal weight was different between sexes ($P = 0.0002$), but *steroid replacement* (*sex*) had no effect ($P = 0.8877$) (Table 2).

MC2R, AR, ERS1 and STAR mRNA

The effect of *sex* on the relative amount of *MC2R* transcript was not significant while the effect of *steroid replacement* (*sex*) tended to be significant ($P = 0.0558$). Gonadectomised rams treated with TC presented greater *MC2R* mRNA expression than males without the steroid replacement and ewes treated with EB (Table 3). For the relative amount of the *AR* transcript the effect of *sex* tended to be significant ($P = 0.0844$) while the effect of *steroid replacement*(*sex*) was significant ($P = 0.0421$). Gonadectomised rams with TC had greater *AR* mRNA expression than non-replaced rams and ewes treated with EB. Gonadectomised ewes treated with EB tended to present lower *AR* mRNA than the ewes without steroid replacement ($P = 0.0881$) (Table 3). No significant effects were found on the relative amount of *ERS1* and *STAR* transcripts (Table 3). However, gonadectomised rams with TC had greater *ERS1* mRNA expression than ewes treated with EB. The transcripts of *AR* and *MC2R* were highly, positively and significantly correlated ($r = 0.95$, $P < 0.0001$) and *ERS1* was highly, positively and significantly correlated with *AR* ($r = 0.79$, $P = 0.0008$) and with *MC2R* ($r = 0.80$, $P = 0.0007$). The *STAR* transcript was also positively correlated with the receptor transcripts, although to a lesser extent (*ERS1*: $r = 0.60$, $P = 0.0244$; *AR*: $r = 0.58$, $P = 0.0302$; and *MC2R*: $r = 0.52$, $P = 0.0548$). In rams TC was positively correlated ($r = 0.75$) with the *AR* and *MC2R* transcripts

Table 2 The effect of sex and steroid replacement on mean BW (kg), absolute adrenal weight (g) and relative adrenal weight (mg adrenal tissue/kg BW) in gonadectomised ewes (control and replaced with Oestradiol Benzoate, EB) and gonadectomised rams (control and replaced with Testosterone Cyclopentilpropionate, TC) (experiment 2)

	Control ewes	EB ewes	Control rams	TC rams	SEp	<i>P</i> (sex)	<i>P</i> (replacement(sex))
BW	44.8 ^a	41.5 ^a	72.6 ^b	68.6 ^b	5.8	<0.0001	0.6777
Absolute adrenal weight	3.58	3.26	3.74	3.77	0.52	0.3811	0.8304
Relative adrenal weight	79.5 ^a	78.0 ^a	52.1 ^b	55.0 ^b	6.7	0.0002	0.8877

SEp = pooled standard error.

^{a,b}Values within a row with different superscripts differ significantly at *P* < 0.05.

Table 3 The effect of sex and steroid replacement on relative expression of MC2R, AR, ERS1 and STAR mRNA in gonadectomised ewes (control and replaced with Oestradiol Benzoate, EB) and gonadectomised rams (control and replaced with Testosterone Cyclopentilpropionate, TC) (experiment 2)

Transcript	Control ewes	EB ewes	Control rams	TC rams	SEp	<i>P</i> (sex)	<i>P</i> (replacement(sex))
MC2R	1.53 ^{ab}	1.13 ^a	1.35 ^a	1.95 ^b	0.26	0.1172	0.0558
AR	1.37 ^{ab†}	0.92 ^{a†}	1.19 ^a	1.73 ^b	0.24	0.0844	0.0421
ERS1	1.44 ^{ab}	1.07 ^a	1.29 ^{ab}	1.63 ^b	0.21	0.2042	0.1127
STAR	1.43	1.06	1.54	1.37	0.32	0.3900	0.4972

AR = androgen receptor; ERS1 = estrogen receptor α ; MC2R = melanocortin 2 receptor (ACTH receptor); STAR = steroid acute regulatory protein; SEp = pooled standard error.

^{a,b}Values within a row with different superscripts differ significantly at *P* < 0.05, [†]*P* = 0.0881.

(*P* = 0.0538 and 0.0522, respectively), and in ewes EB was negatively correlated (*r* = -0.78) with the AR transcript (*P* = 0.0384).

Discussion

Steroid replacement

Before steroid replacement testosterone was not detectable in the blood samples of the rams indicating successful castration. However, the ovariectomised ewes had low levels of 17 β -oestradiol, which has been observed (Adams *et al.*, 1990; Van Lier *et al.*, 2003b) and suggests an extragonadal source of 17 β -oestradiol (or its precursors), possibly the adrenal cortex. Both EB and TC were effective in increasing 17 β -oestradiol and testosterone concentrations, respectively, in all of the treated animals. The levels of 17 β -oestradiol in the animals of experiment 1 were pharmacological and the dose of EB was therefore reduced in experiment 2, resulting in lower, yet still pharmacological levels of 17 β -oestradiol in those animals. Testosterone levels of the animals in both experiments were within the physiological range, although by intramuscular administration the active compound is usually released into the bloodstream at a relatively steady rate and not in discrete pulses, as occurs in intact rams (Sanford *et al.*, 1977).

Cortisol response to ACTH

The differences in cortisol response to ACTH in gonadectomised sex steroid replaced sheep confirm our previous findings of sex differences in cortisol secretion in intact sheep (Van Lier *et al.*, 2003a) and expand these findings as in the present study we found differences in cortisol levels of gonadectomised sheep. This is in contrast with earlier studies

(Tilbrook *et al.*, 1999; Turner *et al.*, 2002) where no difference in cortisol response was observed between gonadectomised male and female sheep. In our earlier work (Van Lier *et al.*, 2003a), cortisol levels of the gonadectomised rams and ewes were in between the levels of the intact animals and tended to be different (*P* = 0.0701). However, differences in cortisol secretion in gonadectomised sheep have been found (Broadbear *et al.*, 2005), which were not reverted by sex steroid replacement, although the ewes were replaced with luteal phase levels of progesterone and oestrogens, which means oestrogens were at nadir. In rats, gonadectomy reduces or eliminates sexual differences in HPA function (Nowak *et al.*, 1995). This discrepancy in the effect of gonadectomy on cortisol secretion in sheep suggests that other factors are involved in the cortisol response to ACTH administration. Sex steroids present during the perinatal period can organise neuronal substrates, resulting in lifelong alterations in endocrine function (McCormick *et al.*, 1998). This might also cause permanent 'organisational' differences in peripheral tissues such as the adrenal cortex, although it is not clear why the differences in cortisol secretion in gonadectomised male and female sheep are not always observed. On the other hand, life cycle stage of the animals also modulates cortisol secretion (Korte *et al.*, 2005) and might conceal differences in gonadectomised ewes and rams.

Hormone replacement affected the ACTH-induced cortisol secretion only in ewes. EB increased the cortisol response to ACTH in female sheep, confirming our hypothesis that oestrogens are involved in the regulation of the response of the adrenal cortex to ACTH, at least at the dose of EB used in our experiment. The effect of oestrogens on the secretion of cortisol has been described in rodents since the 1960s (Kitay, 1961) and later the presence of ER in the adrenal cortex of rats and primates was demonstrated (Cutler *et al.*, 1978;

Hirst *et al.*, 1992). Since then, ER has also been described in the sheep adrenal gland (Van Lier *et al.*, 2003b), supporting a possible role for oestrogens as modulators of cortisol secretion.

Although in our experiment TC did not modify the cortisol response to ACTH in the rams, it cannot be ruled out that this might be due to the type of steroid replacement leading to sustained testosterone levels and not discrete pulses (Sanford *et al.*, 1977). However, Broadbear *et al.* (2005) did not detect differences in cortisol response after HPA stimulation of castrated rams with or without testosterone implant replacement (mean testosterone concentrations 6.11 nmol/l; lower than in the present study). Androgen receptors have been found in adrenal glands from rats (Calandra *et al.*, 1978; Bentvelsen *et al.*, 1996), monkeys (Hirst *et al.*, 1992) and sheep (Gay *et al.*, 2007), indicating that androgens can act on the tissue. Van Lier *et al.* (2003a) reported that while ovariectomised ewes had lower cortisol levels than the intact ewes, castration of rams did not affect cortisol secretion compared with the intact rams, suggesting that cortisol secretion may be regulated by oestrogens – but not testosterone – at the adrenal cortex. However, in other species, androgens have been reported to inhibit corticosteroid secretion (rodents: Handa *et al.*, 1994; Ogilvie and Rivier, 1997; McCormick *et al.*, 1998; cattle: Verkerk and Mcmillan, 1997), probably through action on other parts of the HPA axis.

Adrenal weight

The absolute adrenal weight did not show sex differences and confirms our earlier observations (Van Lier *et al.*, 2003b). However, when the adrenal weight is related to the live weight, ewes have more adrenal mass per unit of live weight than the rams. Absolute and relative adrenal weights seem not to be affected by gonadectomy (Canny *et al.*, 1999; Van Lier *et al.*, 2003b), which would be confirmed by our results since treatment with EB or TC were without effect. Female rats had 30% more adrenal mass than male rats, and this difference appeared after sexual maturation (Rifka *et al.*, 1978). In that study gonadectomised hypophysectomised rats were used and oestrogens did not affect the adrenal weight of female rats, while testosterone induced a decrease in the adrenal weight of male rats. However, Bentvelsen *et al.* (1996) did not find differences in adrenal weight of castrated androgen-replaced male rats. In sheep it seems that the sex steroids do not have trophic effects on the adrenal glands. It is not clear when the differences in relative adrenal weight appear in sheep: before birth (as an organisational effect of sex) or after puberty, as in rats, due to gonadal steroid modulation.

Adrenal ERS1, AR, MC2R and STAR mRNA

This study confirms earlier observations of ERS1 presence in the sheep adrenal gland (Van Lier *et al.*, 2003b) and confirms other reports on the presence of AR (Gay *et al.*, 2007). No differences were found in AR or ERS1 mRNA expression in gonadectomised rams and ewes. Similarly, Van Lier *et al.* (2003b) did not find differences in ERS1

transcript (measured by solution hybridisation) in the adrenals of intact and gonadectomised ewes and rams, while the ER content (measured by ligand-binding assay) differed between males and females and intact and gonadectomised sheep. Thus, data suggest that steady state levels of receptor protein are regulated at other levels of protein synthesis (e.g. efficiency of translation).

Sex steroid replacement was successful in inducing differences in adrenal ERS1 and AR mRNA expression, confirming a role for the sex steroids in regulating tissue sensitivity to hormones (e.g. hormone receptors) in the adrenal gland as has been reported for other tissues (Spencer and Bazer, 1995). EB replacement tended to reduce the relative expression of AR mRNA in ewes and TC increased the relative expression of AR mRNA in rams. The increase of AR mRNA expression in the adrenal gland by androgens has also been observed in the male rat (Bentvelsen *et al.*, 1996). A down regulation of steroid receptors (ER and progesterone receptor-binding activity) by oestrogen treatment has been reported in sheep uteri (Meikle *et al.*, 2000), and the authors suggested that this down regulation is a mechanism to limit hormone action in the target cells. An oestrogen down regulation of ERS1 expression was also found in rat uteri, and the regulation is dose dependent, as demonstrated by the stimulating or inhibitory effects of low or high doses of oestrogens, respectively (Medlock *et al.*, 1994).

EB increased the cortisol response to ACTH in female gonadectomised sheep, but no effect of TC was found in male sheep. Several mechanisms have been suggested by which sex steroids might affect adrenal steroidogenesis such as adrenocortical sensitivity, steroid precursor availability or enzyme induction (Van Lier *et al.*, 2003b). In this study we have looked into the first two of these mechanisms. The most straightforward mechanism would be for the sex steroids to affect adrenocortical sensitivity to ACTH (Atkinson and Waddell, 1997). A stimulating action of oestrogens and an inhibiting action of testosterone on the adrenal have been previously proposed in rats (Kitay, 1963; Nowak *et al.*, 1995). Therefore, one of our target genes in experiment 2 was of the ACTH receptor (MC2R). Contrary to what we expected to find based on the cortisol results of experiment 1 (increased after EB and no change after TC), we found no differences in MC2R expression in EB-treated ewes and higher MC2R expression in TC-treated rams. However, adrenal sensitivity depends on actual MC2R protein content and even if mRNA is frequently used as a predictor of protein content it may not always present a direct relationship as has been described before (Meikle *et al.*, 2000), thus, post-transcriptional events might affect receptor activation and/or receptor turnover. Although the time between steroid replacement and adrenal sampling was determined to allow for protein synthesis and has been used in other studies (Clark *et al.*, 1992; Meikle *et al.*, 1997, 2000), it cannot be ruled out that a critical window was missed.

Steroidogenesis could also be modulated by influencing steroid precursor (cholesterol) availability to the mitochondrial enzymes. The STAR protein participates in the transport of

cholesterol towards the inner mitochondrial membrane where the first enzymatic reaction of steroidogenesis occurs. This transfer of cholesterol from the outer to the inner mitochondrial membrane is the rate-limiting step in steroidogenesis (Manna *et al.*, 2009). Both *STAR* gene expression and transcription are regulated by protein kinase activation in response to ACTH stimulation of cAMP (Dyson *et al.*, 2009). In our study, we have found no evidence that sex or sex steroids differentially regulate *STAR* expression, since no difference in relative abundance of *STAR* mRNA was observed. However, the amount of *STAR* protein is primarily regulated by transcription, which then also regulates steroidogenesis (Hiroi *et al.*, 2004). Therefore, the possibility exists that the observed sex differences in cortisol secretion post-ACTH administration might be due to differences in *STAR* translation rate (caused by the sex steroids) and thus its availability either limiting or stimulating steroidogenesis.

The high and positive correlation between the transcripts is indicative of a regulatory role of the sex steroid on the adrenal gland. TC showed a high and positive correlation with the transcripts of *AR* and *MC2R*; EB showed a high and negative correlation with the transcript of *AR*. This contrast between a positive correlation between TC and *AR* and a negative one for EB and *AR* further supports the concept of opposite adrenal regulation of testosterone and 17β -oestradiol. However, while the sheep adrenal gland expresses the receptor proteins for both 17β -oestradiol and testosterone, and therefore may be subject to sex steroid modulation, the underlying mechanisms remain unresolved.

Conclusions

Gonadectomised steroid replaced ewes and rams in this study showed the most divergent cortisol profiles as well as the most divergent relative *ERS1*, *AR* and *MC2R* mRNA expression, confirming a modulatory role of the sex steroids on cortisol secretion acting directly on the adrenal glands. In ewes, 17β -oestradiol modulated cortisol secretion by increasing the response to exogenous ACTH administration, but only tended to decrease the relative adrenal *AR* expression in ovariectomised ewes. A role for testosterone in influencing cortisol secretion in castrated rams was not confirmed, however, it did increase the relative adrenal *MC2R* and *AR* expression compared with non-steroid replaced castrated rams.

Acknowledgements

The authors thank Nelson Villegas, Daniel Agüero, Dana Montedónico, Alejandra Carbone, Angélica de Larrobla, Carolina Chiesa, Hugo Petrocelli, Stella Lanzzeri, Andrea Fernández, Beatriz Carracelas, Alejandro Castillo, Carlos Pérez, Andrés López for their help with the field work, Oscar Bentancur for statistical assistance, and Yerú Pardiñas and Enrique Rodríguez Grasso for donating part of the animals. Elize van Lier received a grant from the Uruguayan Ministry of Education and Culture, DINACYT, Fondo Profesor Clemente Estable (10083), Uruguay.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S1751731114000780>

References

- Adams NR, Atkinson HC, Hoskinson RM, Abordi JA, Briegel JR, Jones M and Sanders MR 1990. Immunization of ovariectomized ewes against progesterone, oestrogen or cortisol to detect effects of adrenal steroids on reproduction. *Journal of Reproduction and Fertility* 89, 477–483.
- Atkinson HC and Waddell BJ 1997. Circadian variation in basal plasma corticosterone and adrenocorticotropin in the rat: sexual dimorphism and changes across the estrous cycle. *Endocrinology* 138, 3842–3848.
- Bentvelsen FM, McPhaul MJ, Wilson CM, Wilson JD and George FW 1996. Regulation of immunoreactive androgen receptor in the adrenal gland of the adult rat. *Endocrinology* 137, 2659–2663.
- Broadbear JH, Pierce BN, Clarke IJ and Canny BJ 2005. Role of sex and sex steroids in mediating pituitary-adrenal responses to acute buspirone treatment in sheep. *Journal of Neuroendocrinology* 17, 804–810.
- Burgess LH and Handa RJ 1992. Chronic estrogen-induced alterations in adrenocorticotropin and corticosterone secretion, and glucocorticoid receptor-mediated functions in female rats. *Endocrinology* 131, 1261–1269.
- Calandra RS, Purvis K, Naess O, Attramadal A, Djøseland O and Hansson V 1978. Androgen receptors in the rat adrenal gland. *Journal of Steroid Biochemistry* 9, 1009–1015.
- Canny BJ, O'Farrell KA, Clarke IJ and Tilbrook AJ 1999. The influence of sex and gonadectomy on the hypothalamo-pituitary-adrenal axis of the sheep. *Journal of Endocrinology* 162, 215–225.
- Carriquiry M, Weber WJ, Fahrenkrug SC and Crooker BA 2009. Hepatic gene expression in multiparous Holstein cows treated with bovine somatotropin and fed n-3 fatty acids in early lactation. *Journal of Dairy Science* 92, 4889–4900.
- Chen Y, Green JA, Antoniou E, Ealy AD, Mathialagan N, Walker AM, Avalle MP, Rosenfeld CS, Hearne LB and Roberts RM 2006. Effect of interferon-tau administration on endometrium of nonpregnant ewes: a comparison with pregnant ewes. *Endocrinology* 147, 2127–2137.
- Clark JH, Schrader WT and O'Malley BW 1992. Mechanisms of steroid hormone action. In *Textbook of endocrinology* (ed. JD Wilson and DW Foster), pp. 35–90. WB Saunders, Philadelphia, PA, USA.
- Cutler GB Jr, Barnes KM, Sauer MA and Loriaux DL 1978. Estrogen receptor in rat adrenal gland. *Endocrinology* 102, 252–257.
- Dyson MT, Kowalewski MP, Manna PR and Stocco DM 2009. The differential regulation of steroidogenic acute regulatory protein-mediated steroidogenesis by type I and type II PKA in MA-10 cells. *Molecular and Cellular Endocrinology* 300, 94–103.
- Figueiredo HF, Ulrich-Lai YM, Choi DC and Herman JP 2007. Estrogen potentiates adrenocortical responses to stress in female rats. *American Journal of Physiology Endocrinology and Metabolism* 292, E1173–E1182.
- Gay E, Campbell S, McNeilly A and Duncan WC 2007. Sexually dimorphic expression of androgen receptors in the fetal sheep adrenal gland. *Biology of Reproduction* 77, 119–120.
- Handa RJ, Burgess LH, Kerr JE and O'Keefe JA 1994. Gonadal steroid hormone receptors and sex differences in the hypothalamo-pituitary-adrenal axis. *Hormones and Behavior* 28, 464–476.
- Hiroi H, Christenson LK and Strauss JF III 2004. Regulation of transcription of the steroidogenic acute regulatory protein (StAR) gene: temporal and spatial changes in transcription factor binding and histone modification. *Molecular and Cellular Endocrinology* 215, 119–126.
- Hirst JJ, West NB, Brenner RM and Novy MJ 1992. Steroid hormone receptors in the adrenal glands of fetal and adult Rhesus monkeys. *Journal of Clinical Endocrinology and Metabolism* 75, 308–314.
- Jaquiere AL, Oliver MH, Bloomfield FH, Connor KL, Challis JRG and Harding JE 2006. Fetal exposure to excess glucocorticoid is unlikely to explain the effects of periconceptual undernutrition in sheep. *Journal of Physiology* 572, 109–118.
- Kitay JI 1961. Sex differences in adrenal cortical secretion in the rat. *Endocrinology* 68, 818–824.

- Kitay JI 1963. Pituitary-adrenal function in the rat after gonadectomy and gonadal hormone replacement. *Endocrinology* 73, 253–260.
- Korte SM, Koolhaas JM, Wingfield JC and McEwen BS 2005. The Darwinian concept of stress: benefits of allostasis and costs of allostatic load and the trade-offs in health and disease. *Neuroscience and Biobehavioral Reviews* 29, 3–38.
- Manna P, Dyson M and Stocco D 2009. Regulation of the steroidogenic acute regulatory protein gene expression: present and future perspectives. *Molecular Human Reproduction* 15, 321–333.
- McCormick CM, Furey BF, Child M, Sawyer MJ and Donohue SM 1998. Neonatal sex hormones have 'organizational' effects on the hypothalamic-pituitary-adrenal axis of male rats. *Developmental Brain Research* 105, 295–307.
- Medlock KL, Forrester MT and Sheehan DM 1994. Progesterone and estradiol interaction in the regulation of rat uterine weight and estrogen receptor concentrations. *Proceedings of the Society for Experimental Biology and Medicine* 205, 145–153.
- Meikle A, Tasende C, Rodríguez M and Garófalo EG 1997. Effects of oestradiol and progesterone on the reproductive tract and on uterine sex steroid receptors in female lambs. *Theriogenology* 48, 1105–1113.
- Meikle A, Forsberg M, Sahlin L, Masironi B, Tasende C, Rodriguez-Pinon M and Garófalo E 2000. A biphasic action of estradiol on estrogen and progesterone receptor expression in the lamb uterus. *Reproduction Nutrition Development* 40, 283–294.
- Myers DA, Hyatt K, Mlynarczyk M, Bird IM and Ducsay CA 2005. Long-term hypoxia represses the expression of key genes regulating cortisol biosynthesis in the near-term ovine fetus. *American Journal of Physiology Regulatory Integrative and Comparative Physiology* 289, R1707–R1714.
- Nowak KW, Neri G, Nussdorfer GG and Malendowicz LK 1995. Effects of sex hormones on the steroidogenic activity of dispersed adrenocortical cells of the rat adrenal cortex. *Life Sciences* 57, 833–837.
- Ogilvie KM and Rivier C 1997. Gender difference in hypothalamic-pituitary-adrenal axis response to alcohol in the rat: activational role of gonadal steroids. *Brain Research* 766, 19–28.
- Rifka SM, Cutler GB Jr, Sauer MA and Loriaux DL 1978. Rat adrenal androgen receptor: a possible mediator of androgen-induced decrease in rat adrenal weight. *Endocrinology* 103, 1103–1110.
- Sanford LM, Palmer WM and Howland BE 1977. Changes in the profiles of serum LH, FSH and testosterone, and in mating performance and ejaculate volume in the ram during the ovine breeding season. *Journal of Animal Science* 45, 1382–1391.
- Schams D, Kohlenberg S, Amselgruber W, Berisha B, Pfaffl MW and Sinowatz F 2003. Expression and localisation of oestrogen and progesterone receptors in the bovine mammary gland during development, function and involution. *Journal of Endocrinology* 177, 305–317.
- Spencer TE and Bazer FW 1995. Temporal and spatial alterations in uterine estrogen receptor and progesterone receptor gene expression during the estrous cycle and early pregnancy in the ewe. *Biology of Reproduction* 53, 1527–1543.
- Tilbrook AJ, Canny BJ, Serapiglia MD, Ambrose TJ and Clarke IJ 1999. Suppression of the secretion of luteinizing hormone due to isolation/restraint stress in gonadectomised rams and ewes is influenced by sex steroids. *Journal of Endocrinology* 160, 469–481.
- Turner A, Canny BJ, Hobbs RJ, Bond JD, Clarke IJ and Tilbrook AJ 2002. Influence of sex and gonadal status of sheep on cortisol secretion in response to ACTH and on cortisol and LH secretion in response to stress: importance of different stressors. *Journal of Endocrinology* 173, 113–122.
- Uhart M, Chong RY, Oswald L, Lin P-I and Wand GS 2006. Gender differences in hypothalamic-pituitary-adrenal (HPA) axis reactivity. *Psychoneuroendocrinology* 31, 642–652.
- Van Lier E, Pérez-Clariget R and Forsberg M 2003a. Sex differences in cortisol secretion after administration of an ACTH analogue in sheep during the breeding and non-breeding season. *Animal Reproduction Science* 79, 81–92.
- Van Lier E, Meikle A, Bielli A, Akerberg S, Forsberg M and Sahlin L 2003b. Sex differences in oestrogen receptor levels in adrenal glands of sheep during the breeding season. *Domestic Animal Endocrinology* 25, 373–387.
- Verkerk GA and Mcmillan KL 1997. Adrenocortical responses to an adrenocorticotrophic hormone in bulls and steers. *Journal of Animal Science* 75, 2520–2525.
- Viau V and Meaney MJ 1991. Variations in the hypothalamic-pituitary-adrenal response to stress during the estrous cycle in the rat. *Endocrinology* 129, 2503–2511.
- Viau V and Meaney MJ 1996. The inhibitory effect of testosterone on hypothalamic-pituitary-adrenal responses to stress is mediated by the medial preoptic area. *Journal of Neuroscience* 16, 1866–1876.
- Young EA 1995. The role of gonadal steroids in hypothalamic-pituitary-adrenal axis regulation. *Critical Reviews in Neurobiology* 9, 371–381.