THE

BRITISH JOURNAL OF NUTRITION

DIRECTIONS TO CONTRIBUTORS

(REVISED NOVEMBER 1970)

Papers submitted for publication in the *British Journal of Nutrition* should be as concise as possible. Economy of space should not, however, be achieved by suppressing useful results. Authors are invited to preserve experimental results too expensive for publication but deemed of importance and to indicate in the paper submitted their willingness to make such results available to others.

Papers should be accompanied by a signed statement to the effect that the author accepts the conditions laid down in Directions to Contributors. Special attention is directed to the sections below about the preparation of the typescript, and care in this matter will hasten publication. The Editors will return any typescript that does not conform to these conditions.

Communications. Papers submitted for publication should be sent to Professor T. G. Taylor (*British Journal of Nutrition*), Department of Physiology & Biochemistry, The University, Southampton SO₉ 5NH.

General. Submission of a paper to the Editorial Board will be held to imply that it represents the results of original research not previously published; that it is not under consideration for publication elsewhere; and that if accepted for the *British Journal of Nutrition* it will not be published elsewhere in the same form, in English or any other language, without the consent of the Editorial Board.

Authors' names should be given without titles or degrees. Women are requested to give one forename in full to avoid confusion. The name and address of the laboratory or institution where the work was performed should be given. Any necessary descriptive material about the author, e.g. Beit Memorial Fellow, should appear in parentheses after the author's name or at the end of the paper, and not in the form of a footnote.

Typescripts should bear the name and address of the person to whom the proof of the paper is to be sent and should also give a shortened version of the paper's title, not exceeding forty-five letters and spaces in length, suitable for a running title in the published pages of the work.

Form of Papers Submitted for Publication. The onus of preparing a paper in a form suitable for sending to press lies in the first place with the author. Authors should consult a current issue in order to make themselves familiar with the practice of the British Journal of Nutrition as to typographical and other conventions, use of cross-headings, layout of tables, etc. Attention to these and other details (mentioned below) in the preparation of the typescript before it is sent to the Editors will shorten the time required for publication: the need for undue amounts of editorial revision caused by badly prepared typescript will lead to delay in publication. Papers on specialized aspects of the subject should be so presented as to make them intelligible, without undue difficulty, to the ordinary reader of the Journal. Sufficient information should be given to permit repetition of the published work by any competent reader of the Journal.

Papers should be in double-spaced typing on one side

of sheets of uniform size with large margins. A space of 5 cm should be left at the top of the first sheet. Top copies only should be submitted, packed flat. The paper should be written in English, the spelling being that of the Oxford English Dictionary, and should, in general, be divided into the following parts. (a) Synopsis: each paper must open with a synopsis not more than 5 % of the length of the following text. This synopsis should aim at giving a picture in miniature of the entire article. The past tense should be used in referring to the author's experimental work. The present tense may be used where reference to existing knowledge is necessary, or where the author is stating what is shown or concluded. The change of tense should clearly differentiate the author's contribution from what is already known. The sequence in the synopsis should be the same as that in the paper. It is desirable to divide the synopsis into a series of numbered paragraphs, giving, where relevant, the following information: a succinct account of the experimental work with essential facts about apparatus, chemicals, methods and animals; the results, singling out new information; the conclusions from the results. (b) Introductory paragraph: it is not now customary to introduce a paper with a full account of the relevant literature, but the introductory paragraph should help the reader by indicating briefly the nature of the question asked and the reasons for asking it. The answer obtained should be indicated if it is possible to do so shortly. (c) Experimental methods adopted: with chemical papers, the Experimental part will normally appear towards the end, but with other types of publication Methods should appear after the introduction. (d) Results: these should be given as concisely as possible, with the help of figures or tables. (e) Discussion: it is desirable that the presentation of the results and the discussion of their significance should be considered separately. (f) References: these should be given in the text thus: Osborne & Mendel (1916), (Culbertson & Thomas, 1934); where a paper to be cited has more than two authors, the names of all the authors should be given when reference is first made, e.g. (Osborne, Mendel & Ferry, 1919); subsequent citations should appear thus (Osborne et al. 1919). Where more than one paper by the same authors has appeared in one year the reference should be given as follows: Osborne & Mendel (1914a); Osborne & Mendel (1914b); or Osborne & Mendel (1914a, b); (Osborne & Mendel, 1914b, 1916; Culbertson & Thomas, 1934). Where the lists of authors of two papers appearing in the same year begin with the same name but differ subsequently, references after the first full reference should include sufficient names to differentiate between the two papers, e.g. Thomas, Smith et al. (1960), Thomas, Wilson et al. (1960). At the end of the paper, references should be given in alphabetical order according to the name of the first author of the publication quoted, names with prefixes being entered under the prefix, and should include the author's initials; the title of the paper (except that of a thesis) should not be included. Names and initials of authors of unpublished work should be given in the text and not included in the References. Titles of journals should be abbreviated as in the World List of Scientific Periodicals, 1963-5, 4th ed. London: Butterworths. Examples of such abbreviations will be found in the current numbers of the British Journal of Nutrition. References to books and monographs should include the town of publication and the name of the publisher, as well as the date of publication and the number of the edition to which reference is made. Thus:

Ahrens, E. H. Jr, Hirsch, J., Insull, W. Jr & Peterson, M. L. (1958). In *Chemistry of Lipids as Related to Atherosclerosis* p. 222 [I. H. Page, editor]. Springfield, Ill.: C. C. Thomas.

Culbertson, C. C. & Thomas, B. H. (1934). Rep. Iowa agric. Exp. Stn 1933-4, p. 51.

Doisy, E. A., Somogyi, M. & Shaffer, P. A. (1923). J. biol. Chem. 55, xxxi.

Fairey, N. H. (1938). Nature, Lond. 142, 1156.

King, H. (1941). J. chem. Soc. p. 338.

Osborne, T. B. & Mendel, L. B. (1914a). J. biol. Chem. 17, 325.

Osborne, T. B. & Mendel, L. B. (1914b). J. biol. Chem. 18, 1.

Osborne, T. B. & Mendel, L. B. (1916). *Biochem. J.* 10, 534. Osborne, T. B., Mendel, L. B. & Ferry, E. L. (1919). *J. biol. Chem.* 37, 233.

Starling, E. H. (1915). Principles of Human Physiology 2nd ed. London: J. and A. Churchill.

Wilson, J. (1965). Leber's disease. PhD Thesis, University of London.

Units. Results should be presented in metric units according to the Système International d'Unités (see *Metrication in Scientific Journals*: The Royal Society, 1968).

Energy measurements should be expressed in joules or calories, but important mean values and other important values should be given in both units, the second being added in parentheses (4·184 J = 1 cal).

Time. The 24 h clock, should be used, e.g. 15.00 hours.

Statistical Treatment of Results. In general, it is not necessary to publish all the individual results of replicated tests. A statement of the number, their mean value and some appropriate measure of their variability is usually sufficient.

The methods of analysis followed should be indicated, but statistical details, such as tables of analysis of variance, should be given only if they are relevant to the discussion. A statement that the difference between the means for two groups of values is statistically significant should include an indication of the level of significance attained.

Where means are quoted, normally the measure of variation most appropriate is the standard error of the mean; occasionally, as for instance where there is specific interest in the distribution of the individual values in the sample on which the mean is based, the standard deviation may be more appropriate. The measure adopted must be clearly stated in the text and in each table. Where the measure of variation is presented as a separate column \pm need not be repeated before each value.

Illustrations. As a rule, illustrations should be about twice the size of the finished block, each on a separate sheet, not larger overall than the sheets on which the paper itself is typed, and packed flat. Legends should be typed on a separate sheet and numbered corresponding to the relevant illustrations. Each illustration, with its legend, should be comprehensible without reference to the text. The approximate position of each should be indicated in the text thus: 'Fig. 1 near here'.

Diagrams and line-drawings, which must all be originals, should be drawn in indian ink on Bristol-board or on cartridge-, tracing- or faintly blue-lined paper. For general guidance in preparing diagrams, it is suggested that for a figure measuring 250 mm × 150 mm all lines, axes and curves should have a thickness of 0.4 mm, thus: -. In curves presenting experimental results, the determined points should be clearly shown, the symbols used being, in order of preference, $\bigcirc \bullet$, $\triangle \blacktriangle$, $\square \blacksquare$, $\times +$. For a 250 mm × 150 mm figure, the diameter of the circles, the vertical height of the squares and equilateral triangles, and the span of the crosses should be 3 mm. Curves and symbols should be drawn with a mechanical aid and not freehand, and should not extend beyond the experimental points. Scale marks on the axes should be on the inner side of each axis, 3 mm long, and should extend beyond the last experimental point. For plates, glossy photographs (unmarked by paper-clips) are required.

A flyleaf of tracing-paper should be firmly attached to each illustration. On it should be written in ink: (a) the title of the paper and names of the authors; (b) the figure or plate number; (c) the figures and lettering which are to appear on the finished block, in their correct positions relative to the illustration.

Tables. Tables should carry headings describing their content and should be comprehensible without reference to the text. The dimensions of the values, e.g. g/10 ml, should be given at the top of each column and not repeated on each line of the table. Tables should not normally be included in the body of the text, but should be typed on separate sheets. Tables should not be subdivided by ruled lines. Their approximate position in the text should be indicated thus: 'Table 1 near here'.

Chemical Formulas. These should be written as far as possible on a single horizontal line. With inorganic substances, formulas may be used, particularly in the Experimental part, at the discretion of the Editors. With salts, it must be stated whether or not the anhydrous material is used, e.g. anhydrous CuSO₄, or which of the different crystalline forms is meant, e.g. CuSO₄.5H₂O, CuSO₄.H₂O.

Description of Solutions. Solutions of common acids, bases and salts are preferably defined in terms of normality (N) or molarity (M), e.g. N-HCl; O·IM-NaH2PO4. The term' %' must be used in its correct sense, i.e. g/100 g of solution. 10 % HCl means 10 g of hydrogen chloride in 100 g of aqueous solution, and should never be used to indicate a tenfold dilution of laboratory concentrated hydrochloric acid. For 'per cent by volume', i.e. ml/100 ml, the term ' %(v/v)' may be used. To indicate that a given weight of substance is contained in 100 ml of solution, the term '% (w/v)' may be used.

Nomenclature of Vitamins. Most of the names for vitamins and related compounds that are accepted by the Editors are those recommended by the IUNS Committee on Nomenclature. These have been published in Nutr. Abstr. Rev., 1970, 40, 395.

Previous name Vitamin A₁ Retinene or retinal Vitamin A acid Vitamin A2 or 3-dehydroretinol Retinene 2 or 3-dehydroretinal Vitamin D₂ or calciferol Vitamin D₃ Vitamins E Vitamin K₁ or phylloquinone Vitamin K2 series Vitamin K, menadione or menaphthone Vitamin B₁, aneurin(e) or thiamine Vitamin B₂ or riboflavine Nicotinic acid or niacin Niacinamide or nicotinic acid amide Folic acid or folacin(e) Vitamin B₆, adermin or pyridoxol Pyridoxal Pyridoxamine Vitamin B₁₂ or cobalamin Vitamin B_{12a}, B_{12b} or hydroxocobamide Vitamin B₁₂₀ Inositol or meso-inositol Pantothenic acid Biotin Choline p-Aminobenzoic acid Vitamin C or L-ascorbic Ascorbic acid acid

L-Dehydroascorbic acid

Recommended name Retinol Retinaldehyde Retinoic acid Dehydroretinol

Dehydroretinaldehyde

Ergocalciferol Cholecalciferol See Generic descriptors Phylloquinone

*Menaguinones Menaphthone

Thiamin

Riboflavin Nicotinic acid Nicotinamide

Pteroylmonoglutamic acid Pyridoxine

Pyridoxal Pyridoxamine Cyanocobalamin Hydroxocobalamin

Nitritocobalamin myo-Inositol Pantothenic acid **Biotin** Choline p-Aminobenzoic acid

Dehydroascorbic acid

Generic descriptors. The terms vitamin A, vitamin C and vitamin D may still be used where appropriate, for example in phrases such as 'vitamin A deficiency', 'vitamin D activity'.

Vitamin E. The term vitamin E should be used as the descriptor for all tocol and tocotrienol derivatives exhibiting qualitatively the biological activity of α-tocopherol. The term tocopherols should be used as the generic descriptor for all methyl tocols. Thus, the term tocopherol is not synonymous with the term vitamin E.

Vitamin K. The term vitamin K should be used as the generic descriptor for 2-methyl-1,4-naphthoquinone (menaphthone) and all derivatives exhibiting qualitatively the biologalic activity of phylloquinone (phytylmenaquinone).

Niacin. The term niacin should be used as the generic descriptor for pyridine 3-carboxylic acid and derivatives exhibiting qualitatively the biological activity of nicotin-

Folic acids. The term folic acid may be used to designate the naturally occuring pteroylglutamic acids.

Vitamin B₆. The term vitamin B₆ should be used as the generic desciptor for all 2-methylpyridine derivatives exhibiting qualitatively the biological activity of pyridoxine.

Vitamin B₁₂. The term vitamin B₁₂ should be used as the generic descriptor for all corrinoids exhibiting qualitatively the biological activity of cyanocobalamin. The term corrinoids should be used as the generic descriptor for all compounds containing the corrin nucleus and thus chemically related to cyanocobalamin. The term cor**rinoid** is not synonymous with the term **vitamin** B_{12} .

Summation. For vitamin A and its provitamins, summation should be made in mg (μ g) retinol equivalent. (In mg (μ g) β -carotene equivalent for separate summation of provitamins.) For vitamin E, summations should be made in mg (μ g) α -tocopherol equivalent. Other members of vitamin families suggested for summation are:

Vitamin D Cholecalciferol equivalent Vitamin K Phylloquinone equivalent Vitamin B₁₂ Cyanocobalamin equivalent Folic acid Folic acid equivalent Niacin Nicotinic acid equivalent

Nomenclature of Fatty Acids. In the description of results obtained for the analysis of fatty acids by conventional gas-liquid chromatography, the shorthand designation proposed by Farquhar, J. W., Insull, W., Rosen, P., Stoffel, W. & Ahrens, E. H. (Nutrition Reviews, 1959, 17, Supplement) for individual fatty acids should be used in the text, tables and figures. Thus 18: I should be used to represent a fatty acid with 18 carbon atoms and 1 double bond; if the position and configuration of the double bond is unknown, this fatty acid should not be referred to as oleic acid. The shorthand designation should also be used in the synopsis but sentences should be constructed so that it is clear to the non-specialist reader that 18: 1 refers to a fatty acid; for example, '...resulted in an increase in the concentration of the fatty acid 18: 1 in the liver triglycerides...'. If the positions and configurations of the double bonds are known, and these are important to the discussion, then a fatty acid such as linoleic acid may be referred to as 18:2 \triangle 9-cis, 12-cis (positions of double bonds related to the α-carbon atom). However, when essential and related fatty acids derived from animal tissues are being considered, it is preferable to refer to fatty acids such as linoleic acid as 18:206 and arachidonic acid as 20:406 (position of double bonds related to the ω -carbon atom); it is assumed that the double bonds are methyleneinterrupted and are of the cis-configuration (see Holman,

Details of the nomenclature for these and other naturally occuring quinones should follow the Tentative Rules of the IUPAC-IUB Commission on Biochemical Nomenclature (see Biochem. J. 1967, 102, 15).

R. T. in Progress in the Chemistry of Fats and Other Lipids, 1966, Vol. 9, part 1, p. 3. Oxford: Pergamon Press). Groups of fatty acids that have a common chain length but vary in their double bond content or double bond position should be referred to, for example, as C_{20} fatty acids or C_{20} polyunsaturated fatty acids. Impure samples of fatty acids such as those used in the preparation of diets should be referred to, for example, as 'linoleic acid'.

Nomenclature of Enzymes. The nomenclature should be that of the Recommendations of the Commission on Enzymes of the International Union of Biochemistry, 1964 (Enzyme Nomenclature, 1965. Amsterdam: Elsevier Publishing Co.).

Nomenclature \mathbf{of} Micro-organisms. Scientific names of bacteria should be binominals, the generic name only with a capital, and should be underlined once (for italic) in the typescript. Names for new species or genera, or new combinations of generic and specific names, should be formed in accordance with the International Bacteriological Code published in J. Bact. 1948, 55, 287, and (in 1949) in Int. Congr. Microbiol. IV (1948), Copenhagen, p. 587. Wherever possible, the names of recognized species should be those used in Topley & Wilson's Principles of Bacteriology and Immunity (1964: 5th ed. London: Edward Arnold and Co.). However, where authors wish for good reasons to use some other name (e.g. one used in Bergey's Manual of Determinative Bacteriology, 1957: 7th ed. London: Baillière, Tindall and Cox), the Topley & Wilson name should be inserted in parentheses at the first citation thus: Serratia marcescens (Chromobacterium prodigiosum).

A name must be given in full at the first mention in a paper; in subsequent mention the generic name (e.g. Staphylococcus, Streptococcus), may be abbreviated, but the abbreviation must be unambiguous. Single letter abbreviations should, in general, be avoided (thus: Staph. aureus, Strep. pyogenes, not S. aureus, S. pyogenes). When the generic name is used to define a group, it should have a capital but should not be italicized; trivial names, or generic names used as adjectives, should not have capitals or be italicized. Examples of 'trivial names' are: staphylococci, streptococci, and meningococci or meningococcus (generic name is Neisseria). Staphylococcus and Streptococcus are generic names and the following passage illustrates the correct usage. 'This investigation is concerned with Salmonella enteritidis, because this Salmonella is important as a cause of disease in man and, because of experimental salmonella infections in the mouse, it is the most easily controlled. Tests with staphylococci, including Staphylococcus aureus, have not contributed to the understanding of susceptibility to staphylococcal infection. Thus, further tests with Staph. aureus have not been made.'

Microfungi should be designated as in Ainsworth & Bisby's A Dictionary of the Fungi (1954: 4th ed. Kew: Commonwealth Mycological Institute).

Other Nomenclature, Symbols and Abbreviations. Authors should follow current numbers of the *British Journal of Nutrition* in this connexion. The chemical nomenclature adopted is that of the Chemical Society (see *Handbook for Chemical Society Authors* (Special Publication no. 14, 1961, 2nd ed. London: The Chemi-

cal Society)). For nomenclature of amino acids, this Handbook (p. 186) and Br. J. Nutr. 1953, 7, 1 should be consulted. The symbols and abbreviations, other than units, are essentially those listed in British Standard 1991: 1967 Letter Symbols, Signs and Abbreviations: Part 1. General (incorporating amendment no. 1, issued 7 July 1970). Day should be abbreviated to d, for example 7 d; except for example, 'each day', '7th day' and 'day 1'.

When an element e.g. nitrogen, is referred to frequently, the first mention, both in the synopsis and the main text, should be in full, thereafter by its chemical symbol N. Well-known abbreviations for chemical substances may be used without explanation, thus: RNA for ribonucleic acid and DNA for deoxyribonucleic acid. Other substances that are mentioned frequently may also be abbreviated, the abbreviation being placed in parentheses at the first mention, thus: free fatty acids (FFA), thereafter FFA. Well-known terms which are used frequently may be abbreviated in a similar fashion (the abbreviation being in small capitals), thus: basal metabolic rate or basal metabolism (BMR); biological value (BV); dry matter (DM); metabolizable energy (ME). Abbreviations in tables must be defined in footnotes. Signs for footnotes should be used in the sequence: * † ‡ § || ¶, then ** etc.; (omit * or †, or both, from the sequence if they are used to indicate levels of significance).

Spectrophotometric terms and symbols are those proposed by the Society of Public Analysts and other Analytical Chemists (see Analyst, Lond. 1942, 67, 164). For mathematical notation and numerals, the rules laid down in Proc. R. Soc. A, 1909, 82, 14 should be followed. The attention of authors is particularly drawn to the following symbols: m(= milli) = 10^{-8} , μ (= micro) = 10^{-6} , n (= nano) = 10^{-9} and p (= pico) = 10^{-12} . Note also that ml (millilitres) should be used instead of c.c., μ m (micro metre) instead of μ (micron) and μ g (microgramme) instead of γ .

Numbers. Figures should be used with units, for example, 10 g, 7 d, 4 years (except when beginning a sentence, thus 'Four years ago...'); otherwise, words (except when 100 or more), thus: one man, ten ewes, ninety-nine flasks, three times (but with decimals, 2.5 times), 100 patients, 120 cows, 136 samples.

Proofs. Proofs are sent to authors in order that they may make sure that the paper has been correctly set up in type, and not that they may add new material. Otherwise, increased printing charges are inevitable. Excessive alteration may have to be disallowed. The symbols used to indicate corrections should be those laid down in *British Standard* 1219C: 1958.

Corrected proofs and typescripts should be returned without delay to Mrs B. Smyth, National Institute for Research in Dairying, Shinfield, Reading, RG2 9AT.

Contributors who reside outside Europe will receive, by airmail, proofs of their papers, but not the typescripts.

Reprints. Orders for reprints should be addressed to The University Press, Cambridge. Where at least one author of a paper is a member of The Nutrition Society, twenty-five reprints are supplied free of cost. Additional reprints may be purchased if the Press are notified on the appropriate form immediately the proof of the paper is received.