

Current dietary supplement use of Australian military veterans of Middle East operations

Jolieke C van der Pols^{1,2,*}, Jeeva Kanesarajah¹, Alison Bell³ and Chi-Wai Lui¹

¹The University of Queensland, School of Public Health, Herston, QLD 4006, Australia: ²Queensland University of Technology, School of Exercise and Nutrition Sciences, Kelvin Grove, QLD, Australia: ³Recover Injury Research Centre, Faculty of Health and Behavioural Sciences, The University of Queensland, Herston, QLD, Australia

Submitted 18 July 2016: Final revision received 26 May 2017: Accepted 19 June 2017: First published online 15 August 2017

Abstract

Objective: To assess patterns and levels of dietary supplement use among Australian Defence Forces, previously deployed to the Middle East Area of Operations.

Design: A cross-sectional study. Participants of a large survey self-completed questions about dietary supplement use, health status, personal and job-related characteristics, and lifestyle factors. Frequency of current use of supplements was assessed in three categories (bodybuilding, energy and weight loss).

Setting: Middle East Area of Operations post-deployment health survey.

Subjects: Current and ex-serving Australian Defence Force personnel (n 14 032) who deployed to the Middle East between 2001 and 2009.

Results: Bodybuilding supplements were used by 17.5% of participants, energy supplements by 24.5% and weight-loss supplements by 7.6%. Overall, 32.3% of participants used any of these supplements. Bodybuilding and energy supplements were more often used by men, younger persons and those in the Army, while weight-loss supplements were more commonly used by women and Navy personnel. Supplements in all three categories were more commonly used by persons in lower ranks, active service and combat roles. Users of bodybuilding supplements had healthier lifestyles and better health status, while users of energy and weight-loss supplements had less healthy lifestyles and poorer mental and physical health status. Overall, 11.7% of participants used supplements containing caffeine and 3.6% used a creatine-containing product.

Conclusions: Use of dietary supplements among Australian Defence Force personnel is common, and patterned by lifestyle factors and health status.

Keywords
Dietary supplements
Military personnel
Australia
Creatine
Caffeine
Weight loss
Mental health

Use of dietary supplements is common in many Western countries including Australia, where national data indicate that almost a third of adults use these products⁽¹⁾. However, the efficacy and safety of dietary supplements have long been debated⁽²⁾. Carbohydrate and/or protein drinks can offer benefits during an extended period of intensive exercise^(3,4), or iron and calcium supplements can be beneficial for persons with low dietary intakes of these nutrients⁽⁵⁾. Due to the high physical demands commonly experienced by military personnel, they choose to use dietary supplements to improve energy levels, performance, muscle strength and endurance, and to promote general health⁽⁶⁾. Overweight and obesity are also increasingly affecting military populations^(7,8), which may lead to use of supplements marketed for weight loss in this population group, to help them to meet the fitness standards required for employment in military services.

However, many of the available supplements offer no health benefits⁽⁹⁾ and some may cause harm when multiple supplements are used, when intake exceeds the recommended dose⁽¹⁰⁾ or when their use causes a person's total intake to exceed recommended levels. For example, caffeine and creatine supplements can have ergogenic and performance-enhancing effects^(11,12), but caffeine-containing supplements add to the caffeine intake from drinks and caffeine intake levels in US military personnel have been reported to be high⁽¹³⁾. Some studies have shown possible risks associated with high caffeine intake, including a negative effect on sleep patterns⁽¹⁴⁾, although the effects of high caffeine intake on sleep quality in a combat environment have not been established definitively⁽¹⁵⁾. Certain weight-loss related supplements have been associated with gastrointestinal distress and nervousness^(16,17).

*Corresponding author: Email j.vanderpols@qut.edu.au

Current evidence indicates that use of dietary supplements by military personnel is higher than that in the general population in Western countries. More than 50% of US soldiers⁽⁶⁾ and almost 40% of British soldiers in training⁽¹⁸⁾ reported use of dietary supplements, with many using multiple products; and use is higher in army personnel serving in conflict zones^(19,20). Products targeting muscle strength, muscle mass and endurance are favoured by British and US military personnel^(6,19,21–23), and weight-loss supplements are also commonly used by US military personnel⁽²⁴⁾.

Most of the evidence on use of dietary supplements by military personnel comes from data collected on US and British military personnel, but evidence for Australian military personnel, who can take typical multivitamin preparations approved by the Australian Therapeutic Goods Administration at recommended doses without prior approval, is lacking. We therefore investigated levels and patterns of use of dietary supplements by Australian military personnel, particularly those who have experienced deployment to conflict zones. We studied the characteristics of dietary supplement users by assessing general demographic factors, service and lifestyle factors, and mental and physical health status.

Methods

Study design and participants

We analysed data from a large cross-sectional survey of current and ex-serving Australian Defence Force (ADF) members who deployed to the Middle East Area of Operations (MEAO) between 1 October 2001 and 31 December 2009⁽²⁵⁾. This survey included MEAO veterans deployed to Iraq or areas supporting operations in Iraq (including ships in the Persian Gulf), those deployed to Afghanistan or areas supporting operations in Afghanistan, and those attached to foreign militaries or the UN in the above areas. A small proportion (6%) of participants had left the ADF since serving in the Middle East. These ex-serving personnel were included in the present study to capture current supplement use of all persons who had served in the Middle East previously and improve representation of all those who served.

Data collection

Data were collected in 2010 and 2011 through online surveys or hard copy questionnaires that were sent to all eligible current and former ADF members for completion. Participation was voluntary and anonymous. Full details on the participant recruitment, response rates, survey design and methodology are documented elsewhere⁽²⁵⁾. In brief, all participants completed a questionnaire that asked about demographic characteristics, their role during the most recent deployment, current health status, lifestyle behaviours and health symptoms. Participants were asked the

following questions in relation to their supplement use: 'How often do you currently take any of the following supplements? Body building supplements (such as amino acids, weight gain products, creatine etc.); Energy supplements (such as energy drinks, pills, or energy enhancing herbs); Weight loss supplements?', with response options of 'never', 'less than once a month', 'monthly', 'weekly' and 'daily or almost daily'. The respondent was then asked to record the name (generic or brand name) of the supplement they used, for each of the three categories.

The supplement brand name and type listed by the participant were entered into an Internet search engine to verify the product was a dietary supplement and to obtain product information. Products in each of the three supplement categories (bodybuilding, energy and weight loss) were coded into sub-categories based on the most prominent type of ingredients, including protein and amino acids, multivitamins and herbs, carbohydrates and sugars, stimulants, and creatine monohydrate. We further identified which products contained either creatine or caffeine in each sub-category, given the concern about possible side-effects associated with these ingredients. Manual coding of the data was undertaken by one author and incongruities were resolved through discussion with all authors.

Lifetime smoking history was self-reported in three categories (current smoker; ex-smoker; never smoker) and caffeine use was ascertained as the reported average daily consumption of caffeine-containing beverages (250–375 ml) grouped into four categories (none; 1–2 per day; 3–5 per day; 6 or more per day). Participants were asked to report whether they experienced selected health symptoms (e.g. sleeping difficulties, headaches, vomiting) in the past month. The Short Form-12 Survey (SF-12)⁽²⁶⁾ was completed as part of the study questionnaire, and summary component scores for physical and mental health calculated using standardised methodology. Major depressive syndrome was derived using the Patient Health Questionnaire (PHQ-9) Depression module, which evaluates the frequency of nine DSM-IV (the Diagnostic and Statistical Manual of Mental Disorders, 4th edition) criteria for major depression syndrome in the last two weeks⁽²⁷⁾. Major depression was considered present if five or more of the nine depressive symptoms were reported more than half the time in the past two weeks, and if one of the symptoms reported was depressed mood or anhedonia.

Ethical approval for the present study was obtained from the Australian Department of Defence (protocol number LREP 14-010) and The University of Queensland (protocol number 2009001441) ethics committees. Participants of the MEAO Survey provided consent in writing or electronically, and all aspects of this research complied with the Declaration of Helsinki.

Statistical analysis

We assessed the proportion of respondents who reported using supplements in each of the three supplement

categories by subgroups of general demographic characteristics, service and lifestyle characteristics, and the presence or absence in the past month of a number of specified health symptoms. Differences between subgroups were ascertained by using binary logistic regression analysis to calculate OR and 95% CI. Univariate comparisons were followed by multivariable-adjusted models with adjustment for possible confounding by all other demographic, service and lifestyle factors considered (only factors with evidence of confounding, based on evidence from the literature and changes in the risk estimate when the covariate was included, were retained in the model). Physical and mental health scores from the SF-12 instrument were continuous variables and comparisons between subgroups carried out using multivariable linear regression weighted for non-response. The total number of health symptoms was modelled as a count variable, using the negative binomial model with robust SE. All data were weighted for non-response, based on the response rates in the three service groups (Navy, Army, Air Force), four service status groups (active regulars, active reserves, inactive reserve, ex-serving), three rank categories (commissioned officers, non-commissioned officers, other ranks), and among males and females. This resulted in seventy-two strata and weights calculated were the inverse of the response rate in each stratum. Count data (number of participants) in subgroups of supplement use are presented as unweighted data, but all summary statistics such as percentages, means and OR were based on weighted data. All analyses were carried out using the statistical software package SAS version 9.3.

Results

A total of 26 239 eligible ADF members were invited to participate and 14 032 (53%) responded to the survey. Most of the respondents were male (88.9%), mean age was 37 years (range 18–68 years), and 49.3% were Army, 23.6% Navy and 27.1% Air Force personnel. Alcohol use was common (48.3% consumed 1–2 alcoholic drinks/d, 33.9% drank 3–5 alcoholic drinks/d and 6.1% drank ≥ 6 alcoholic drinks/d), and 28.2% were current smokers. A small proportion (6.0%) of the participants had left the ADF since serving in the Middle East. Mean age of ex-serving participants (36.4 years) was similar to that of active regular personnel (35.8 years), while the average age of active reserve and inactive reserve personnel was slightly higher (39.7 and 37.2 years, respectively).

Analysis of non-response was reported previously⁽²⁵⁾ and showed that response rates were higher among females (59% of females responded *v.* 52% of males), older persons (67% of those aged ≥ 45 years responded *v.* 39% of those aged 18–25 years old), personnel of the Air Force (60% responded *v.* 50% of those in the Navy and 51% of those in the Army), those ranked as officer

(63% of officers responded *v.* 35% of those in other ranks) and personnel actively serving in the military (58% of those in active employment responded *v.* 32% of ex-serving personnel).

Use of dietary supplements was common among the study participants, with 21.5% of respondents using a dietary supplement in any of the three categories once weekly or more often. Overall, 32.3% of participants used any of the three types of supplement investigated (at any frequency of use), with bodybuilding supplements (17.5%) and energy supplements (24.5%) used more often than weight-loss supplements (7.6%); 11.6% of participants used supplements in more than one category. A large proportion of users of bodybuilding supplements used these daily (47.9%), while 21.7% used these weekly and 30.4% monthly. Energy supplements were used daily by 28.8% of users of these supplements, weekly by 27.1% and monthly by 44.2%. Weight-loss supplements were used daily by 42.5% of respondents who used such supplements, weekly by 15.5% and monthly by 42.0%. The number of different bodybuilding supplements used per person ranged from 1 to 5 different products (average 1.1 products per person), and this ranged from 1 to 4 different products (average 1.0 product per person) for energy supplements, and from 1 to 2 different products (average 1.0 product per person) for weight-loss products.

Types of supplements

In terms of the types of supplements used, amino acid and protein supplements were most commonly used. Overall, 51.5% of supplement users used amino acid and protein products, followed by multivitamin or herb compounds (14.9%) and carbohydrates and sugars (11.9%). Overall, 40.0% of all supplement users used a supplement containing caffeine (11.7% of all study participants) and 12.2% of supplement users took a product containing creatine (3.6% of all study participants), including 4.1% who used a pure creatine monohydrate product.

Table 1 and Fig. 1 show the distribution of supplement types reported within the three categories of supplements investigated. Users of bodybuilding supplements most commonly used amino acid and protein supplements, typically taken as bars or powder (90.1% of all bodybuilding supplement users used these products). Multivitamin and herb compounds marketed to aid bodybuilding were also commonly used as a bodybuilding supplement (9.3%), followed by products predominantly containing creatine monohydrate (8.5%) and other products (2.8%). Besides creatine, multivitamin and herb products marketed for bodybuilding also commonly contained caffeine (24.0% of such products).

Users of energy supplements most commonly used products that contained stimulants (54.6% of all energy supplement users), almost always including caffeine. Users of energy supplements also often reported using amino acid and protein products (21.2%), as well as

Table 1 Subgroups of supplements used by current and ex-serving Australian Defence Force personnel (*n* 14 032) who deployed to the Middle East between 2001 and 2009

	Total		Contains creatine		Contains caffeine	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Bodybuilding supplements						
Amino acids/protein products	1702	90.1	86	5.1	17	1.0
Multivitamin/herb compound	175	9.3	72	41.1	42	24.0
Carbohydrates and sugars	0	0.0				
Stimulants	0	0.0				
Creatine monohydrate	160	8.5	160	100.0	0	0.0
Other (non-prescription)	9	0.5	0	0.0	0	0.0
Other (prescription medicine)	0	0.0				
Total users of bodybuilding supplements*	1890	100.0	307	16.2	59	3.1
Energy supplements						
Amino acids/protein products	568	21.2	201	35.4	19	3.3
Multivitamin/herb compound	505	18.9	0	0.0	1	0.2
Carbohydrates and sugars	249	9.3	0	0.0	7	2.8
Stimulants	1460	54.6	1	0.1	1442	98.8
Creatine monohydrate	0	0.0				
Other (non-prescription)	0	0.0				
Other (prescription medicine)	0	0.0				
Total users of energy supplements*	2676	100.0	202	7.5	1463	54.7
Weight-loss supplements						
Amino acids/protein products	386	47.5	3	0.8	13	3.4
Multivitamin/herb compound	99	12.2	0	0.0	18	18.2
Carbohydrates and sugars	223	27.4	0	0.0	0	0.0
Stimulants	78	9.6	0	0.0	76	97.4
Creatine monohydrate	0	0.0				
Other (non-prescription)	7	0.9	0	0.0	0	0.0
Other (prescription medicine)	47	5.8	0	0.0	0	0.0
Total users of weight-loss supplements*	813	100.0	3	0.4	105	12.9

*Total number of Australian Defence Force personnel in these categories do not equal to the sum of column responses because participants may have used more than one type of bodybuilding/energy/weight-loss supplement.




Fig. 1 Subgroups of supplements (■, bodybuilding supplements; □, energy supplements; ▨, weight-loss supplements) used by current and ex-serving Australian Defence Force personnel (*n* 14 032) who deployed to the Middle East between 2001 and 2009

multivitamin and herb products (18.9%), but less often (9.3%) products that predominantly contained carbohydrates and sugars. Users of weight-loss supplements most often used products that supplied protein and amino acids (47.5%), carbohydrates and sugars (27.4%), or multivitamin and herb compounds marketed to support weight loss (12.2%). Stimulants (9.6%), mainly containing caffeine (97.4% of such products), were also used as weight-

loss supplements, while a small proportion of weight-loss supplement users (5.8%) reported using prescription medicines to support weight loss.

Factors associated with supplement use

Men were more likely than women to use bodybuilding and energy supplements, while women were more likely to use weight-loss supplements. There was a clear

age-related pattern of use, particularly for bodybuilding and energy supplements, with around a third of the youngest participants (aged 18–29 years) using such supplements, compared with less than 15% of the oldest (≥ 40 years of age) study participants (all $P < 0.001$; Table 2).

Army personnel were more likely to use of bodybuilding and energy supplements (about 25% for both types of supplements) compared with Navy and Air Force personnel (12.2–24.0%), while use of weight-loss supplements was somewhat more common in Navy personnel (9.2%) compared with Army and Air Force (8.2 and 5.0%, respectively). Use of supplements (any type) was generally highest among personnel in lower ranks, those deployed in active regular service and in combat positions (Table 2).

There was some evidence of an association between supplement use and other lifestyle factors. Current smokers were more likely to use energy and weight-loss supplements than never smokers, although differences between these groups were small (Table 2). Participants with a high intake of caffeine-containing drinks (≥ 6 cups/d) were more likely to use energy and weight-loss supplements compared with persons who did not consume caffeine-containing drinks, while bodybuilding supplements were less frequently used by participants with moderate (3–5 cups/d) intake levels of caffeine-containing drinks compared with respondents who did not consume such drinks (Table 2).

Health status

In terms of recent symptoms reported by study participants, there were consistent patterns within each of the three categories of supplements (Table 3). Participants who used bodybuilding supplements reported less frequently than non-users that they suffered from a range of health problems including headaches, sleeping difficulties, flatulence, diarrhoea and indigestion, whereas such problems were more common among users of energy supplements and weight-loss supplements compared with non-users (Table 3). Multivariable modelling showed that these patterns were independent of the general demographic and job characteristics associated with supplement use, as identified in Table 2.

Major depressive syndrome was more common in users *v.* non-users of energy supplements (4.2 *v.* 3.4%) and weight-loss supplements (6.3 *v.* 3.4%), but did not differ between users and non-users of bodybuilding supplements after adjustment for confounders (Table 3). Also, in terms of summary measures of physical and mental health (using the SF-12 assessment tool), physical health scores were lower on average for users of weight-loss supplements compared with non-users, but physical health was better in users compared with non-users of bodybuilding supplements (Table 3). Mental health summary scores were lower in users of energy and weight-loss supplements compared with non-users. The frequency of supplement use among users was lower for bodybuilding

supplements and higher for energy supplements if the participant reported recent health symptoms; there was no such pattern for weight-loss supplements (detailed results not shown).

Discussion

Our findings show that bodybuilding, energy and weight-loss supplements are commonly used by Australian military personnel who have served in the Middle East: about one in three persons reported using at least one of these types of supplements, with one in five persons using a supplement at least once weekly. This level of supplement use appears to be higher than that in the general Australian population. National survey data showed that 19% of Australian males aged 19–30 years, and 25% of women in the same age group, use dietary supplements⁽¹⁾, although specific national data on the types of supplements assessed in our study are not available for direct comparisons. We did not ask participants to report their reasons for dietary supplement use, but data from British Army soldiers suggest that general improvement of physical performance, and preparation prior to and recovery after training or physical activity may be the main motivating factors for choosing to use dietary supplements among military personnel⁽¹⁸⁾. US data indicate that energy drinks tend to be used by Army personnel to boost energy and enhance mental alertness⁽²⁸⁾.

Compared with other military populations, supplement use was not as common in our study of Australian military personnel as in similar groups in other countries. A large survey of more than 16 000 US military personnel in 2005 indicated that 25% used bodybuilding supplements (compared with 17% in our study) and 21% used weight-loss supplements (compared with only 7% in our study), while use of energy supplements could not be compared directly between the two studies⁽²⁹⁾. Very similar to our observations, use of bodybuilding supplements was more common in US Army than Navy and Air Force personnel, and more common in younger than older persons, in men, and in non-smokers compared with smokers⁽²⁹⁾. Results from the US Millennium Cohort Study also indicated a higher overall use of supplements in US compared with Australian military personnel, with 47% of US military personnel reporting use of at least one type of bodybuilding, energy or weight-loss supplement (compared with 32% in our study). In particular, the use of energy supplements (38%) and weight-loss supplements (19%) was much higher in US compared with Australian military personnel⁽³⁰⁾. The level of dietary supplement use in our study participants appears more similar to the level of use reported by British than US military personnel. In a survey of UK-based soldiers in training, 38% reported current use of supplements, most commonly protein supplements and energy drinks⁽¹⁸⁾, which is comparable to the level of use reported in our study.

Table 2 Associations between general demographic, service and lifestyle factors and use of dietary supplements by current and ex-serving Australian Defence Force personnel (*n* 14 032) who deployed to the Middle East between 2001 and 2009

	Bodybuilding supplement use					Energy supplement use					Weight-loss supplement use				
	Total*	% User†	AOR‡	95 % CI	<i>P</i>	Total*	% User†	AOR§	95 % CI	<i>P</i>	Total*	% User†	AOR	95 % CI	<i>P</i>
Overall	13 308	17.5	–	–	–	13 305	24.5	–	–	–	13 290	7.6	–	–	–
Sex															
Male	11 667	19.0	3.67	3.03, 4.45	<0.001	11 665	25.0	1.23	1.09, 1.39	0.001	11 646	7.0	0.51	0.43, 0.59	<0.001
Female	1641	6.3	Ref.			1640	20.7	Ref.			1644	12.0	Ref.		–
Age group (years)															
18–29	3002	34.6	7.65	6.67, 8.76	<0.001	2995	36.4	2.63	2.37, 2.92	<0.001	2997	9.6	1.41	1.20, 1.67	<0.001
30–39	5385	17.2	3.29	2.91, 3.72	<0.001	5385	25.6	1.86	1.70, 2.04	<0.001	5377	7.8	1.18	1.03, 1.36	0.02
≥40	4921	5.8	Ref.		–	4925	14.7	Ref.		–	4916	5.8	Ref.		–
Service															
Army	6214	22.3	1.27	1.14, 1.42	<0.001	6212	27.3	1.12	1.03, 1.23	0.01	6211	8.2	1.51	1.29, 1.75	<0.001
Navy	2987	13.8	0.80	0.68, 0.93	0.004	2985	24.0	0.98	0.87, 1.10	0.69	2981	9.2	1.57	1.30, 1.88	<0.001
Air Force	4107	12.2	Ref.		–	4108	19.7	Ref.		–	4098	5.0	Ref.		–
Rank															
Commissioned officer	3985	11.1	Ref.		–	3981	16.7	Ref.		–	3972	4.8	Ref.		–
Non-commissioned officer	7881	17.1	1.34	1.21, 1.48	<0.001	7886	24.8	1.40	1.28, 1.52	<0.001	7879	8.4	1.65	1.44, 1.89	<0.001
Lower ranks	1442	29.1	1.39	1.19, 1.63	<0.001	1438	35.5	1.59	1.38, 1.83	<0.001	1439	8.9	1.34	1.06, 1.68	0.01
Service status															
Active regular	9583	18.5	1.39	1.10, 1.76	0.006	9581	24.6	0.85	0.72, 1.02	0.08	9574	8.0	1.31	0.97, 1.76	0.08
Active reserve	1613	14.5	1.32	1.01, 1.76	0.04	1612	22.3	0.94	0.77, 1.15	0.54	1605	7.1	1.21	0.87, 1.69	0.26
Inactive reserve	1318	18.0	1.31	1.01, 1.71	0.05	1321	23.6	0.82	0.66, 1.00	0.05	1319	6.5	1.06	0.75, 1.50	0.76
Ex-serving	794	14.2	Ref.		–	791	27.7	Ref.		–	792	6.9	Ref.		–
Main role on deployment															
Combat	3739	25.7	1.84	1.62, 2.10	<0.001	3736	31.5	1.31	1.18, 1.46	<0.001	3731	9.6	1.53	1.29, 1.82	<0.001
Combat support	2384	14.5	1.22	1.06, 1.40	0.007	2381	19.6	0.85	0.75, 0.95	0.005	2383	6.2	1.04	0.86, 1.25	0.69
Medical/welfare	495	12.4	1.73	1.33, 2.25	<0.001	495	16.6	0.89	0.72, 1.12	0.32	494	7.3	1.22	0.89, 1.67	0.22
Air crew	709	9.0	0.89	0.71, 1.13	0.34	709	18.7	0.98	0.83, 1.17	0.85	708	3.7	0.87	0.63, 1.20	0.39
Maritime operations	304	10.1	1.17	0.80, 1.73	0.42	304	20.3	0.93	0.70, 1.24	0.61	304	8.3	1.20	0.79, 1.81	0.39
Logistics	2412	13.5	Ref.		–	2419	23.5	Ref.		–	2415	6.2	Ref.		–
Administration	228	8.2	0.92	0.62, 1.36	0.66	227	17.3	0.78	0.59, 1.04	0.09	228	8.3	1.11	0.76, 1.62	0.6
Other roles	1296	11.4	1.44	1.20, 1.72	<0.001	1297	17.9	1.00	0.87, 1.16	0.96	1294	5.8	1.16	0.92, 1.47	0.21
Smoking history															
Current	3535	17.4	0.81	0.73, 0.90	<0.001	3534	28.6	1.17	1.07, 1.27	<0.001	3536	8.5	1.14	0.97, 1.34	0.1
Ex-smoker	2345	15.0	1.12	0.98, 1.27	0.09	2345	20.8	0.97	0.87, 1.08	0.54	2341	7.3	1.11	0.96, 1.28	0.16
Never smoked	7154	18.5	Ref.		–	7152	23.6	Ref.		–	7141	7.2	Ref.		–
Intake of caffeine-containing drinks															
None	1529	23.0	Ref.		–	1528	20.7	Ref.		–	1526	7.9	Ref.		–
1–2/d	6374	20.1	0.99	0.86, 1.13	0.85	6372	26.5	1.62	1.42, 1.86	<0.001	6361	7.5	1.05	0.87, 1.27	0.61
3–5/d	4557	12.7	0.77	0.67, 0.90	0.001	4557	22.6	1.63	1.41, 1.88	<0.001	4558	7.1	1.11	0.91, 1.37	0.3
≥6/d	813	13.8	0.90	0.71, 1.13	0.34	814	26.2	1.83	1.50, 2.22	<0.001	812	10.1	1.49	1.13, 1.95	0.004

AOR, adjusted odds ratio; ref., reference category.

*Unweighted totals; totals may not add up due to missing responses.

†Percentages weighted for non-response; percentages may not sum to 100 due to rounding.

‡Adjusted binary logistic regression models include all demographic, service and lifestyle variables listed in the table.

§Adjusted binary logistic regression models include the variable of interest adjusted for the confounding effects of gender, age group, service, rank, smoking history at survey completion and main role on deployment.

|| Adjusted binary logistic regression models include the variable of interest adjusted for the confounding effects of gender, age group, service, rank, caffeine use at survey completion and main role on deployment.

Table 3 Associations between physical and mental health and use of dietary supplements by current and ex-serving Australian Defence Force personnel (*n* 14 032) who deployed to the Middle East between 2001 and 2009

	Bodybuilding supplement use					Energy supplement use					Weight-loss supplement use				
	Users (<i>n</i> 2169)	Non-users (<i>n</i> 11 029)	AOR*	95% CI	<i>P</i>	Users (<i>n</i> 3068)	Non-users (<i>n</i> 10 128)	AOR*	95% CI	<i>P</i>	Users (<i>n</i> 980)	Non-users (<i>n</i> 12 201)	AOR*	95% CI	<i>P</i>
Recent health symptoms (%)															
Headaches	40.5	49.6	0.80	0.73, 0.87	<0.001	51.8	46.7	1.32	1.23, 1.43	<0.001	53.2	47.5	1.18	1.05, 1.33	0.006
Sleeping difficulties	53.7	56.0	1.10	1.00, 1.20	0.05	61.0	53.9	1.46	1.35, 1.58	<0.001	67.2	54.7	1.73	1.53, 1.96	<0.001
Constipation	7.7	10.5	1.02	0.86, 1.21	0.83	10.5	9.8	1.30	1.14, 1.47	<0.001	13.1	9.7	1.37	1.14, 1.63	<0.001
Flatulence or burping	23.9	30.7	0.87	0.79, 0.97	0.01	31.5	28.8	1.35	1.24, 1.47	<0.001	33.3	29.1	1.26	1.11, 1.43	<0.001
Stomach cramps	12.6	14.0	1.01	0.88, 1.16	0.88	15.9	13.1	1.37	1.23, 1.53	<0.001	18.4	13.4	1.37	1.17, 1.61	<0.001
Diarrhoea	17.9	20.8	0.88	0.78, 0.99	0.04	23.0	19.4	1.31	1.20, 1.44	<0.001	25.6	19.9	1.39	1.22, 1.59	<0.001
Indigestion	13.3	18.6	0.85	0.74, 0.97	0.01	19.0	17.3	1.34	1.21, 1.48	<0.001	21.2	17.4	1.30	1.12, 1.51	<0.001
Nausea	6.0	7.6	0.91	0.75, 1.10	0.34	8.9	6.8	1.48	1.28, 1.71	<0.001	9.5	7.1	1.17	0.95, 1.44	0.14
Vomiting	3.2	3.6	0.92	0.71, 1.19	0.29	4.2	3.2	1.39	1.14, 1.70	0.001	4.8	3.4	1.23	0.92, 1.64	0.17
Total number of health symptoms, mean†	1.8	2.1	0.95	0.90, 1.01	0.0824	2.2	1.9	1.20	1.16, 1.25	<0.001	2.4	2.0	1.19	1.12, 1.27	<0.001
sd	1.8	1.9				2.0	1.9				2.1	1.9			
Health status															
Physical health, mean‡	53.1	51.2	1.1	0.8, 1.5	<0.001	51.6	51.5	-0.3	-0.7, -0.01	0.05	49.9	51.7	-1.6	-2.2, -1.1	<0.001
sd	7.4	8.4				8.4	8.2				9.2	8.2			
Mental health, mean§	49.7	49.3	-0.01	-0.5, 0.5	0.96	48.1	49.7	-1.7	-2.1, -1.3	<0.001	46.9	49.5	-2.2	-2.9, -1.6	<0.001
sd	10.0	10.2				10.7	10.0				10.9	10.1			
Major depressive syndrome (%)	2.4	3.9	0.79	0.57, 1.10	0.16	4.2	3.4	1.33	1.06, 1.66	0.01	6.3	3.4	1.91	1.45, 2.53	<0.001

AOR, adjusted odds ratio.

*Binary logistic regression models comparing users *v.* non-users, adjusting for sex, age group, service, rank, service status, main role during most recent deployment, current smoking status (bodybuilding and energy supplements only) and consumption of caffeine-containing drinks (bodybuilding and weight-loss supplements only).†Unweighted means and *sd*.‡Unweighted means and *sd* of Short Form-12 Health Survey Physical Component Score.§Unweighted means and *sd* of Short Form-12 Health Survey Mental Component Score.

Our data indicated quite clear patterns of personal characteristics that were associated with use of supplements. Bodybuilding supplements tended to be used more often by individuals in active service, with better health behaviours and of better self-reported health (with a lower level of supplement use if the person reported health symptoms). In contrast, use of energy and weight-loss supplements was not associated with service status (active or inactive), and these supplements were more commonly used by participants with poorer health behaviours and with poorer physical and mental health. However, differences in mental and physical health between groups were generally very small and the clinical significance of these differences is uncertain.

This patterning of demographic characteristics with supplement type is very similar to that observed in other studies. For example, in the US Millennium Cohort Study, users of bodybuilding supplements were generally younger, more physically active and healthier, while users of energy and weight-loss supplements tended to be less physically active and to have poorer health indicators⁽³⁰⁾. Similarly in British Army soldiers in training, overall use of dietary supplements was associated with higher levels of physical activity, although supplement use was higher in smokers compared with non-smokers⁽¹⁸⁾; we could not directly compare data for subgroups of supplements.

Associations with recent health symptoms and summary measures of health status also showed fairly consistent patterning, in that users of bodybuilding supplements had better physical and mental health, in contrast to users of energy and weight-loss supplements. This pattern of associations is very similar to that reported for US Army staff⁽³⁰⁾, including that of a large survey of US Armed Forces which found that multivitamin/minerals and protein/amino acid supplements were associated with better general health, eating habits and fitness levels⁽³¹⁾. Recent US data also indicated that the use of a combination of dietary supplement products is most commonly associated with side-effects^(32,33). However, because details of health symptoms in our study participants were not specifically reported in the context of dietary supplement use, we cannot draw any conclusions about side-effects from our data.

We found that the prevalence of major depressive syndrome was twice as high in users of weight-loss supplements than among non-users of weight-loss supplements. Due to the cross-sectional nature of our data collection we are unable to ascertain which factors are causes and which factors are consequences in these associations, but this observation raises the possibility that the increasing prevalence of overweight and obesity among military populations^(7,8) may have impacts on mental as well as physical health in this group. Measures of overweight or obesity were unfortunately not available for our study participants.

We were particularly interested in the use of supplements that contain caffeine. Military personnel may have

higher levels of caffeine intake than the general population because caffeine alleviates the adverse cognitive consequences of sleep deprivation (e.g. due to long work shifts) and it enhances physical performance⁽¹³⁾. However, there is also concern about possible side-effects of high caffeine intake^(34,35), such as disturbed sleep patterns⁽¹⁴⁾ as a result of use of caffeinated products during overnight periods of sustained wakefulness.

In our study, sleeping difficulties were higher in users of energy supplements (55% of which contained caffeine) and weight-loss supplements (13% of which contained caffeine) compared with non-users of such supplements. However, because of the cross-sectional nature of our data, and because we were unable to quantify caffeine intake, it is not possible to draw any firm conclusions about the role of caffeine-containing supplements in these health symptoms. Nevertheless, our data show that use of caffeine-containing supplements was common (11.7% of all participants used these), which adds to the caffeine intake obtained from caffeine-containing drinks (40% of respondents consumed three or more caffeine drinks daily), raising the possibility of very high caffeine intake levels in some persons. Observations from US military personnel have also indicated that use of caffeine-containing energy drinks is very common, particularly among young males, and that sleeping difficulties may be associated with this⁽²⁸⁾. Our data did not facilitate calculation of total caffeine intake for the study participants. However, it is known that the caffeine content of supplements is highly variable and poorly listed in product information⁽³⁴⁾, and thus difficult to quantify. A more detailed study on levels of caffeine intake in the Australian military, or monitoring of levels of intake at an individual level, and its impact on the health of serving personnel may therefore be warranted.

Our study is the first to report supplement use in Australian military personnel, and a major strength of the study is its large number of participants and comprehensive assessment of physical and mental health, as well as demographic and lifestyle factors. Other strengths include the fact that the response rate compared favourably with other, similar studies (as discussed in detail in the MEAO Survey report⁽²⁵⁾) and the results were weighted for non-response to control for potential response bias. Furthermore, standard, validated instruments were used, which allows comparison with other military and civilian studies.

However, the study has some limitations that need to be considered when interpreting the results. The data on dietary supplement use were self-reported by the study participants and such self-reporting may be influenced by reporting biases⁽³⁶⁾. It is a cross-sectional study and thus we were unable to ascertain whether health symptoms were a cause or a consequence of supplement use. Health status indicators were based on self-reported data, which may be affected by recall bias. However, these health data were collected as part of a larger health survey and not

specifically in the context of a study of dietary supplements, thus it is very unlikely that misreporting (if any) caused a systematic bias in our present analyses. Full details on the frequency dose of supplements taken, and use of dietary supplements other than those in the categories reported here, were not ascertained in the MEAO Survey questionnaire. Thus, unfortunately, we cannot directly compare absolute amounts of supplement use (of any type) with other populations.

There is a variety of reasons why patterns of dietary supplement use may be different among military (and general) populations of different nations, for example due to different availability of supplements, diverse product ranges, different marketing practices, different regulatory laws, differences in dietary and related health habits, etc. One of the problems with comparisons between different studies is the differences in the definition of 'supplement user'. For example, we have chosen to focus our analyses on supplement users at any frequency of use, similar to others^(18,30), while some different studies have focused on persons who use a dietary supplement at least once weekly^(6,32). The questionnaire items relating to dietary supplement use in the larger health questionnaire of the MEAO Survey, from which our data were extracted, were designed to focus on the intended use of supplements (bodybuilding, energy or weight loss). However, in our analyses we have also incorporated a more ingredient-focused categorisation (amino acids, multivitamins, etc.), to the best of our ability given the available data and resources. While direct comparisons of dietary supplement use in different nations will always have limitations due to the inherent differences in dietary supplement products available around the world, it would be very useful if an agreement could be reached over a standardised method of categorising these products, as well as the frequency of use that would characterise someone as a regular supplement user.

We were able to find product information for almost all supplements reported by the study participants, but a very small number of misclassifications may have occurred. We observed that products that may be quite similar in composition are sometimes marketed for divergent purposes of use (e.g. protein supplements are marketed as bodybuilding supplements as well as weight-loss supplements), and conversely there are products that fall into the same category (e.g. weight loss) but have very little overlap in terms of ingredients⁽³⁷⁾. This makes it difficult to study health effects of specific supplement ingredients in detail, but at the same time this emphasises the need to monitor for possible side-effects of dietary supplements at an individual level, including among military personnel⁽³⁸⁾. General provision of information about dietary supplements⁽³⁹⁾ would be prudent for military personnel. Military personnel are known to be healthier on average than members of the general population (the healthy soldier effect⁽⁴⁰⁾), but our findings indicate

that regardless of their better health, use of dietary supplements is very common among Australian military personnel. Our data may help those in charge of reviewing the health status of ADF personnel to identify priorities for further, more detailed investigations of supplement use by military personnel, and to review the potential risks associated with use of certain dietary supplements in this large population group. The recent banning of all DMAA (dimethylamylamine) containing supplements by the Australian Therapeutic Goods Administration illustrates the need for regulation to prevent avoidable risks to the health of military personnel and the population at large.

Conclusion

In conclusion, our study shows that bodybuilding and energy supplements are commonly used by ADF personnel who have served in the Middle East. Our data indicate that patterns of use are associated with employment and lifestyle factors, as well as health symptoms. Although we were unable to ascertain absolute amounts of intake of dietary supplements and ingredients, our findings may help identify subgroups of Australian military personnel who may be at risk of possible side-effects.

Acknowledgements

Acknowledgements: The authors thank the veterans who participated in the survey. Helpful comments on this manuscript by Lieutenant Colonel Peter Nasveld are gratefully acknowledged. *Financial support:* The MEAO Census Study was funded by the Australian Government Department of Defence. The authors used data from the MEAO Census Study during the course of their employment with The University of Queensland, School of Public Health. The authors received no funding from the Australian Government Department of Defence for this specific work. The opinions or assertions contained herein are the private views of the authors and are not to be construed as official, or representing the views of the Australian Defence Force or Department of Defence. The Department of Defence had no role in the design, analysis or writing of this article. *Conflict of interest:* None. *Authorship:* J.C.v.d.P. oversaw data extraction and analyses, and wrote the paper; J.K. supported extraction of product data, carried out statistical analyses and helped write the paper; A.B. extracted product data and helped write the paper; C.-W.L. conceptualised the idea and helped write the paper. All authors read and approved the final manuscript. *Ethics of human subject participation:* Ethical approval for the present study was obtained from the Australian Department of Defence (protocol number LREP 14-010) and The University of Queensland (protocol number 2009001441) ethics committees. Participants of

the MEAO Survey provided consent in writing or electronically, and all aspects of this research complied with the Declaration of Helsinki.

References

- Australian Bureau of Statistics (2014) Australian Health Survey: Nutrition First Results – Foods and Nutrients, 2011–12. <http://www.abs.gov.au/australianhealthsurvey/> (accessed April 2016).
- Hathcock JN (1997) Vitamins and minerals: efficacy and safety. *Am J Clin Nutr* **66**, 427–437.
- Jimenez-Flores R, Heick J, Davis SC *et al.* (2012) A comparison of the effects of a high carbohydrate vs. a higher protein milk supplement following simulated mountain skirmishes. *Mil Med* **177**, 723–731.
- Naclerio F & Larumbe-Zabala E (2016) Effects of whey protein alone or as part of a multi-ingredient formulation on strength, fat-free mass, or lean body mass in resistance-trained individuals: a meta-analysis. *Sports Med* **46**, 125–137.
- Craig WJ, Mangels AR & American Dietetic Association (2009) Position of the American Dietetic Association: vegetarian diets. *J Am Diet Assoc* **109**, 1266–1282.
- Lieberman HR, Stavinocha TB, McGraw SM *et al.* (2010) Use of dietary supplements among active-duty US Army soldiers. *Am J Clin Nutr* **92**, 985–995.
- Peake J, Gargett S, Waller M *et al.* (2012) The health and cost implications of high body mass index in Australian defence force personnel. *BMC Public Health* **12**, 451.
- Reyes-Guzman CM, Bray RM, Forman-Hoffman VL *et al.* (2015) Overweight and obesity trends among active duty military personnel: a 13-year perspective. *Am J Prev Med* **48**, 145–153.
- Jenkinson DM & Harbert AJ (2008) Supplements and sports. *Am Fam Physician* **78**, 1039–1046.
- Bishop D (2010) Dietary supplements and team-sport performance. *Sports Med* **40**, 995–1017.
- Trexler ET & Smith-Ryan AE (2015) Creatine and caffeine: considerations for concurrent supplementation. *Int J Sport Nutr Exerc Metab* **25**, 607–623.
- Spriet LL (2014) Exercise and sport performance with low doses of caffeine. *Sports Med* **44**, Suppl. 2, S175–S184.
- Lieberman HR, Stavinocha T, McGraw S *et al.* (2012) Caffeine use among active duty US Army soldiers. *J Acad Nutr Diet* **112**, 902–912, 912.e1–e4.
- Clark I & Landolt HP (2016) Coffee, caffeine, and sleep: a systematic review of epidemiological studies and randomized controlled trials. *Sleep Med Rev* **31**, 70–78.
- Waits WM, Ganz MB, Schillreff T *et al.* (2014) Sleep and the use of energy products in a combat environment. *US Army Med Dep J* Oct–Dec issue, 22–28.
- Burke J, Seda G, Allen D *et al.* (2007) A case of severe exercise-induced rhabdomyolysis associated with a weight-loss dietary supplement. *Mil Med* **172**, 656–658.
- Manore MM (2012) Dietary supplements for improving body composition and reducing body weight: where is the evidence? *Int J Sport Nutr Exerc Metab* **22**, 139–154.
- Casey A, Hughes J, Izard RM *et al.* (2014) Supplement use by UK-based British Army soldiers in training. *Br J Nutr* **112**, 1175–1184.
- Boos CJ, Simms P, Morris FR *et al.* (2011) The use of exercise and dietary supplements among British soldiers in Afghanistan. *J R Army Med Corps* **157**, 229–232.
- Austin KG, McLellan TM, Farina EK *et al.* (2016) Soldier use of dietary supplements, including protein and body building supplements, in a combat zone is different than use in garrison. *Appl Physiol Nutr Metab* **41**, 88–95.
- Boos CJ, Wheble GAC, Campbell MJ *et al.* (2010) Self-administration of exercise and dietary supplements in deployed British military personnel during Operation TELIC 13. *J R Army Med Corps* **156**, 32–36.
- Cassler NM, Sams R, Cripe PA *et al.* (2013) Patterns and perceptions of supplement use by US Marines deployed to Afghanistan. *Mil Med* **178**, 659–664.
- Austin K, Price L, McGraw S *et al.* (2014) Longitudinal trends of dietary supplement use by US Army soldiers differ from the US civilian population. *FASEB J* **28**, 245–246.
- Greenwood MRC & Oria M (2008) *Use of Dietary Supplements by Military Personnel*. Washington, DC: National Academies Press.
- Dobson A, Treloar S, Zheng WY *et al.* (2013) *The Middle East Area of Operations (MEAO) Health Study: Census Study Report*, vol. 1. Brisbane: The The University of Queensland, Centre for Military and Veterans' Health.
- Ware J, Kosinski M & Keller S (1996) A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. *Med Care* **34**, 220–233.
- Kroenke K, Spitzer RL & Williams JBW (2001) The PHQ-9. *J Gen Intern Med* **16**, 606–613.
- Stephens MB, Attipoe S, Jones D *et al.* (2014) Energy drink and energy shot use in the military. *Nutr Rev* **72**, 72–77.
- Kao T-C, Deuster PA, Burnett D *et al.* (2012) Health behaviors associated with use of body building, weight loss, and performance enhancing supplements. *Ann Epidemiol* **22**, 331–339.
- Jacobson IG, Horton JL, Smith B *et al.* (2012) Bodybuilding, energy, and weight-loss supplements are associated with deployment and physical activity in US military personnel. *Ann Epidemiol* **22**, 318–330.
- Austin KG, McGraw SM & Lieberman HR (2014) Multivitamin and protein supplement use is associated with positive mood states and health behaviors in US Military and Coast Guard personnel. *J Clin Psychopharmacol* **34**, 595–601.
- Knapik JJ, Trone DW, Austin KG *et al.* (2016) Prevalence, adverse events, and factors associated with dietary supplement and nutritional supplement use by US Navy and Marine Corps personnel. *J Acad Nutr Diet* **116**, 1423–1442.
- Austin KG, Farina EK & Lieberman HR (2016) Self-reported side-effects associated with use of dietary supplements in an armed forces population. *Drug Test Anal* **8**, 287–295.
- Cohen PA, Attipoe S, Travis J *et al.* (2013) Caffeine content of dietary supplements consumed on military bases. *JAMA Intern Med* **173**, 592–594.
- Wesensten NJ (2014) Legitimacy of concerns about caffeine and energy drink consumption. *Nutr Rev* **72**, 78–86.
- Sikkens JJ, van Eijnsden M, Bonsel GJ *et al.* (2011) Validation of self-reported folic acid use in a multiethnic population: results of the Amsterdam Born Children and their Development study. *Public Health Nutr* **14**, 2022–2028.
- Geller AI, Shehab N, Weidle NJ *et al.* (2015) Emergency department visits for adverse events related to dietary supplements. *N Engl J Med* **373**, 1531–1540.
- Siano KA (2014) Renal failure in a soldier taking N.O.-Xplode. *J Am Board Fam Med* **27**, 565–569.
- The Canadian Forces Health Services Group (2004) The Dietary Supplement Dilemma: Helpful or Harmful? <http://publications.gc.ca/site/fra/9.835612/publication.html> (accessed May 2016).
- McLaughlin R, Nielsen L & Waller M (2008) An evaluation of the effect of military service on mortality: quantifying the healthy soldier effect. *Ann Epidemiol* **18**, 928–936.