

- group A *Streptococcus* traced to a vaginal carrier. *J Infect Dis* 1978;138:287.
9. Larson E. Guideline for use of topical antimicrobial agents. *Am J Infect Control* 1988;16:253.
 10. Groschel DHM, Pruett TL. Surgical antisepsis. In: Block SS, ed. *Disinfection, Sterilization and Preservation*. 4th ed. Philadelphia, PA Lea & Febiger; 1991.
 11. Culver DH, Horan TC, Gaynes RP, et al. Surgical wound infection rates by wound class, operative procedure, and patient risk index. *Am J Med* 1991;91(suppl 3B):152S.
 12. Nosocomial infection rates for interhospital comparison: limitations and possible solutions. *Infect Control Hosp Epidemiol* 1991;12:609.
 13. Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori TG. CDC definitions of nosocomial surgical site infections, 1992: a modification of CDC definitions of surgical wound infections. *Am J Infect Control* 1992;20:271.
 14. Webster J, Faoagali JL. An in-use comparison of chlorhexidine gluconate 4% w/v, glycol-poly-siloxane plus methylcellulose and a liquid soap in a special care baby unit. *J Hosp Infect* 1989;14:141.
 15. Sanderson PJ, Weissler S. A comparison of the effect of chlorhexidine antisepsis, soap and antibiotics on bacteriuria, perineal colonization and environmental contamination in spinally injured patients. *J Hosp Infect* 1990;15:235.
 16. Nicoletti G, Boghossian V, Borland R. Hygienic hand disinfection: a comparative study with chlorhexidine detergents and soap. *J Hosp Infect* 1990;15:323.
 17. Ayliffe GAF, Babb JR, Davies JG, Lill HA. Hand disinfection: a comparison of various agents in laboratory and ward studies. *J Hosp Infect* 1988;11:226.
 18. Hayek LJ, Emerson JM. Preoperative whole body disinfection—a controlled clinical study. *J Hosp Infect* 1988;11(suppl B): 15.
 19. Aly R, Maibach HI. Comparative efficacy of a 2-minute surgical scrub with chlorhexidine gluconate, povidone-iodine and chloroxlenol sponge-brushes. *Am J Infect Control* 1988;16:173.
 20. Blech ME, Hartemann P, Paquin JL. Activity of nonantiseptic soaps and ethanol for hand disinfection. *Zbl Bakt Hyg I Abt Orig B* 1985;181:496.
 21. Larson EL, Butz AM, Gullette DL, Laughon BA. Alcohol for surgical scrubbing? *Infect Control Hosp Epidemiol* 1990;11:139.
 22. Bendig JWA. Surgical hand disinfection: comparison of 4% chlorhexidine detergent solution and 2% triclosan detergent solution. *J Hosp Infect* 1990;15:143.
 23. Babb JR, Davies JG, Ayliffe GAF. A test procedure for evaluating surgical hand disinfection. *J Hosp Infect* 1991;18(suppl B):41.
 24. Rotter ML, Koller W, Wewalka G. Evaluation of procedures for hygienic hand-disinfection: controlled parallel experiments on the Vienna test model. *J Hyg Camb* 1986;96:27.
 25. Kobayashi H. Evaluation of surgical scrubbing. *J Hosp Infect* 1991;11(suppl B):29.
 26. Holloway PM, Platt JH, Reybrouk G, Lilly HA, Mehtar S, Drabu Y. A multicentre evaluation of two chlorhexidine-containing formulations for surgical hand disinfection. *J Hosp Infect* 1990;16:151.

US Approves First Vaccine for Hepatitis A

by **Gina Pugliese, RN, MS**
Medical News Editor

The US Food and Drug Administration (FDA) gave SmithKline Beecham PLC approval to distribute Havrix, the first vaccine licensed for preventing hepatitis A. The US was the 41st country to license Havrix. An FDA spokesperson

said the agency wanted to be sure that the vaccine caused no significant adverse effects in recipients. An estimated 1.3 million people in Europe and Asia have received Havrix since 1991.

The Centers for Disease Control and Prevention (CDC) has reported that hepatitis A cases are under-

reported considerably. About 23,000 cases are officially reported each year, but CDC estimates that the virus infects 150,000 Americans each year.

FROM: Altman L. U.S. approves first vaccine for hepatitis A. *New York Times* February 23, 1995.