March 1964

Vol. LXXVIII No. 3

The Journal of Laryngology and Otology

EDITED BY G. H. BATEMAN

> ASSISTANT EDITOR LIONEL TAYLOR

Contents

THE RECEPTOR ORGAN IN THE COCHLEA—2	L. NAFTALIN
THE USE OF THE C.A.L. ULTRASONIC GENERATOR IN MfiNiÈRE's DISEASE	G. KOSSOFF, A. E. KHAN and R. H. FARRANT
THE TREATMENT OF MENIÈRE'S DISEASE WITH ULTRA- SOUND	. R. B. LUMSDEN
R6LE OF COMPRESSION IN THE ^ETIOLOGY OF BELL'S	
PALSY	N. JAIN and A. P. SHARMA
PRESBYACUSIS.	
EAR PHOTOGRAPHY. K.	MCLAY and R. J. RUSSELL
CLINICAL RECORDS—	
A SOLITARY NEURILEMMOMA CAUSING PARALYSIS OF THE IXTH, XTH, XITH AND XIITH CRANIAL NERVES AND SYMPATHETIC CHAIN	P. M. BINNS
TRACHEAL STENOSIS FOLLOWING TRACHEOTOMY	P. M. BINNS
LOCALIZED LARYNGEAL AMYLOIDOSIS, A REPORT OF A CASE WITH A REVIEW OF THE LITERATURE J. C. MCA	ALPINB and A. P. FULLER
A CASE OF CHONDROMA OF THE EPIGLOTTIS	. A. B. N. RAO

CHRONIC SINUSITIS WITH ALVEOLAR FISTULAE O F MEDIAEVAL DATE

GENERAL NOTES

London

Headley Brothers

109 Kingsway WC2

Annual Subscription £5/5/0 net, U.S.A. \$15 Monthly, 12/6 net post

CALVIN WELLS

free

https://doi.org/10.1017/S0022215100062010 Published online by Cambridge University Press

ADVERTISEMENTS

The Journal of Laryngology and Otology

(FOUNDED in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY

G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the under-standing that they are contributed to this *journal* solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgement must be made. 2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and

wen spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc. The Harvard system of recording references should be used, e.g. GREEN, C, and BROWN, D. (1951) /. Laryng., 65, 33. Abbreviations of Journals should follow the style recommended in *World Medical Periodicals*, published by World Health Organization, 1952.

It is most important that authors should verify *personally* the accuracy of every reference before submitting a paper for publication. 3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks are provided free up to the limit of $\pounds \rangle o$ per article; beyond this authors are expected to pay half the cost. Coloured illustrations will be charged in full to authors. Blocks will normally be held by the Printers for three years after which they will be destroyed. Any author who has borne a part of the cost of his blocks is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article to HEADLEX BROTHERS. 100 Kingway three years of the appearance of the article, to HEADLEY BROTHERS, 109 Kingsway, London, WC2

5. Orders for reprints must be sent when returning galley proofs, and for this

5. Orders for reprints must be sent when returning galey proofs, and for this purpose special forms are supplied. 6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to THE EDITOR, *Journal of Laryngology*, c/o HEADLEY BROTHERS, 109 Kingsway, London, WC2.

8. The annual subscription is five guineas sterling (U.S.A. \$15) post free, and payable in advance.
9. Single copies will be on sale at 12s. 6d. each; copies of parts up to Vol. LXIII which are available may be purchased at 7s. 6d. each.
10. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, WC2.

United States of America

Orders for this *Journal* may be sent through local bookseller, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, NXC2, England. © Journal of Laryngology and Otology, 1964

CONTENTS

PAGE

The Transmission of Acoustic Energy from Air to the Receptor Organ in the Cochlea—2. L. Naftalin (Lincoln)	239
THE USE OF THE C.A.L. ULTRASONIC GENERATOR IN MÉNIÈRE'S DISEASE. G. Kossoff, A. E. Khan and R. H. Farrant (Sydney)	245
THE TREATMENT OF MÉNIÈRE'S DISEASE WITH ULTRA-SOUND. R. B. Lumsden	258
Rôle of Compression in the Ætiology of Bell's Palsy. S. N. Jain and A. P. Sharma	266
PRESBYACUSIS. T. E. T. Weston	273
EAR PHOTOGRAPHY. K. McLay (Edinburgh) and R. J. Russell (London)	287
CLINICAL RECORDS	
A Solitary Neurilemmoma Causing Paralysis of the IXth, Xth, XIth and XIIth Cranial Nerves and Sympathetic Chain. P. M. Binns (Leeds)	290
TRACHEAL STENOSIS FOLLOWING TRACHEOTOMY. P. M. Binns (Leeds)	292
Localized Laryngeal Amyloidosis, A Report of a Case with a Review of the Literature. J. C. McAlpine	
and A. D. Fuller (London)	
and A. P. Fuller (London)	296
A Case of Chondroma of the Epiglottis. A. B. N. Rao	296 315
A Case of Chondroma of the Epiglottis. A. B. N. Rao . Chronic Sinusitis with Alveolar Fistulae of Mediæval	315
A Case of Chondroma of the Epiglottis. A. B. N. Rao	-

THE LARYNGOSCOPE

A Monthly Journal devoted to the disease of EAR, NOSE AND THROAT

Official organ for the American Laryngological Rhinological and Otological Society

Price \$15.00 per year Canada \$15.50 per year Foreign \$16.00 per year

MAX A. GOLDSTEIN, M.D. FOUNDER THEODORE E. WALSH, M.D. EDITOR

640 SOUTH KINGSHIGHWAY SAINT LOUIS 10, MO.



'She gave me ears'

William Wordsworth (1807)

Another satisfied customer, perhaps. Dorothy, 'the Blessing of my later years,' made the world come alive again for Wordsworth, and a new invisible hearing aid has often done just this for the clients of Ingrams.

You can help your patients to take the first step by telling them about Ingrams, the firm of independent—and thus completely impartial—hearing aid consultants.

Ingrams will help your patient to choose from twentysix different makes. This is the largest selection in the country—and some of the most advanced aids are exclusive to Ingrams.

Wordsworth was so grateful to Dorothy that he made her immortal. We cannot guarantee that your patients will do the same but we know they will be pleased that you—like so many other E.N.T. Specialists—recommended the Ingrams service.

INGRAMS

THE INDEPENDENT HEARING AID CONSULTANTS 2 SHEPHERD ST., SHEPHERD MARKET, LONDON, W.1

ORBENIN SODIUM CLOXACILLIN

ORAL PENICILLIN ACTIVE AGAINST THE 'HOSPITAL STAPHYLOCOCCUS'

Orbenin is highly bactericidal to both resistant and sensitive staphylococci, and also to other Gram positive organisms such as streptococci and pneumococci.

Dosage: 500mg. orally (capsules or syrup) 6-hourly. Orbenin may also be given by Intramuscular injection.

Orbenin is a product of British research at BEECHAM RESEARCH LABORATORIES



COMMONWEALTH ACOUSTIC LABORATORIES

ULTRASONIC GENERATOR

FOR TREATMENT OF MENIERE'S DISEASE

The C.A.L. Ultrasonic Generator designed by The Commonwealth Acoustic Laboratories, Sydney, is manufactured in Australia by Hawker de Havilland.

It permits simple ultrasound therapy to be employed in the treatment of Meniere's disease using a proven instrument measuring only 9" x 8" x 7", incorporating a voltage stabilised, fully transistorised circuit.

The treatment probe is small enough to provide good visualisation during therapy and supplies concentrated ultrasonic energy in a focused beam free from spurious side lobes. Probe diameter is only 7.5 mm. tapering to a tip diameter of 2 mm.

Each instrument is supplied with its own calibration chart plotting manually adjustable current readings against measured acoustic output in watts from the treatment probe. Guaranteed maximum output of not less than 1 watt. The beam from all probes is carefully monitored by Schlieren visualisation during final tests.

HAWKER de HAVILLAND AUSTRALIA PTY. LTD. P.O. Box 30, Bankstown, N.S.W., Australia

220/64





Penbritin is a product of British research at BEECHAM RESEARCH LABORATORIES



IN ALL **INDICATIONS** FOR LOCAL **ANAESTHESIA** Citanest is a new local anaesthetic possessing all the excellent features of lignocaine with additional advantages. Its superiority to other agents has been repeatedly confirmed during many years of clinical trials. □ Citanest is 40% less toxic than lignocaine.

☐ Citanest is 40% less toxic than inglocanie. ☐ Citanest exerts no cumulative action on repeated injections.

□ Citanest has only slight effect on blood pressure and respiration.







The **AMPLIVOX** Model 90 transistor audiometer with **Bone Conduction & Masking**


Ideal for hospitals, hearing aid clinics, consulting rooms, industrial medical departments, schools, etc., when a full clinical instrument (such as the Amplivox Model 82) is not essential but where bone conduction measurements with masking are often required to supplement the air conduction audiogram.

- * Nine frequencies for air conduction 250-8000 c/s
- * Seven frequencies for bone conduction 250-4000 c/s
- * White noise masking up to 70 dB in 10 dB increments
- * Independent of mains supply
- * British Standard calibration
- * Maximum intensity 100 dB re BS
- * Positive battery check ensures constant output accuracy
- * Weighs only 14 lb
- # Equally suitable for rapid screening OR individual diagnostic audiometry



This new Audiometer is housed in a durable fibreglass cabinet, $12'' \times 10'' \times 8\frac{3''}{4}$, with roller blind shutter to panel and storage compartment, eliminating extra carrying case.

SEND FOR TECHNICAL LEAFLET describing this outstanding addition to the Amplivox range: Amplivox Limited, Medical Acoustic Division, 80 New Bond St., London W1



'OTOSPORIN'

The modern method of treatment in otitis externa, chronic suppurative otitis media, and infected mastoid cavities, is 'Otosporin' Drops. 'Otosporin' is an aqueous suspension containing neomycin and polymyxin B which are together effective against practically all the bacteria found in ear infections. In addition, it contains hydrocortisone to reduce inflammation and swelling, thereby providing easier access for the antibiotics.

'OTOSPORIN' BRAND Drops ISSUED IN BOTTLES OF 5 ml.

港 BURROUGHS WELLCOME & CO. (THE WELLCOME FOUNDATION LTD.) LONDON