

ON THE COHOMOLOGICAL COMPLETENESS OF q -COMPLETE DOMAINS WITH CORNERS

KAZUKO MATSUMOTO

Abstract. We prove the vanishing and non-vanishing theorems for an intersection of a finite number of q -complete domains in a complex manifold of dimension n . When q does not divide n , it is stronger than the result naturally obtained by combining the approximation theorem of Diederich-Fornaess for q -convex functions with corners and the vanishing theorem of Andreotti-Grauert for q -complete domains. We also give an example which implies our result is best possible.

Introduction

Let D be a complex manifold of dimension n and let q be an integer with $1 \leq q \leq n$. A continuous function from D to \mathbb{R} is called q -convex with corners if it is locally a maximum of a finite number of q -convex functions. In [D-F] Diederich-Fornaess proved that every q -convex function with corners defined on D can be approximated by \tilde{q} -convex functions whole on D , where $\tilde{q} := n - [n/q] + 1$ and $[x]$ denotes the integral part of x . They moreover showed that the number \tilde{q} is best possible for any (n, q) , i.e., there exist an open subset D in \mathbb{C}^n and a finite number of q -convex functions $\varphi_1, \dots, \varphi_s$ defined on D such that the function $\varphi := \max\{\varphi_1, \dots, \varphi_s\}$ cannot be approximated by $(\tilde{q} - 1)$ -convex functions.

A complex manifold D is called q -complete (resp. q -complete with corners) if D has an exhaustion function which is q -convex (resp. q -convex with corners) on D . Combining the above theorem of Diederich-Fornaess with the theorem of Andreotti-Grauert ([A-G]) it follows at once that if D is q -complete with corners then D is cohomologically \tilde{q} -complete.

Now the following problem arises naturally.

PROBLEM. *Is there a complex manifold which is q -complete with corners but not cohomologically $(\tilde{q} - 1)$ -complete?*

Received May 15, 2001.

2000 Mathematics Subject Classification: 32F10.

It is easy to find such examples if q divides n (cf. [S-V], [E-S], [M-1] and [M-2]). However, it seems that such an example is still unknown if q does not divide n .

The purpose of this article is to prove the following.

THEOREM. *Let M be a complex manifold of dimension n and let D_1, \dots, D_t be q -complete open subsets in M . Let \mathcal{F} be a coherent analytic sheaf on M such that $H^n(M, \mathcal{F}) = 0$. Then*

$$H^j(D_1 \cap \dots \cap D_t, \mathcal{F}) = 0 \quad \text{if } j \geq \widehat{q}_t.$$

Here

$$\widehat{q}_t := \min\{\widehat{q}, t(q - 1) + 1\}$$

and

$$\widehat{q} := n - \left[\frac{n-1}{q} \right] = \begin{cases} \widetilde{q} & \text{if } q \mid n \\ \widetilde{q} - 1 & \text{if } q \nmid n. \end{cases}$$

Moreover, the number \widehat{q}_t in the above theorem is best possible for any (n, q, t) . In particular, for any (n, q) there exist a finite number of q -complete open subsets D_1, \dots, D_s in \mathbb{C}^n such that $H^{\widehat{q}-1}(D_1 \cap \dots \cap D_s, \mathcal{O}) \neq 0$, where \mathcal{O} denotes the sheaf of germs of holomorphic functions on \mathbb{C}^n (see §3).

The author, in general, does not know whether the cohomologically \widehat{q} -complete set $D_1 \cap \dots \cap D_t$ in the above theorem is \widehat{q} -complete, i.e., it has a \widehat{q} -convex exhaustion function, even in the case $M = \mathbb{C}^n$.

§1. The key proposition

First we show the following proposition which is a key step to prove Theorem.

PROPOSITION 1. *Let M be a topological space, let $\{D_1, \dots, D_t\}$ be a family of open subsets in M and let \mathcal{S} be a sheaf of Abelian groups on M . Let $p \in \mathbb{N}$ be fixed and suppose that for any k with $1 \leq k \leq t - 1$ the family $\{D_1, \dots, D_t\}$ satisfies the condition*

$$C(k, p) \quad H^j(D_{i_1} \cap \dots \cap D_{i_k}, \mathcal{S}) = 0$$

for all $j \geq p$ and all $i_1, \dots, i_k \in \{1, 2, \dots, t\}$.

Then

- (1) $H^j(D_1 \cap \dots \cap D_t, \mathcal{S}) \cong H^{j+t-1}(D_1 \cup \dots \cup D_t, \mathcal{S}) \quad \text{if } j \geq p;$
- (2) $H^{p-1}(D_1 \cap \dots \cap D_t, \mathcal{S}) \twoheadrightarrow H^{p+t-2}(D_1 \cup \dots \cup D_t, \mathcal{S})$

Remark. The family $\{D_1, \dots, D_t\}$ satisfies the condition $C(k, p)$ for all k with $1 \leq k \leq t - 1$ if it satisfies only $C(t - 1, p)$.

Proof of Proposition 1. We shall prove the proposition by induction on $t \in \mathbb{N}$.

Step 1. When $t = 1$, (1) and (2) are trivial.

Step 2. When $t \geq 2$, it follows by Mayer-Vietories that the sequence

$$\begin{aligned} &H^j(D_1 \cap \dots \cap D_{t-1}, \mathcal{S}) \oplus H^j(D_t, \mathcal{S}) \\ &\longrightarrow H^j((D_1 \cap \dots \cap D_{t-1}) \cap D_t, \mathcal{S}) \\ &\longrightarrow H^{j+1}((D_1 \cap \dots \cap D_{t-1}) \cup D_t, \mathcal{S}) \\ &\longrightarrow H^{j+1}(D_1 \cap \dots \cap D_{t-1}, \mathcal{S}) \oplus H^{j+1}(D_t, \mathcal{S}) \end{aligned}$$

is exact for each j . Since $\{D_1, \dots, D_t\}$ satisfies $C(t - 1, p)$ and $C(1, p)$ by assumption, we have

$$H^j(D_1 \cap \dots \cap D_{t-1}, \mathcal{S}) = H^j(D_t, \mathcal{S}) = 0 \quad \text{if } j \geq p.$$

Therefore, if we put $E_i := D_i \cup D_t$ for $i = 1, 2, \dots, t - 1$, then

$$(3) \quad H^j(D_1 \cap \dots \cap D_t, \mathcal{S}) \cong H^{j+1}(E_1 \cap \dots \cap E_{t-1}, \mathcal{S}) \quad \text{if } j \geq p;$$

$$(4) \quad H^{p-1}(D_1 \cap \dots \cap D_t, \mathcal{S}) \twoheadrightarrow H^p(E_1 \cap \dots \cap E_{t-1}, \mathcal{S}).$$

In particular, this means that the proposition holds in the case $t = 2$.

Step 3. When $t \geq 3$, we assume that the proposition has been proved for $1, 2, \dots, t - 1$. We first show the following.

LEMMA 1. *Under the above situation, the family $\{E_1, \dots, E_{t-1}\}$ satisfies the condition $C(t - 2, p + 1)$.*

Proof. We shall prove by induction that for any l with $1 \leq l \leq t - 2$ the family $\{E_1, \dots, E_{t-1}\}$ satisfies the condition

$$\begin{aligned} &C(l, p + 1) \quad H^{j+1}(E_{i_1} \cap \dots \cap E_{i_l}, \mathcal{S}) = 0 \\ &\quad \text{for all } j \geq p \text{ and all } i_1, \dots, i_l \in \{1, 2, \dots, t - 1\}. \end{aligned}$$

By the assumption of the proposition $\{D_1, \dots, D_t\}$ satisfies $C(t-1, p)$ and particularly $C(1, p)$ and $C(2, p)$. Since the proposition holds in the case $t = 2$ we have

$$\begin{aligned} H^{j+1}(E_{i_1}, \mathcal{S}) &= H^{j+1}(D_{i_1} \cup D_t, \mathcal{S}) \\ &\cong H^j(D_{i_1} \cap D_t, \mathcal{S}) = 0 \quad \text{if } j \geq p. \end{aligned}$$

Therefore, $\{E_1, \dots, E_{t-1}\}$ satisfies $C(1, p+1)$.

Next let $2 \leq l \leq t-2$ and assume that the lemma has been proved for all m with $1 \leq m \leq l-1$. Then the family $\{E_{i_1}, \dots, E_{i_l}\}$, where $i_1, \dots, i_l \in \{1, 2, \dots, t-1\}$, also satisfies the condition $C(m, p+1)$ for all m with $1 \leq m \leq l-1$. Moreover, since $\{E_1, \dots, E_{t-1}\}$ satisfies $C(l-1, p+1)$ by the inductive hypothesis and since the proposition holds for l ,

$$\begin{aligned} H^{j+1}(E_{i_1} \cap \dots \cap E_{i_l}, \mathcal{S}) &\cong H^{(j+1)+l-1}(E_{i_1} \cup \dots \cup E_{i_l}, \mathcal{S}) \\ &= H^{j+(l+1)-1}(D_{i_1} \cup \dots \cup D_{i_l} \cup D_t, \mathcal{S}) \quad \text{if } j \geq p. \end{aligned}$$

On the other hand, $\{D_1, \dots, D_t\}$ satisfies $C(l, p)$ and $C(l+1, p)$ because $l+1 \leq t-1$. Since the proposition holds for $l+1$,

$$\begin{aligned} H^{j+(l+1)-1}(D_{i_1} \cup \dots \cup D_{i_l} \cup D_t, \mathcal{S}) &\cong H^j(D_{i_1} \cap \dots \cap D_{i_l} \cap D_t, \mathcal{S}) = 0 \quad \text{if } j \geq p. \end{aligned}$$

Hence we obtain

$$H^{j+1}(E_{i_1} \cap \dots \cap E_{i_l}, \mathcal{S}) = 0 \quad \text{if } j \geq p,$$

which proves that $\{E_1, \dots, E_{t-1}\}$ satisfies $C(l, p+1)$ for all l with $1 \leq l \leq t-2$. □

End of Proof of Proposition 1. If $t \geq 3$ and if $\{D_1, \dots, D_t\}$ satisfies $C(t-1, p)$ then $\{E_1, \dots, E_{t-1}\}$ satisfies $C(t-2, p+1)$, where $E_i := D_i \cup D_t$ for $i = 1, 2, \dots, t-1$. Therefore, by the inductive hypothesis, we have

(5) $H^{j+1}(E_1 \cap \dots \cap E_{t-1}, \mathcal{S}) \cong H^{j+t-1}(E_1 \cup \dots \cup E_{t-1}, \mathcal{S}) \quad \text{if } j \geq p;$

(6) $H^p(E_1 \cap \dots \cap E_{t-1}, \mathcal{S}) \twoheadrightarrow H^{p+t-2}(E_1 \cup \dots \cup E_{t-1}, \mathcal{S}).$

Notice here that $E_1 \cup \dots \cup E_{t-1} = D_1 \cup \dots \cup D_t$. Then we can obtain (1) and (2) by (3), (4), (5) and (6).

This completes the proof of the proposition. □

§2. Proof of Theorem

Let M be a complex manifold of dimension n , let D_1, \dots, D_t be q -complete open subsets in M and let \mathcal{F} be a coherent analytic sheaf on M such that $H^n(M, \mathcal{F}) = 0$.

Since the intersection $D_1 \cap \dots \cap D_t$ is q -complete with corners it follows from the theorem of Diederich-Fornaess and the theorem of Andreotti-Grauert that

$$H^j(D_1 \cap \dots \cap D_t, \mathcal{F}) = 0 \quad \text{if } j \geq \tilde{q}_t.$$

Here $\tilde{q}_t := \min\{\tilde{q}, t(q - 1) + 1\}$ and $\tilde{q} := n - [n/q] + 1$.

We put

$$\hat{q} := n - \left\lfloor \frac{n-1}{q} \right\rfloor = \begin{cases} \tilde{q} & \text{if } q \mid n \\ \tilde{q} - 1 & \text{if } q \nmid n. \end{cases}$$

For the proof of Theorem it is enough to prove the following.

LEMMA 2. *Under the above situation,*

$$H^j(D_1 \cap \dots \cap D_t, \mathcal{F}) = 0 \quad \text{if } j \geq \hat{q}.$$

Proof. We put $m := [n/q]$ and $r := n - mq$. Then $n = mq + r$ and $0 \leq r \leq q - 1$. We shall prove the lemma by induction on $t \in \mathbb{N}$.

First if $t \leq m$,

$$t(q - 1) + 1 \leq m(q - 1) + 1 = n - m + 1 - r = \tilde{q} - r.$$

If $q \mid n$ or $r = 0$ then $\tilde{q} - r = \tilde{q} = \hat{q}$; and if $q \nmid n$ or $r \geq 1$ then $\tilde{q} - r \leq \tilde{q} - 1 = \hat{q}$. Hence if $t \leq m$ we have $t(q - 1) + 1 \leq \hat{q} \leq \tilde{q}$ and

$$\tilde{q}_t := \min\{\tilde{q}, t(q - 1) + 1\} = t(q - 1) + 1 \leq \hat{q}.$$

Therefore, by the theorem of Diederich-Fornaess, the lemma holds if $t \leq m$.

Next if $t \geq m + 1$ and if the lemma holds for $1, 2, \dots, t - 1$, then for any k with $1 \leq k \leq t - 1$ the family $\{D_1, \dots, D_t\}$ satisfies the condition

$$C(k, \hat{q}) \quad \begin{aligned} &H^j(D_{i_1} \cap \dots \cap D_{i_k}, \mathcal{F}) = 0 \\ &\text{for all } j \geq \hat{q} \text{ and all } i_1, \dots, i_k \in \{1, 2, \dots, t\}. \end{aligned}$$

Hence by Proposition 1

$$H^j(D_1 \cap \dots \cap D_t, \mathcal{F}) \cong H^{j+t-1}(D_1 \cup \dots \cup D_t, \mathcal{F}) \quad \text{if } j \geq \hat{q}.$$

Notice here that if $t \geq m+1$ and $j \geq \widehat{q}$ then $j+t-1 \geq \widehat{q}+m \geq \widetilde{q}-1+m = n$.

Since the set $D_1 \cup \dots \cup D_t$ is open in M and since $H^n(M, \mathcal{F}) = 0$ by assumption we have $H^n(D_1 \cup \dots \cup D_t, \mathcal{F}) = 0$ (see Remark below). Therefore we obtain

$$H^j(D_1 \cap \dots \cap D_t, \mathcal{F}) = 0 \quad \text{if } j \geq \widehat{q},$$

which proves the lemma. □

Theorem is the direct result of the above lemma and the theorem of Diederich-Fornaess (cf. [D-F], §5).

Remark. By the theorem of Greene-Wu ([G-W]), a connected complex manifold of dimension n is n -complete if and only if it is noncompact. Therefore, if D is noncompact complex manifold of dimension n then by the theorem of Andreotti-Grauert $H^n(D, \mathcal{F}) = 0$ for any coherent analytic sheaf \mathcal{F} on D . It is obvious that if $H^n(M, \mathcal{F}) = 0$ then $H^n(D, \mathcal{F}) = 0$ for any connected (and not necessarily noncompact) component D of M .

§3. Example

As in Section 2 we put $n = mq + r$. In \mathbb{C}^n , consider the complex linear subspaces defined by

$$L_i := \{(z_1, \dots, z_n) \in \mathbb{C}^n \mid z_{(i-1)q+1} = \dots = z_{iq} = 0\}$$

and put $D_i := \mathbb{C}^n \setminus L_i$ for $i = 1, 2, \dots, m$. Then each D_i is q -complete but not $(q - 1)$ -complete (cf. [W]). If $q \nmid n$ or $r \geq 1$, we moreover put

$$L_{m+1} := \{(z_1, \dots, z_n) \in \mathbb{C}^n \mid z_{mq+1} = \dots = z_n = 0\}$$

and $D_{m+1} := \mathbb{C}^n \setminus L_{m+1}$. Then D_{m+1} is r -complete and particularly q -complete because $r < q$.

The number \widehat{q}_t in Theorem is best possible for any (n, q, t) , where

$$\widehat{q}_t := \min\{\widehat{q}, t(q - 1) + 1\} = \begin{cases} t(q - 1) + 1 & \text{if } t \leq m \\ \widehat{q} & \text{if } t > m \end{cases}$$

and

$$\widehat{q} := n - \left\lfloor \frac{n - 1}{q} \right\rfloor = \begin{cases} n - m + 1 & \text{if } q \mid n \\ n - m & \text{if } q \nmid n. \end{cases}$$

In fact, we have the following.

EXAMPLE. Under the above notations, $H^{t(q-1)}(D_1 \cap \dots \cap D_t, \mathcal{O}) \neq 0$ for $t = 1, 2, \dots, m$. Moreover, $H^{n-m-1}(D_1 \cap \dots \cap D_{m+1}, \mathcal{O}) \neq 0$ if $q \nmid n$.

In the example above, \mathcal{O} denotes the sheaf of germs of holomorphic functions on \mathbb{C}^n . The example is a part of the following.

PROPOSITION 2. Let $\alpha_0, \alpha_1, \dots, \alpha_t$ and n_0 be integers such that $0 = \alpha_0 < \alpha_1 < \dots < \alpha_t = n_0 \leq n$. In \mathbb{C}^n , consider the complex linear subspaces defined by

$$L_i := \{(z_1, \dots, z_n) \in \mathbb{C}^n \mid z_{\alpha_{i-1}+1} = z_{\alpha_{i-1}+2} = \dots = z_{\alpha_i} = 0\}$$

and put $D_i := \mathbb{C}^n \setminus L_i$ for $i = 1, 2, \dots, t$. Then

$$\begin{cases} H^{n_0-t}(D_1 \cap \dots \cap D_t, \mathcal{O}) \neq 0 \\ H^j(D_1 \cap \dots \cap D_t, \mathcal{O}) = 0 \end{cases} \quad \text{if } j \geq n_0 - t + 1.$$

Proof. Since $\text{codim } L_i \leq n_0 - (t - 1)$ each D_i is at least $(n_0 - t + 1)$ -complete. Hence if we put $p := n_0 - t + 1$ then $H^j(D_i, \mathcal{O}) = 0$ for all $j \geq p$ and all i with $1 \leq i \leq t$.

We shall now prove by induction that for any k with $1 \leq k \leq t - 1$ the family $\{D_1, \dots, D_t\}$ satisfies the condition

$$C(k, p) \quad \begin{aligned} &H^j(D_{i_1} \cap \dots \cap D_{i_k}, \mathcal{O}) = 0 \\ &\text{for all } j \geq p \text{ and all } i_1, \dots, i_k \in \{1, 2, \dots, t\}. \end{aligned}$$

First $\{D_1, \dots, D_t\}$ satisfies $C(1, p)$. Next if it satisfies $C(k - 1, p)$ where $k \geq 2$, it follows from Proposition 1 that

$$H^j(D_{i_1} \cap \dots \cap D_{i_k}, \mathcal{O}) \cong H^{j+k-1}(D_{i_1} \cup \dots \cup D_{i_k}, \mathcal{O}) \quad \text{if } j \geq p.$$

Since $D_{i_1} \cup \dots \cup D_{i_k} = \mathbb{C}^n \setminus (L_{i_1} \cap \dots \cap L_{i_k})$ and since $\text{codim } (L_{i_1} \cap \dots \cap L_{i_k}) \leq n_0 - (t - k) = p + k - 1$, the set $D_{i_1} \cup \dots \cup D_{i_k}$ is at least $(p + k - 1)$ -complete. Hence for any k with $1 \leq k \leq t - 1$ we have

$$H^j(D_{i_1} \cap \dots \cap D_{i_k}, \mathcal{O}) = 0 \quad \text{if } j \geq p,$$

which implies that $\{D_1, \dots, D_t\}$ satisfies $C(t - 1, p)$.

Therefore, by Proposition 1 we obtain

$$(7) \quad H^j(D_1 \cap \dots \cap D_t, \mathcal{O}) \cong H^{j+t-1}(D_1 \cup \dots \cup D_t, \mathcal{O}) \quad \text{if } j \geq p;$$

$$(8) \quad H^{p-1}(D_1 \cap \dots \cap D_t, \mathcal{O}) \twoheadrightarrow H^{p+t-2}(D_1 \cup \dots \cup D_t, \mathcal{O}).$$

On the other hand,

$$\begin{cases} H^{n_0-1}(D_1 \cup \cdots \cup D_t, \mathcal{O}) \neq 0 \\ H^j(D_1 \cup \cdots \cup D_t, \mathcal{O}) = 0 \end{cases} \quad \text{if } j \geq n_0$$

because $D_1 \cup \cdots \cup D_t = \mathbb{C}^n \setminus (L_1 \cap \cdots \cap L_t)$ and $\text{codim}(L_1 \cap \cdots \cap L_t) = n_0$. Since $p := n_0 - t + 1$ we thus obtain

$$\begin{cases} H^{n_0-t}(D_1 \cap \cdots \cap D_t, \mathcal{O}) \neq 0 \\ H^j(D_1 \cap \cdots \cap D_t, \mathcal{O}) = 0 \end{cases} \quad \text{if } j \geq n_0 - t + 1.$$

This completes the proof of the proposition. \square

Acknowledgement. The author expresses her sincere thanks to the referee for the useful comment.

REFERENCES

- [A-G] A. Andreotti and H. Grauert, *Théorèmes de finitude pour la cohomologie des espaces complexes*, Bull. Soc. Math. France, **90** (1962), 193–259.
- [D-F] K. Diederich and J. E. Forneaess, *Smoothing q -convex functions and vanishing theorems*, Invent. Math., **82** (1985), 291–305.
- [E-S] M. G. Eastwood and G. V. Suria, *Cohomologically complete and pseudoconvex domains*, Comment. Math. Helv., **55** (1980), 413–426.
- [G-W] R. E. Greene and H. Wu, *Embedding of open Riemannian manifolds by harmonic functions*, Ann. Inst. Fourier (Grenoble), **25** (1975), 215–235.
- [M-1] K. Matsumoto, *Pseudoconvex domains of general order in Stein manifolds*, Mem. Fac. Sci. Kyushu Univ., **43** (1989), 67–76.
- [M-2] ———, *Boundary distance functions and q -convexity of pseudoconvex domains of general order in Kähler manifolds*, J. Math. Soc. Japan, **48** (1996), 85–107.
- [S-V] G. Sorani and V. Villani, *q -complete spaces and cohomology*, Trans. Amer. Math. Soc., **125** (1966), 432–448.
- [W] K. Watanabe, *Pseudoconvex domains of general order and vanishing cohomology*, Kobe J. Math., **10** (1993), 107–115.

Department of Applied Mathematics
Osaka Women's University
Daisen-cho, Sakai 590-0035
Japan
kazuko@appmath.osaka-wu.ac.jp