

## ERRATUM

# *Anobothrus amourouxi* sp. nov., a new species of Ampharetidae (Polychaeta) from the Capbreton Canyon (Bay of Biscay, NE Atlantic Ocean) – ERRATUM

PAULO BONIFÁCIO, NICOLAS LAVESQUE, GUY BACHELET AND JULIO PARAPAR

doi: 10.1017/S0025315414002094, published by Cambridge University Press, 9 February 2015

The 'Key for Identification of the Species of the Genus *Anobothrus* in the World' has been corrected as follows:

- |  |                           |   |                              |
|--|---------------------------|---|------------------------------|
| 1. No paleae.....  | 2                         | diameter, 13 AU.....  |                              |
| - Paleae present.....  | 3                         | .....   | <i>A. rubropaleatus</i>      |
| 2. Modified notopodia with hirsute tips notochaetae.....   |                           | 9. Modified notopodia without notochaetae with hirsute tips .....   | 10                           |
| .....  | <i>A. fimbriatus</i>      | - Only the modified notopodia with notochaetae with hirsute tips.....   | 11                           |
| - All notopodia without hirsute tips notochaetae.....  | <i>A. apaleatus</i>       | - All notopodia with notochaetae with hirsute tips.....   |                              |
| 3. 3 pairs of branchiae .....  | 4                         | .....   | <i>A. gracilis</i>           |
| - 4 pairs of branchiae.....  | 6                         | 10. 3 teeth on uncini from TU1 (lateral view), paleae abruptly tapering to delicate tip .....   | <i>A. pseudoampharete</i>    |
| 4. Modified notopodia on TU8.....  | 5                         | - 8–9 teeth on uncini from TU1 (lateral view), diameter of all branchiophores more or less equal, less than 10 paleae, very conspicuous stout and long paleae .....   | <i>A. paleodiscus</i>        |
| - Modified notopodia on TU9.....   | <i>A. flabelligerulus</i> | - 5 teeth on uncini from TU1 (lateral view), fourth pair of branchiophores two times slimmer and shorter than others reduced and their branchiostyles many times shorter than others, paleae longer than best-developed notochaeta, gradually tapering..... | <i>A. patersoni</i>          |
| 5. Notochaetae present on fused segments II- III, prostomium <i>Ampharete</i> -type, branchiae forming transversal line .....  | <i>A. laubieri</i>        | - 5 teeth on uncini from TU1 (lateral view), inner and middle pairs of branchiophores shorter and slimmer than others .....   | <i>A. mironovi</i>           |
| .....  |                           | .....   |                              |
| - Notochaetae absent on fused segments II-III, prostomium conical, wide gap between groups of branchiae .....  | <i>A. dayi</i>            | 11. 6 teeth on uncini from TU1 (lateral view), surface of branchiostyle smooth, thoracic arrangement (fused SII-III with paleae from SII and without notochaetae from SIII).....  | <i>A. glandularis</i>        |
| .....  |                           | - 4–5 teeth on uncini from TU1 (lateral view), surface of all branchiostyles papillated, thoracic arrangement (fused SII-III with paleae from SII and notochaetae from SIII) .....  | <i>A. antarctica</i>         |
| 6. Modified notopodia on TU6.....  | <i>A. bimaculatus</i>     | - 6–7 teeth on uncini from TU1 (lateral view), surface of inner pair of branchiostyle with transversal ciliated ridges, thoracic arrangement (fused SII-III with paleae from SII and with reduced notopodia without notochaetae from SIII).....             | <i>A. amourouxi</i> sp. nov. |
| - Modified notopodia on TU7.....   | <i>A. mancus</i>          | .....   |                              |
| - Modified notopodia on TU8.....   | 7                         | .....   |                              |
| - Modified notopodia on TU9.....   | <i>A. paleatus</i>        | .....   |                              |
| 7. Circular band on TU1.....   | <i>A. patagonicus</i>     | .....   |                              |
| - Circular band on TU2 (modified notochaetae without hirsute tips) .....   | 8                         | .....   |                              |
| - Circular band on TU3 .....   | 9                         | .....   |                              |
| 8. Paleae colourless, fine, with base slimmer than (or equal to) most developed notochaetae, outer pairs of branchiae distinctly narrower than inner, 12–13 AU ..... | <i>A. wilhelmi</i>        | .....   |                              |
| - Paleae reddish, stout, with base stouter than most developed notochaetae, branchiae with almost the same   |                           | .....   |                              |