

EPP522

Evaluating Cognitive Bias in Psychosis: A Novel Approach Using Large Language Models on Spanish Speech Data

E. Gutierrez Alvarez^{1,2}, A. Barajas^{3,4*} and R. Ayesa^{5,6}

¹Universidad Politécnica de Madrid, NEBULA group, Madrid, Spain; ²MIT linQ - Massachusetts Institute of Technology, Cambridge, United States; ³Department of Clinical and Health Psychology, Autonomous University of Barcelona; ⁴Government of Catalonia, Serra Húnter Programme, Barcelona; ⁵Mental Illness Research Group, Valdecilla Biomedical Research Institute (IDIVAL), Santander and ⁶Biomedical Research Center in Mental Health Network (CIBERSAM), Health Institute Carlos III, Madrid, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2025.771

Introduction: Jumping to conclusions (JTC) is a cognitive bias strongly involved in the genesis of psychotic symptoms. Accurate evaluation of JTC is crucial for early intervention and treatment planning. However, traditional assessment methods are time-consuming and subject to human error. This study leverages state-of-the-art Large Language Models (LLMs) to evaluate JTC in a unique Spanish population database collected through the DISCOURSE protocol at the Instituto de Investigación Marqués de Valdecilla (IDIVAL).

Objectives: Our primary objectives were to:


1. Assess the efficacy of LLMs in evaluating JTC bias from transcribed speech.
2. Compare different LLM models and prompting techniques for optimal performance.
3. Explore the potential of AI-assisted cognitive bias evaluation in clinical settings.

Methods: We utilized a database of approximately 170 participants, including patients, controls, and relatives, collected through the DISCOURSE protocol. This protocol is particularly valuable as it includes tasks designed to elicit JTC behaviors, such as ambiguous picture interpretation. Audio recordings were automatically transcribed using two speech-to-text algorithms and manually revised for accuracy.

We investigated various LLM models ("gpt4o", "claude-sonnet-3.5", "llama3", "gemini pro") and experimented with different prompting techniques, including instruction combinations and reasoning scratchpads (Chain of Thoughts).

Results: Our evaluation has provided valuable insights into the potential of LLMs for assessing Jumping to Conclusions (JTC) bias. We observed varying degrees of effectiveness across different LLM models in identifying JTC behaviors from transcribed speech, with some showing promise in capturing subtle linguistic cues. Prompting techniques, particularly Chain of Thought reasoning, demonstrated potential in enhancing the models' analytical capabilities. Given our Spanish-language database, we gained important insights into LLM performance in non-English contexts. Error analysis identified common limitations, informing future refinements. Preliminary findings suggested performance variations across demographic subgroups, highlighting areas for further investigation.

Image 1:



Conclusions: This study represents a step towards integrating AI and automation into clinical workflows for psychosis evaluation and treatment. The understanding of the ability of LLMs to assess JTC from speech samples could significantly enhance the objectivity of cognitive bias evaluations. These findings lay the groundwork for future research exploring the integration of AI in psychosocial interventions for psychosis, including potential applications in cognitive remediation, metacognitive training, and personalized treatment planning.

Disclosure of Interest: None Declared

EPP525

Pathways to care for patients with schizophrenia and severe social impairment

I.-M. Mølstrøm

Psychiatry East, Region Sealand Psychiatry, Research Unit for Early Detection and Psychopathology, Roskilde, Denmark

doi: 10.1192/j.eurpsy.2025.772

Introduction: This presentation sheds light on a group of patients with schizophrenia that are often overlooked. These minority groups include homeless patients and patients with very poor social functioning. Homeless patients with schizophrenia often have complex problems and are difficult to diagnose and treat. While many resources are allocated to early detection and prevention of psychosis programs, few resources are spent on treating the patients who have fallen through the cracks of society and various support systems. These patients are undeniably among the most severely ill psychiatric patients of our time. Patients who are not homeless (domiciled patients) with schizophrenia and very poor social functioning are also often challenging to treat, and they often end up living an isolated existence in their own homes.