

- implications for prevention and management. *Am J Med* 1993;94:313-328.
6. Ayliffe GAJ. The progressive intercontinental spread of methicillin-resistant *Staphylococcus aureus*. *Clin Infect Dis* 1997;24(suppl 1):S74-S79.
  7. Boyce JM, Jackson MM, Pugliese G, Batt MD, Fleming D, Garner JS, et al. Methicillin-resistant *Staphylococcus aureus*: a briefing for acute care hospitals and nursing facilities. *Infect Control Hosp Epidemiol* 1994;15:105-115.
  8. Crossley K, Landesman B, Zaske D. An outbreak of infections caused by strains of *Staphylococcus aureus* resistant to methicillin and aminoglycosides, II: epidemiologic studies. *J Infect Dis* 1979;139:280-287.
  9. Duckworth GJ. Diagnosis and management of methicillin resistant *Staphylococcus aureus* infection. *BMJ* 1993;307:1049-1052.
  10. Hershov RC, Khayr WF, Smith NL. A comparison of clinical virulence of nosocomially acquired methicillin-resistant and methicillin-sensitive *Staphylococcus aureus* infections in a university hospital. *Infect Control Hosp Epidemiol* 1992;13:587-593.
  11. Boyce JM. Methicillin-resistant *Staphylococcus aureus*: detection, epidemiology and control measures. *Infect Dis Clin North Am* 1989;3:901-913.
  12. Nicolle LE, Bialkowska-Hobrzanska H, Romance L, Harry VS, Parker S. Clonal diversity of methicillin-resistant *Staphylococcus aureus* in an acute care institution. *Infect Control Hosp Epidemiol* 1992;13:33-37.
  13. Dammann TA, Wiens RM, Taylor GD. Methicillin-resistant *Staphylococcus aureus*: identification of a community outbreak by monitoring of hospital isolates. *Can J Public Health* 1988;79:312-314.
  14. Romance L, Nicolle LE, Ross J, Law B. An outbreak of methicillin-resistant *Staphylococcus aureus* in a pediatric hospital. How it got away and how we caught it. *Can J Infect Control* 1991;6:11-13.
  15. Taylor G, Kirkland T, Kowalewska-Grochowska K, Wang Y. A multistrain cluster of methicillin-resistant *Staphylococcus aureus* based in a native community. *Can J Infect Dis* 1990;1:121-126.
  16. Laboratory for Disease Control. Methicillin-resistant *Staphylococcus aureus*—Canada. *Canada Diseases Weekly Report* 1984;10:26-28.
  17. Embil J, Ramotar K, Romance L, Alfa M, Conly J, Cronk S, et al. Methicillin-resistant *Staphylococcus aureus* in tertiary care institutions on the Canadian prairies, 1990-1992. *Infect Control Hosp Epidemiol* 1994;15:646-651.
  18. Tokue Y, Shoji S, Satoh K, Watanabe A, Motomiya M. Comparison of a polymerase chain reaction assay and a conventional microbiologic method for detection of methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 1992;36:6-9.
  19. Geha DJ, Uhl JR, Gustaferrero CA, Persing DH. Multiplex PCR for identification of methicillin-resistant staphylococci in the clinical laboratory. *J Clin Microbiol* 1994;32:1768-1772.
  20. Tenover FC, Arbeit RD, Goering RV, Michelson PA, Murray BE, Persing DH, et al. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J Clin Microbiol* 1995;33:2233-2239.
  21. Hicks NR, Moore EP, Williams EW. Carriage and community treatment of methicillin resistant *Staphylococcus aureus*: what happens to colonized patients after discharge? *J Hosp Infect* 1991;19:17-24.
  22. Mylotte JM. Control of methicillin-resistant *Staphylococcus aureus*: the ambivalence persists. *Infect Control Hosp Epidemiol* 1994;15:73-77.
  23. Klimek JJ, Marsik FJ, Bartlett RC, Weir B, Shea P, Quintiliani R. Clinical, epidemiologic and bacteriologic observations of an outbreak of methicillin-resistant *Staphylococcus aureus* at a large community hospital. *Am J Med* 1976;61:340-345.
  24. Lugeon C, Blanc DS, Wenger A, Francioli P. Molecular epidemiology of methicillin-resistant *Staphylococcus aureus* at a low-incidence hospital over a 4-year period. *Infect Control Hosp Epidemiol* 1995;16:260-267.
  25. Hiramatsu K, Hanaki H, Ino T, Yabuta K, Oguri T, Tenover FC. Methicillin-resistant *Staphylococcus aureus* clinical strain with reduced vancomycin susceptibility. *J Antimicrob Chemother* 1997;40:135-136.
  26. Centers for Disease Control and Prevention. *Staphylococcus aureus* with reduced susceptibility to vancomycin—United States, 1997. *MMWR* 1997;46:765-766.

## Integrity of Latex Gloves

Gina Pugliese, RN, MS  
Martin S. Favero, PhD

Researchers from John Hopkins school of medicine recently conducted a study to assess the loss of integrity of the latex-glove barrier during use in the emergency department (ED) setting. A prospective observational study was conducted in an urban ED and level I trauma center. Procedures performed by ED healthcare workers (HCWs) were directly observed, timed, and categorized. The gloves used by the HCWs were collected and subjected to the standard US FDA leak test.

The results showed that 99 (7.9%) of 1,254 pairs of gloves used for observed procedures leaked, compared with 2 of 200 unworn control pairs (1.0%) and 12 of 300 pairs that were worn but not used (4.0%). Leak rates varied

by manufacturer and were higher for gloves worn 20 minutes or longer (13.7%;  $P=.015$ ), used for four or more procedures (50%;  $P<.01$ ), or used for critical-care procedures (23.5%;  $P<.01$ ). Sixty-six of an additional 325 pairs of gloves collected from unobserved critical-care procedures (20.3%) leaked.

The researchers concluded that loss of glove integrity occurs during the performance of ED procedures, subjecting the HCW to possible infectious-fluid exposure. Risk of glove perforation increases with duration of wear, number of procedures performed, and the performance of critical-care procedures.

FROM: Hansen KN, Korniewicz DM, Hexter DA, Kornilow JR, Kelen GD. Loss of glove integrity during emergency department procedures. *Ann Emerg Med* 1998;31(1):65-72.