

Temperature Evolution of Molecular Clouds in the Central Molecular Zone

Nico Krieger¹, Jürgen Ott², Fabian Walter^{1,2}, J. M. Diederik
Kruijssen^{3,1}, Henrik Beuther¹ and the SWAG team

¹Max Planck Institute for Astronomy, Heidelberg, Germany

²National Radio Astronomy Observatory, Socorro, NM, USA

³Astronomisches Rechen-Institut, Universität Heidelberg, Germany

Abstract. We infer the absolute time dependence of kinematic gas temperature along a proposed orbit of molecular clouds in the Central Molecular Zone (CMZ) of the Galactic Center (GC). Ammonia gas temperature maps are one of the results of the “Survey of Water and Ammonia in the Galactic Center” (SWAG, PI: J. Ott); the dynamical model of molecular clouds in the CMZ was taken from Kruijssen *et al.* (2015). We find that gas temperatures increase as a function of time in both regimes before and after the cloud passes pericenter on its orbit in the GC potential. This is consistent with the recent proposal that pericenter passage triggers gravitational collapse. Other investigated quantities (line width, column density, opacity) show no strong sign of time dependence but are likely dominated by cloud-to-cloud variations.

Keywords. ISM: clouds, ISM: evolution, ISM: structure, radio lines: ISM, stars: formation

The Central Molecular Zone (CMZ) in the Galactic Center (GC) contains large amounts of dense molecular gas, but observed star formation rates are lower than expected by a factor of 10 – 100 (Longmore *et al.* 2013a). Part of this gas is projected onto an ∞ -like shape and has been modeled as open-ended streams wrapping around the GC (Longmore *et al.* 2013b; Kruijssen *et al.* 2015). The streams repeatedly pass close (60 pc) to the GC where cloud collapse is expected to be triggered by tidal interactions. Downstream of the near side pericenter, a sequence of advancing star formation (SF) tracers is present that may indicate a sequence of progressing star formation states (Longmore *et al.* 2013b). We aim to understand the conditions of molecular gas, which is the fuel of SF, at the different stages of the sequence.

In the “Survey of Water and Ammonia in the Galactic Center” (SWAG, cf. overview by Ott *et al.* in this volume) targeted the CMZ at six meta-stable ammonia lines among a variety of other spectral lines. Fig. 1 (left panel) shows a peak intensity map of NH₃ (1,1) highlighting the ring-like gas streams and other relevant objects like molecular clouds and star clusters. The right panel of Fig. 1 represents the same view in kinetic gas temperature T₂₄ which can be derived from the relative line fluxes through hyperfine structure fits of NH₃ (2,2) and NH₃ (4,4) on a pixel by pixel basis (Krieger 2016a, 2016b).

The kinematic model of Kruijssen *et al.* (2015, see marked orbit in fig. 1, left) allows us to assign time stamps to CMZ clouds if their 3D position (longitude, latitude, line-of-sight velocity) is known. Thus, mapped quantities like temperature can be plotted as a function of time as in Fig. 2 for kinetic gas temperature T₂₄. Temperatures increase in the dust ridge (-1.7 to -1.3 Myr) and near side pericenter passage (+1.8 to +2.0 Myr) as shown by the fits at consistent slopes of ~ 46 K/Myr and ~ 42 K/Myr. At 0.0 Myr (far side pericenter), we find a similar trend, although the statistical basis is weak because of the short sampled time range and few measurements due to low SNR.




Figure 1. *left:* Overview of the central CMZ. The NH_3 (1,1) peak flux map extends from -0.65° to $+0.80^\circ$ in Galactic longitude and $\pm 0.35^\circ$ in Galactic latitude. Gas streams are highlighted (dark band) and overplotted with time (white) since the far side pericenter passage in the model of Kruijssen *et al.* (2015). Pericenter passages occur at ± 2.0 Myr and 0.0 Myr. The “dust ridge” spans between the “Brick” and Sgr B2. *right:* NH_3 (2,2)-(4,4) kinetic temperature (T_{24} in Kelvin) map of the CMZ derived from pixel-by-pixel fitting of the ammonia hyperfine structure (Krieger 2016a, 2016b). Blanked regions cannot be fitted due to low signal-to-noise ratios.




Figure 2. Kinetic ammonia temperature T_{24} [NH_3 (2,2) to (4,4)] as function of time since far side pericenter passage. Individual temperature measurements are overplotted with data density contours, medians of bins of 0.01 Myr width (crosses) and linear fits to these medians.

We also find quantitatively similar results for kinetic ammonia temperatures T_{45} [NH_3 (4,4) to (5,5)] and T_{36} [NH_3 (3,3) to (6,6)]. The physical reasons for differences between temperature slopes of different temperature tracers are currently unknown but likely reflect different conditions in the clouds.

Other mapped quantities (column density, opacity, line width) are dominated by variations of cloud structure and do not show consistent evolution with time.

References

- Krieger, N. 2016a, Master thesis, University of Heidelberg
 Krieger, N., Ott, J., Walter, F., *et al.* 2016b, in prep.
 Kruijssen, J. M. D., Dale, J. E., & Longmore, S. N. 2015, *MNRAS*, 447, 1059
 Longmore, S. N., Bally, J., Testi, L. *et al.* 2013a, *MNRAS*, 429, 987
 Longmore, S. N., Kruijssen, J. M. D., Bally *et al.* 2013b, *MNRAS*, 433, L15