

Spectral analyses of WC stars in the LMC

Götz Gräfenner, Wolf-Rainer Hamann, and Lars Koesterke

Department of Physics, University of Potsdam, Germany

Abstract. Spectra of seven WC stars in the LMC are taken with the *HST-FOS*. They are analyzed by means of non-LTE models for spherically expanding atmospheres with complex model atoms of helium, carbon and oxygen taken into account. We find stellar luminosities in the range from $10^{5.1-5.6} L_{\odot}$. The stellar temperature T_{\star} , defined as the effective temperature related to the stellar core radius, is of the order of 100 kK. The atmospheric compositions show carbon mass fractions of about 0.4 and oxygen mass fractions in the range 0.1–0.5.

1. Analyses

In the observed sample all single WC and WO stars from the catalog of Breyssacher (1981) are considered. Due to the known distance and low interstellar extinction of the LMC, reliable stellar parameters can be obtained by spectral analyses. The results are presented in Table 1. A detailed description of the observations and methods of analysis is given in Gräfenner *et al.* (1998).

Table 1. Model parameters: stellar luminosity L_{\star} ; core radius R_{\star} ; effective core temperature T_{\star} ; mass loss rate \dot{M} ; terminal velocity v_{∞} ; surface mass fraction of carbon β_{C} and oxygen β_{O} .

star	type	$\log L_{\star}$ (L_{\odot})	R_{\star} (R_{\odot})	T_{\star} (kK)	$\log \dot{M}$ ($M_{\odot} \text{ yr}^{-1}$)	v_{∞} (km s^{-1})	β_{C}	β_{O}
Br 7	WC 4	5.29	1.65	94.6	-4.16	2300	0.4	0.2
Br 8	WC 4	5.13	1.13	104.2	-4.06	2300	0.4	0.3
Br 10	WC 4	5.62	2.52	92.5	-3.80	2800	0.5	0.2
Br 43	WC 4	5.26	1.32	103.9	-3.91	2600	0.4	0.3
Br 50	WC 4	5.55	2.25	94.0	-3.72	2800	0.4	0.1
Br 74	WC 4	5.13	1.13	104.2	-4.06	2300	0.4	0.3
Br 93	WO 4	5.10	1.17	101.0	-4.41	3600	0.4	0.5




Figure 1. *Left:* Surface abundances are compared to central abundances of core helium burning stars with different values of the $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ rate; *Right:* Mass-loss rates are compared to a wind momentum-luminosity relation with a force multiplier of $\alpha' = 0.72$. The luminosities L_{nuc} are corrected for the flux of kinetic wind energy.

2. Surface abundances, mass-loss rates

Since for massive stars the central temperature in the helium burning core is nearly independent of the stellar mass, there exists a relation between central helium and carbon mass fractions. The shape of this relation — in particular the maximum value for carbon — mainly depends on the ratio of the $^{12}\text{C} + \alpha \rightarrow ^{16}\text{O}$ rate to the process $3\alpha \rightarrow ^{12}\text{C}$. Because of their large mass-loss rates WC stars reveal layers at their surface which have been part of a convective helium burning core and show the corresponding abundances. As is shown in Figure 1 the derived abundances are in good agreement with the high $^{12}\text{C} + \alpha \rightarrow ^{16}\text{O}$ rate from Caughlan *et al.* (1985).

The right panel of Figure 1 shows that the WC mass-loss rates obey a wind momentum-luminosity relation analogous to the one which was derived by Puls *et al.* (1996) from the theory of line-driven winds for O stars. The fact that our homogeneous sample of WR stars obeys such a relation over a wide range of luminosities is a hint to the line-force as driving mechanism for WR winds.

References

- Breysacher, J. 1981, *A&AS* 43, 203
 Caughlan, G.R., Fowler, W.A., Harris, M.J., Zimmermann, B.A. 1985, *Atomic Data Nuc. Data Tables* 32, 197
 Gräfener, G., Hamann, W.-R., Hillier, D.J., Koesterke, L. 1998, *A&A* 329, 190
 Puls, J., Kudritzki, R.-P., Herrero, A., Pauldrach, A.W.A., Haser, S.M., Lennon, D.J., Gabler, R., Voels, S.A., Vílchez, J.M., Wachter, S., Feldmeier, A. 1996, *A&A* 305, 171