

BULLETIN

OF

ENTOMOLOGICAL RESEARCH

**ISSUED BY THE ENTOMOLOGICAL RESEARCH
COMMITTEE (TROPICAL AFRICA), APPOINTED
BY THE COLONIAL OFFICE.**

EDITOR: THE SCIENTIFIC SECRETARY.

VOL. I.

LONDON:

SOLD BY


LONGMANS, GREEN, & Co., 39 PATERNOSTER ROW, E.C.,

AND

TAYLOR & FRANCIS, RED LION COURT, FLEET STREET, E.C.

1910-1911

ALERE FLAMMAM.



PRINTED BY TAYLOR AND FRANCIS,
RED LION COURT, FLEET STREET.

CONTENTS.

ORIGINAL ARTICLES.

| | Page |
|--|------|
| AUSTEN, E. E. | |
| A new genus and two new species of African Fruit-flies | 71 |
| A new species of <i>Cordylobia</i> | 79 |
| On Tabanidæ collected in North-Eastern Rhodesia and Katanga, Congo Free State, by Dr. S. and Mr. S. A. Neave, with Descriptions of new species | 275 |
| A new species of <i>Tabanus</i> from the Anglo-Egyptian Sudan | 291 |
| A new species of Tsetse-fly allied to <i>Glossina palpalis</i> , Rob.-Desv. | 294 |
| BOYCE, Sir RUBERT. | |
| The prevalence, distribution and significance of <i>Stegomyia fasciata</i> , F. (= <i>calopus</i> , Mg.), in West Africa | 233 |
| DAVEY, Dr. J. B. | |
| Notes on the habits of <i>Glossina fusca</i> | 143 |
| DRAKE-BROCKMAN, Dr. R. E. | |
| Notes on the Blood-sucking Diptera met with in Eastern and South-Eastern Abyssinia | 55 |
| DUDGEON, GERALD C. | |
| Notes on two West African Hemiptera injurious to Cocoa | 59 |
| On the Parasites of two species of West African wild Silkworms | 83 |
| West African Hemiptera injurious to Cocoa | 177 |

| | Page |
|---|------|
| GRAHAM, Dr. W. M. | |
| The Study of Mosquito Larvæ | 51 |
| On West African Trypetidæ (Fruit-flies) | 161 |
| GREEN, E. ERNEST. | |
| Remarks on Coccidæ from Uganda | 201 |
| HALL, P. E. | |
| Notes on the movements of <i>Glossina morsitans</i> in the Lundazi District, North-Eastern Rhodesia | 183 |
| HUGHES, FRANK. | |
| The chemical analysis of the secretory covering of <i>Dactylopius perniciosus</i> | 141 |
| KING, HAROLD H. | |
| Some observations on the bionomics of <i>Tabanus ditentatus</i> , Macquart, and <i>Tabanus kingi</i> , Austen | 265 |
| Some observations on the bionomics of <i>Tabanus par</i> , Walk., and <i>Tabanus teniola</i> , P. de B. | 99 |
| NEAVE, S. A. | |
| Report on a Journey to the Luangwa Valley, North-Eastern Rhodesia, from July to September, 1910 | 303 |
| NEWSTEAD, ROBERT. | |
| On Scale Insects (Coccidæ), &c., from the Uganda Protectorate | 63 |
| Some further observations on the Scale Insects of the Uganda Pro- tectorate | 185 |
| NEWSTEAD, R., and WILLCOCKS, F. C. | |
| Characters and general morphology of the Lebbek Scale | 138 |
| OUDEMANS, Dr. A. C. | |
| A short survey of the more important Families of Acari | 105 |
| ROTHSCHILD, Hon. N. CHARLES. | |
| A synopsis of the Fleas found on <i>Mus norvegicus</i> (= <i>decumanus</i>), <i>Mus rattus</i> (= <i>alexandrinus</i>), and <i>Mus musculus</i> | 89 |
| SANDERSON, Dr. MEREDITH. | |
| Notes on <i>Glossina fusca</i> , Walk., in North Nyasa | 299 |

| | Page |
|---|------|
| SHARPE, Sir ALFRED. | |
| Notes on the habits of <i>Glossina morsitans</i> in Nyasaland and the adjoining Territories | 173 |
| WESCHÉ, W. | |
| On the Larval and Pupal Stages of West African Culicidæ | 7 |
| WILLCOCKS, F. C. | |
| A Mealy Bug injurious to the Lebbek trees of Cairo | 121 |

MISCELLANEOUS.

| | |
|---|--------------|
| Collections received | 87, 159, 231 |
| Current notes | 155, 225 |
| Foreword | 1 |
| <i>Glossina</i> , an insect enemy of, in Dahomey | 219 |
| <i>Glossina morsitans</i> , concerning | 147 |
| Mosquito larvæ and their natural enemies | 213 |
| Sleeping Sickness Commission in Uganda, some entomological observations made by the .. | 221 |
| Sleeping Sickness infection in the Luangwa Valley, investigation into the origin of | 179 |
| Tsetse-flies, a method of destroying | 85 |
| Tsetse-flies (<i>Glossina</i>), an investigation into the habits and distribution of the Cattle | 152 |
| Tsetse-flies, investigation into the habits and distribution of | 319 |
| Tsetse-fly and Cattle disease in the Nyasaland Protectorate, précis of reports submitted by District Residents concerning | 203 |

PLATES.

| | Page |
|---|------|
| I. to VII. Larval and pupal stages of West African Culicidæ to face | 50 |
| VIII. West African Hemiptera injurious to Cocoa | 62 |
| IX. <i>Tabanus tæniola</i> , P. de B., and <i>T. par</i> , Walk. | 104 |
| X. Plan of part of Cairo infested by a Mealy Bug (<i>Dactylopius perniciosus</i> , Newst. & Willcocks) | 136 |
| XI. to XIII. West African Fruit-flies (Trypetidæ) | 170 |

M A P S.

| | Page |
|---|-------------|
| Sketch-map showing the distribution of <i>Glossina morsitans</i> in the Lundazi District, N.E. Rhodesia | to face 183 |
| Map showing Tsetse-fly areas in Nyasaland | 204 |
| Map of West Africa, showing places in which the <i>Stegomyia</i> mosquito has been found | 262 |
| Map of West Africa, showing places in which yellow fever has occurred | 262 |
| Sketch-map of a portion of North Nyasa district, to show the distribution of <i>Glossina fusca</i> | 302 |
| Map of N.E. Rhodesia, to illustrate Mr. S. A. Neave's report on a journey to the Luangwa Valley | 316 |

ILLUSTRATIONS IN THE TEXT.

| | Page |
|--|--------------------|
| Siphon and anal segment of <i>Stegomyia fasciata</i> , F. | 26 |
| <i>Stictococcus dimorphus</i> , Newstead, female and male larvæ | 64 |
| <i>Carpophthoromyia pulchella</i> , Austen, ♀ | 72 |
| <i>Carpophthoromyia formosula</i> , Austen, ♂ | 75 |
| <i>Cordylobia prægrandis</i> , Austen, ♀ | 79 |
| Cocoon-mass of <i>Anopheles venuta</i> , Walk., on a leaf of Cassava | 83 |
| Details of fleas found on domestic rats and mice | 91, 92, 93, 96, 98 |
| Figures of Acari :— | |
| <i>Opiliocarus segmentatus</i> , With, ♀ | 105 |
| <i>Holothyreus longipes</i> , Thor., ♂ | 106 |
| <i>Rhipicephalus sanguineus</i> , Latr., ♀ | 107 |
| <i>Ornithodoros moubata</i> , Murray, ♀ | 107 |
| <i>Leiulus truncatus</i> , Oudms., ♀ | 109 |
| <i>Dermonyssus gallinæ</i> , De Geer, ♀ | 109 |
| <i>Uropoda africana</i> , Oudms., deutonymph | 110 |
| <i>Pediculoides spinosus</i> , Kramer, ♀ | 111 |
| <i>Labidostoma denticulatum</i> , Schrank, ♀ | 112 |
| <i>Tarsotomus parietinus</i> , Herm., ♀ | 113 |
| <i>Tetronychus carpini</i> , Oudms. | 114 |
| <i>Cheletes eruditus</i> , Schrank, ♀ | 114 |
| <i>Myobia affinis</i> , Poppe, ♀ | 114 |
| <i>Bdella rubra</i> , Lam. | 114 |
| <i>Euthrombidium trigonum</i> , Herm., larva | 115 |
| <i>Hydryphantès rubra</i> , De Geer, larva | 115 |
| <i>Eremaeus hessei</i> , Oudms. | 116 |
| <i>Aleurobius africana</i> , Oudms., ♀ | 117 |
| <i>Sarcoptes equi</i> , Gerl., ♀ | 117 |
| <i>Chorioptes spathiferus</i> , Mégnin, ♀ | 118 |
| <i>Demodex canis</i> , Leydig, ♀ | 118 |
| <i>Eriophyes tenuis</i> , Nalepa | 119 |
| Healthy lebbek trees (<i>Albizia lebbek</i>) growing on Ghezireh Island, Cairo | 122 |
| Colonies of the mealy bug situated round the edge of a wound, and in the axils of leaf-petioles | 124 |
| Growing point of lebbek attacked by the mealy bug | 125 |
| Compound leaf of lebbek, showing colonies of the mealy bug on the leaf- petioles and elsewhere | 126 |

| | Page |
|--|-----------------------------------|
| Twigs from lebbek tree badly infested with the mealy bug | 127 |
| Lebbek tree in Chareh Kasr el Nil, attacked by <i>Dactylopius</i> | 128 |
| Colonies of <i>Dactylopius</i> protected by a covering of dead lebbek flowers (2 figures) | 129 |
| Chareh el Manakh as it appeared in August 1909, after the trees had been cut back | 133 |
| <i>Dactylopius perniciosus</i> , Newst. & Willcocks, ♀ | 138 |
| Head of <i>Ceratitis nigra</i> , Graham | 163 |
| <i>Inglisia conchiformis</i> , Newst., adult females | 186 |
| <i>Ceroplastes singularis</i> , Newst.: antennæ of adult female, 188; larvæ, 189; male puparia | 189 |
| <i>Ceroplastes ficus</i> , Newst. | 191 |
| <i>Ceroplastes quadrilineatus</i> , Newst.: adult females (tests), 193; antenna of female | 194 |
| <i>Chionaspis dentilobis</i> , Newst., details | 196 |
| <i>Chionaspis substriata</i> , Newst., detail | 197 |
| ? <i>Diaspis chionaspiformis</i> , Green, detail | 198 |
| <i>Enyallopsis durandi</i> , Luc. | 227 |
| <i>Stegomyia fasciata</i> , F.: a favourite breeding-place for, 235; ♀ imago, 240; larva, 243; characteristics of larva | 244 |
| <i>Tabanus ditæniatus</i> , Macq.: immature larva, 266; mature larva, 267; pupa... .. | 268 |
| <i>Tabanus kingi</i> , Austen: view of Khor Arbat, 269; rock at Khor Arbat, showing sites selected by <i>T. kingi</i> for ovipositing, 270; egg-mass and mature larva, 272; pupal case | 273 |
| <i>Chrysops neavei</i> , Austen, ♀ | 276 |
| <i>Diatomineura virgata</i> , Austen, ♀ | 278 |
| <i>Dorcalcemus auricomus</i> , Austen, ♀ | 287 |
| <i>Rhinomyza umbraticola</i> , Austen, ♀ | 290 |
| <i>Tabanus kingi</i> , Austen, ♀ | 291 |
| Heads of <i>Tabanus kingi</i> , Austen, ♀, and <i>T. tenuola</i> , P. de B., ♀ | 292 |
| Antennæ of— <i>Glossina caliginea</i> , Austen; <i>G. palpalis</i> , Rob.-Desv.; <i>G. pallicera</i> , Bigot | 295 |
| Views of the course of the Mawerela Stream, North Nyasa, a haunt of <i>Glossina fusca</i> , Walk. | 300, 301 |
| Views of the valley of the Luangwa River, Rhodesia, visited by Mr. S. A. Neave | 304, 305, 307, 308, 309, 310, 311 |