

Frequency of meals that includes staple, main and side dishes and nutrient intake: findings from the 2012 National Health and Nutrition Survey, Japan

Kazuko Ishikawa-Takata^{1,*} , Kayo Kurotani² , Miyuki Adachi³ and Hidemi Takimoto²

¹Department of Nutrition and Metabolism, National Institute of Health and Nutrition, National Institutes of Biomedical Innovation, Health and Nutrition, Tokyo 162-8636, Japan; ²Department of Nutritional Epidemiology and Shokuiku, National Institute of Health and Nutrition, National Institutes of Biomedical Innovation, Health and Nutrition, Tokyo, Japan; ³Institute of Health and Nutrition, Nagoya University of Arts and Sciences, Nagoya, Aichi, Japan

Submitted 14 June 2019; Final revision received 25 March 2020; Accepted 1 April 2020; First published online 13 July 2020

Abstract

Objective: To investigate the relationship between nutrient intake and a straightforward Japanese nutritional goal of consuming a meal that includes staple, main and side dishes (SMS meal) more than twice a day.

Design: This was a cross-sectional, observational study using the data from de-identified records of the 2012 National Health and Nutrition Survey. Dietary assessment was conducted using semi-weighed dietary records that were completed on one weekday.

Setting: A home-based setting in Japan.

Participants: 30 269 subjects (male n 14 284; female n 15 985) aged ≥ 6 years (6–103 years, mean 50·4 (SD 22·3) years).

Results: As the frequency of SMS meals increased, we found a higher intake of most nutrients and each food group. Subjects who consumed SMS meals at least twice a day were more likely to meet the Dietary Reference Intakes (DRI) for Japan compared with subjects who consumed an SMS meal once a day or less. However, the frequency of SMS meals did not necessarily ensure that DRI for Japan for saturated fatty acid, vitamin A, vitamin B₁, calcium, iron and salt were met.

Conclusions: Consuming SMS meals at least twice a day is a useful and straightforward nutrition goal to achieve adequate intake of some nutrients. However, it is necessary to provide the public with more information on foodstuff choice to ensure an adequate intake of all nutrients.

Keywords

Dietary goal

Dish-based recommendation

Nutrient intake

Foodstuffs

International interest in the traditional Japanese diet has grown, especially after the registration of UNESCO Intangible Cultural Heritage in 2013⁽¹⁾. The Japanese diet is also believed to have an impact on longevity among the Japanese population. In Japan, several policies and recommendations relating to nutrition have been established. The main dietary policies or recommendations in Japan include the Dietary Reference Intakes (DRI) for Japan⁽²⁾, the Japanese Food Guide Spinning Top⁽³⁾, the second term of the National Health Promotion Movement in the 21st century (Health Japan 21, 2013–2023)⁽⁴⁾, and A Basic Act for Shokuiku (food and nutrition education/promotion)⁽⁵⁾. Adherence to the Japanese Food Guide Spinning Top has already been demonstrated to reduce mortality and metabolic risk factors^(6,7).

The main strategies of these policies and recommendations differ. The relationships between these policies/recommendations and the concepts of nutrient-based, foodstuff-based and dish-based recommendations are shown in Fig. 1^(8,9). DRI for Japan shows the recommended nutrient intake and is usually used in various situations of dietary education and meal planning by dietitians and other health professionals. Nutrient-based recommendations, such as the DRI for Japan, are difficult to apply in daily life by non-professionals because calculating the nutrient contents of meals is complicated. The Japanese Food Guide Spinning Top (2005) presents the recommended intake from three dish groups (grains, vegetables and fish and meat) and two additional food groups (fruit and dairy) per day to facilitate the selection of adequate foods in

*Corresponding author: Email kazu@nibiohn.go.jp

© The Author(s), 2020. Published by Cambridge University Press on behalf of The Nutrition Society




Fig. 1 A framework of food/nutrition education. The stratified framework of level of food intake, target behaviour, examples of leading dietary standards, and their contents (nutrients, food groups and dishes) were categorised according to the level of food/nutrition education

daily life⁽³⁾. In addition, Health Japan 21 (2013–23) has set the dietary goal (DG) to increase the number of people who consume meals that include staple, main and side dishes (an SMS meal that includes SMS dishes in one meal) more than twice a day, every day, from 68·1% in 2011 to 80% in 2022⁽⁴⁾. Moreover, A Basic Act for Shokukku (food and nutrition education/promotion) set a similar goal to increase the proportion of individuals who consume SMS meals more than twice a day among all Japanese people, especially those aged 20–39 years⁽⁵⁾.

A complete SMS meal includes SMS dishes and is the traditional and cultural concept of a single meal in Japan. Adachi first mentioned this concept as a tool for nutrition education with a dish-based recommendation⁽¹⁰⁾. Adachi suggested that a practical recommendation for a daily diet, rather than a nutrients-based recommendation alone, was needed to achieve a healthy diet. In this concept, SMS dishes are defined based on their size and main ingredients. Following this work, the usefulness of this dish-based recommendation has been carefully studied^(10,11).

However, more than 30 years have passed since the concept of SMS meal as a dish-based recommendation

was first established. During this time, dietary intakes, lifestyle and health status in Japan have changed dramatically⁽¹²⁾. Thus, the concept of SMS meal in relation to nutrient intake needs to be explored again, especially given that this dish-based concept is still used in many recent nutritional policies.

In this study, we aimed to clarify the relationship between (i) consuming an SMS meal and the intake of each nutrient, and (ii) consuming an SMS meal and agreement of DRI for Japan, using data from the Japan National Health and Nutrition Survey (JNHNS)⁽¹³⁾. Using JNHNS data enabled us to examine these relationships across broad age groups representing the Japanese population.

Methods

Study design

De-identified records from the 2012 JNHNS were cross-sectionally analysed. Secondary use of the data was permitted by the Japanese Ministry of Health, Labour and Welfare.

Study population

The original JNHNS survey included 32 228 subjects aged 1–103 years. In that survey, targeted households were selected by a stratified single-stage cluster design across all forty-seven prefectures in Japan. The household response rate was 67%. The individual response rate was not officially reported, but has been estimated to be around 60%⁽¹⁴⁾. The following subjects were excluded from the current analysis: 1370 subjects aged ≤ 5 years, 368 subjects who were pregnant or lactating, and 221 subjects with an energy intake ± 3 SD for each sex and age group. Children aged ≤ 5 years were excluded because the Japanese Food Guide Spinning Top used to define SMS dishes can only be applied to subjects aged ≥ 6 years⁽³⁾.

Dietary assessment

Dietary data were collected using semi-weighed dietary records that were completed on one weekday between October and November 2012, as described elsewhere^(15–17). Briefly, dietary intake was recorded by a representative household member who was responsible for preparing meals and beverages consumed by all household members. This representative person was asked to weigh each food before and/or after preparation, and to record the weight of food that was uneaten. When the household members shared foods or dishes, the representative household member was asked to record the approximate proportions of foods consumed by each household member. If the weighing of food was difficult, she/he was asked to record the portion size. When the household members ate out, he/she was asked to record their food intake and portion size. In addition, the staff dietitians collected information about the menus of child participants' school lunches. The staff dietitian confirmed what food had been eaten at outside the home with each household member. However, if this was not possible, the staff dietitian asked the representative person to confirm the dietary record when it was not completely recorded. Then, the staff dietitians converted these portion sizes into an estimated weight, and explained these procedures to the representative person. The participants distinguished meals into breakfast, lunch and dinner, and reported these on separate sheets. All drinks and foods consumed that were not meals were categorised as snacks. Macro- and micronutrients were calculated using the Shokujii Shirabe⁽¹⁸⁾, which is an exclusive software developed for the JNHNS according to the 2010 Standard Tables of Food Composition⁽¹⁹⁾. Foods were grouped according to the food classifications used in the JNHNS (cereals, potatoes, sugar, beans, seeds, vegetables, fruits, mushrooms, seaweed, fish, meat, eggs, dairy products, oils, sweets, drinks and seasonings)⁽¹⁵⁾.

Other assessments

Body height and body mass of approximately 90% of the participants were measured by trained fieldworkers using standardised procedures. For the remaining participants,

height and mass were measured either by other household members at home or were self-reported. BMI was calculated as body mass divided by height-squared.

In addition, the first page of the dietary assessment sheet asked participants about their occupation status and exercise habits. If participants exercised > 2 d/week for > 30 min at a time, and had done so for > 1 year, they were classified as having an exercise habit, and answered questions about the frequency, duration and intensity of their exercise.

Calculation of staple, main and side meal frequency

To assess the consumption of an SMS meal from the JNHNS data, we used the definitions of SMS dishes of the Japanese Food Guide Spinning Top⁽³⁾, because it was difficult to apply the basic concept of each dish developed by Adachi⁽¹⁰⁾ to dietary data from the JNHNS. One serving of each dish was defined as follows: staple dish: 40 g of carbohydrate from grain (including rice, bread and noodles); main dish: 6 g of protein from meat, fish, eggs or beans; side dish: 70 g of vegetables (including mushrooms, potatoes and seaweed)⁽³⁾. As with previous studies^(20,21), if a subject consumed at least half a serving of any of these dishes, they were considered to have consumed that dish. A complete SMS meal was defined as a meal that included SMS dishes in a single meal. The number of servings was calculated using two methods. In the first method, the number of servings of each dish on the same plate was calculated, then the number of servings of each dish at one meal was summed. This is the usual method to calculate the number of servings when using the Japanese Food Guide Spinning Top in nutritional education. In the second method, all foods in a single meal were added together, and the number of servings of each dish was calculated. We found that the second method was easy to apply when calculating the number of servings of each dish from the dietary data, and used this method in the study. There were a total of 30 269 meals recorded by all participants; 91·4% of these meals were similarly distinguished as an SMS meal or not, according to these two methods.

The frequency of consuming SMS meals was classified into ≤ 1 , 2, ≥ 3 times per day. Meals were distinguished into breakfast, lunch and dinner. In addition, drinks and foods consumed at times outside of these meals were categorised as snacks. A small proportion of participants (0·8%) ate an SMS meal as a snack; as a result, 0·1% of participants consumed SMS meals four times a day.

Statistical analyses

Data of subjects' characteristics are expressed as mean and SD for numerical data, and as numbers and percentages for categorical data. Other numerical data, including energy, nutrient intake and intake for each food group, are expressed as adjusted mean and 95% CI. Data were analysed using a linear mixed model, including nine regions

in Japan (Hokkaido, Tohoku, Kanto, Hokuriku, Tokai, Kinki, Chugoku, Shikoku and Kyushu) and age as random effect variables. We divided participants into five age groups, as follows: 6–11 (elementary school), 12–17 (junior and senior high school), 18–40 (early adults), 41–64 (late adults) and ≥ 65 (elderly) years.

To assess the relationship between the frequency of SMS meals and healthy nutrient intake, nutrient intakes were compared with the upper and lower limits of a tentative DG for preventing lifestyle-related diseases. In DRI for Japan, DG was set as a percentage of energy intake (%E) from protein, fat, carbohydrate, SFA and grams per day of dietary fibre and salt. If the nutrient did not have a DG, nutrient intake was compared with the estimated average requirement (EAR) and the tolerable upper intake level (UL) of the DRI for Japan 2015⁽²⁾. The DRI for Japan 2015 presents DG or EAR for a total of thirty-four nutrients. Iodine, Se and Mo were excluded because of insufficient information on these in food composition tables. We applied 0·45 and 0·50 mg and 4·8 mg niacin equivalent per 4184 kJ energy intake for the EAR of vitamin B₁, vitamin B₂ and niacin according to EAR. The tolerable UL for vitamin A, folate, Ca, Fe, Zn and Cu was used to assess excess intake. Age groups in the DRI for Japan 2015 were 6–7, 8–9, 10–11, 12–14, 15–17, 18–29, 30–49, 50–69, and >70 years. DG, EAR and UL for the age groups used in the present study were calculated by the weighted mean of each sex and age group. In this study, EAR of Fe for menstruating women was used as EAR of Fe for all women aged 10–69 years.

Energy adjustment was conducted for vitamin A, vitamin B₆, vitamin B₁₂, folate, vitamin C, minerals, dietary fibre, salt and intake from each food group to avoid the effect of underestimation caused by the dietary survey⁽²²⁾. Energy adjustment was calculated as follows: energy-adjusted nutrient or food intake was calculated as amount per day = (reported intake of nutrient or food) \times (estimated energy requirement)/(reported energy intake)⁽²³⁾. The estimated energy requirement for each subject was calculated as the standard RMR shown in DRI for Japan 2015⁽²⁾ multiplied by the subject's body mass and physical activity level. Physical activity level was determined according to the subjects' occupations and exercise habits based on the DRI for Japan⁽²⁾.

Linear trends in energy-adjusted intakes, %E intake, and per 4184 kJ intake of nutrient- and energy-adjusted food intakes across the frequency of SMS meals were compared using a generalised linear model after adjusting for age and the nine regions in Japan, because these could be potential confounding factors on the relationship between the frequency of SMS meals and nutrient or food intake.

To calculate the proportion of subjects who consumed less than the DG (lower limit) or EAR and more than the DG or UL, the distribution of usual intake in these subjects was estimated using the best power method^(24,25) because the JNHNS only included a 1-d dietary survey. To use the best

power method, power transformations and ratios of within-to between-individual variance were quoted from a previous study that reported dietary intake among the normal Japanese population⁽²⁵⁾. Distribution estimation was conducted using an exclusive software (HabitDist)⁽²⁶⁾. The proportion of subjects who consumed less than the DG (lower limit) or EAR and more than the DG or UL by the frequency of SMS meals was compared using the χ^2 test. All other analyses were conducted using IBM SPSS (version 22·0) for Windows (IBM). Two-sided $P < 0·05$ was considered statistically significant.

Results

The proportion of subjects who ate an SMS meal ≥ 3 times per day was highest among subjects aged ≥ 65 years, for both male and female participants (Table 1), while the proportion of subjects who ate ≤ 1 SMS meal per day was highest among male and female participants aged 18–40 years. Approximately 90% of subjects of both sex and all age groups except for the 18–40 and 41–64 years groups ate home-cooked breakfast and dinner, including meals eaten at home and taken in a lunch box. Over 90% of boys and girls aged 6–11 years ate school lunches. Almost equivalent numbers of 12- to 17-year-old boys and girls ate school lunches and home-cooked meals, either eaten at home or taken to school in a lunch box.

A trend analysis examined the relationship between the frequency of SMS meals and nutrient intake (see online supplementary material, Supplementary Table 1a and 1b). Energy intake was significantly higher with a higher frequency of SMS meals, and this was true for all sex and age groups ($P < 0·01$). Among the male participants, protein, dietary fibre, vitamin A, vitamin B₆, folate, vitamin C, Mg, Fe, Zn, Cu and salt intake were higher with a higher frequency of SMS meals in all age groups. However, vitamin B₁, vitamin B₂ and vitamin B₁₂ intake was not related to the frequency of SMS meals in all age groups. Among the female participants, the intakes of protein, dietary fibre, folate, vitamin C, Ca, Mg, Fe, Zn and Cu were higher with a higher frequency of SMS meals in all age groups. However, vitamin B₂ intake was not related to the frequency of SMS meals in all age groups.

In male participants aged 6–11 years, the intakes of potatoes, beans, vegetables, fish and meat were higher with a higher frequency of SMS meals (see online supplementary material, Supplementary Table 2a). In addition to intakes from these food groups, male participants aged 12–17 consumed more mushrooms and seaweeds, but not fish and meat, as the frequency of SMS meals increased. In subjects aged 6–11, 12–17, 18–40, and 41–64 years, sweet intake was lower with an increasing frequency of SMS meals. Among men aged 18–40, 41–64 and ≥ 65 years, intake from almost all food groups was higher as the frequency of SMS meals increased, except for cereals, sugar,

Table 1 Age, physical characteristics, frequency of SMS meals and type of each meal by age and sex group

		6–11 years		12–17 years		18–40 years		41–64 years		≥65 years	
Male participants											
Number		916		976		3208		4942		4242	
Age* (years)		8.6	1.7	14.5	1.7	30.8	6.7	53.9	7.3	74	6.5
Body weight* (kg)		28.6	7.9	52.8	11.5	68.2	12.3	68.0	10.4	61.9	9.3
Height* (cm)		130.0	11.3	163.1	9.8	171.2	5.9	168.5	6.0	162.4	6.3
BMI* (kg/m ²)		16.7	2.4	19.7	3.0	23.2	3.9	23.9	3.2	23.5	3.0
Frequency of SMS meals† (%)	Less than one time per day	206	22.5	220	22.5	1300	40.5	1700	34.4	1173	27.7
	Two times per day	543	59.3	534	54.7	1462	45.6	2251	45.5	1785	42.1
	Equal or more than three times per day	167	18.2	222	22.7	446	13.9	991	20.1	1248	30.3
Breakfast‡	Home-cooked§	837	91.4	873	89.4	2141	66.7	3910	79.1	3935	92.8
	Prepared food	44	4.8	29	3.0	157	4.9	218	4.4	121	2.9
	Eating out¶	0	0.0	2	0.2	77	2.4	103	2.1	27	0.6
	School lunch**	0	0.0	0	0.0	14	0.4	10	0.2	0	0.0
	Others†	35	3.8	72	7.4	819	25.5	701	14.2	159	3.7
Lunch‡	Home-cooked§	60	6.6	452	46.3	1529	47.7	2632	53.3	3264	76.5
	Prepared food	11	1.2	21	2.2	258	8.0	372	7.5	373	8.8
	Eating out¶	3	0.3	42	4.3	952	29.7	1342	27.2	453	10.7
	School lunch**	842	91.9	457	46.8	320	10.0	444	9.0	41	1.0
	Others†	0	0.0	4	0.4	149	4.6	152	3.1	129	3.0
Dinner‡	Home-cooked§	888	96.9	922	94.5	2690	83.9	4363	88.3	3882	91.5
	Prepared food	15	1.6	25	2.6	195	6.1	282	5.7	249	5.9
	Eating out¶	12	1.3	23	2.4	245	7.6	239	4.8	98	2.3
	School lunch**	0	0.0	2	0.2	44	1.4	23	0.5	1	0.0
	Others†	1	0.1	4	0.4	34	1.1	35	0.7	12	0.3
Female participants											
Number		939		843		3199		5721		5283	
Age* (years)		8.6	1.7	14.3	1.7	31.0	6.7	53.7	7.2	75.1	7.4
Body weight* (kg)		28.3	8.0	47.8	8.4	53.0	9.1	54.6	9.1	51	10.7
Height* (cm)		130.4	11.7	155.8	6.2	157.9	5.4	155.6	5.6	148.9	6.3
BMI* (kg/m ²)		16.3	2.3	19.7	2.8	22.2	3.9	23.1	3.5	23	3.5
Frequency of SMS meals† (%)	Less than one time per day	232	24.7	214	25.4	1574	49.2	2331	40.7	1705	32.3
	Two times per day	560	59.6	449	53.3	1257	39.3	2441	42.7	2263	42.8
	Equal or more than three times per day	147	15.7	180	21.4	368	11.5	949	16.6	1315	24.9
Breakfast‡	Home-cooked§	864	92.0	749	88.8	2428	75.9	4959	86.7	4980	94.3
	Prepared food	38	4.0	29	3.4	173	5.4	194	3.4	113	2.1
	Eating out¶	0	0.0	3	0.4	20	0.6	20	0.3	12	0.2
	School lunch**	0	0.0	0	0.0	4	0.1	2	0.0	5	0.1
	Others***	37	3.9	62	7.4	574	17.9	546	9.5	173	3.3
Lunch‡	Home-cooked§	62	6.6	362	42.9	1887	59.0	3873	67.7	4271	80.8
	Prepared food	5	0.5	19	2.3	328	10.3	545	4.5	426	8.1
	Eating out¶	6	0.6	27	3.2	541	16.9	733	12.8	394	7.5
	School lunch**	865	92.1	430	51.0	271	8.5	393	6.9	62	1.2
	Others***	1	0.1	5	0.6	172	5.4	177	3.1	130	2.5
Dinner‡	Home-cooked§	903	92.2	799	94.8	2861	89.4	5334	93.2	4943	93.6
	Prepared food	19	2.0	21	2.5	137	4.3	193	3.4	247	4.7
	Eating out¶	16	1.7	18	2.1	151	4.7	131	2.3	65	1.2
	School lunch**	0	0.0	0	0.0	10	0.3	14	0.2	5	0.1
	Others***	1	0.1	5	0.6	40	1.3	49	0.9	23	0.4

*Mean values and SD.

†An SMS meal is a meal that includes staple, main, and side dishes together in one meal.

‡Numbers and percentages.

§Home-cooked: Food cooked at home and either eaten at home or taken out in a lunch box.

||Prepared food: Take-away or home-delivered food eaten at home.

¶Eating out: Food eaten outside the home either at a restaurant or as a take-away.

**School lunch: A school-prepared lunch or food bought at a company cafeteria.

***Others: Skipped meal, a supplement, or drink only, or unknown.

seeds, dairy products and oils in those aged 18–40 years; sugar, seeds, fruits, dairy products and oils in those aged 41–64 years; and cereals, sugar, dairy products and oils in those aged ≥65 years. In all female participants, sweet intake was lower as the frequency of SMS meals increased

(see online supplementary material, Supplementary Table 2b). Among 6- to 11-year-old girls, the intake of cereals, sugar, fruits, meat, eggs and oils were not related to the frequency of SMS meals. However, in 18- to 64-year-old women only, the intake of sugar, seeds and oils was not

related to the frequency of SMS meals, while in women aged ≥ 65 years, sugar, fruits, seaweeds and seasonings intakes were not related to the frequency of SMS meals consumed.

The proportion of subjects with a nutrient intake less than the DG (lower) or EAR was compared according to the frequency of SMS meals consumed (Table 2a). For almost all nutrients, the proportion of subjects who had an intake that was less than the DG (lower) or EAR was lower with an increase in the frequency of SMS meals. Almost none of the participants consumed protein, carbohydrate, vitamin B₁₂ or Cu less than the lower limit of DG or EAR. However, $\geq 50\%$ of participants who consumed SMS meals more than twice a day still had an inadequate intake of some nutrients, and this differed between sexes and age groups. This included an inadequate intake of vitamin A in men aged 18–40 and 41–64 years; vitamin B₁ in men aged 12–17, 18–40, 41–64 and ≥ 65 years and women aged 12–17 and 18–40 years; Ca in men aged 12–17, 18–40, 41–64 and ≥ 65 years and women aged 18–40 and 41–64 years; and Fe in female participants <65 years.

Table 2b shows the proportion of subjects with a nutrient intake that was more than the DG or UL. Only a small number of subjects consumed excessive nutrients than the reference values, and this included all nutrients except for SFA and salt. In addition, only the proportion of subjects with an excessive salt intake increased by an increase in the frequency of SMS meals.

Discussion

In this study, we found that as the frequency of SMS meals increased, the intake of most nutrients and food groups also increased. In addition, with an increase in the frequency of SMS meals, there was an increase in subjects who met the DG or EAR for most nutrients. However, even in subjects who consumed more than two SMS meals a day, it seemed difficult to meet the DG or EAR for SFA, vitamin A, vitamin B₁, Ca, Fe and salt. In addition, excessive salt intake increased as the frequency of SMS meals increased.

In this study, the proportion of both male and female subjects who consumed ≥ 2 SMS meals a day was lowest among subjects aged 18–40 years. It is difficult to compare this finding with previous ones because other studies had primarily assessed the number of days per week in which two or more SMS meals were consumed. In addition, the definitions of SMS dishes were not comparable with other studies. The Consciousness Survey for Shokuiku 2018 was conducted among adult men and women, and the results suggested that the proportion of subjects who ate ≥ 2 SMS meals almost every day increased with age⁽²⁷⁾. We also found that the proportion of subjects who ate ≥ 2 SMS meals a day was higher among subjects aged ≥ 65 years than in subjects aged 18–40 and 41–64 years. Similarly, Kakutani *et al.*⁽²⁸⁾ reported that only about half of men and women

aged 18–24 years ate more than two SMS meals a day for >3 d a week. While there is a lack of published data in children, we assume that the high proportion of children who consumed SMS meals is partly a result of the school lunch system in Japan; our results suggest that $>90\%$ of subjects aged 6–11 years and nearly half of subjects aged 12–17 years ate school lunches. A study comparing nutrient intake between school days and non-school days among elementary and junior high school students concluded that school lunches improved nutritional intakes⁽²³⁾.

Intakes of most nutrients were higher with a higher frequency of SMS meals. In addition, a lower proportion of participants consumed less than the lower limit of DG or EAR in most nutrients for subjects who ate SMS meals at least twice a day, compared with subjects who less frequently ate SMS meals. Eating ≥ 2 SMS meals a day may represent a way to increase nutrient intake in the population. However, our results also suggest that it is quite difficult to consume sufficient vitamin A, vitamin B₁, Ca and Fe simply by eating SMS meals. Koyama *et al.*'s⁽²⁹⁾ study suggested that subjects with a higher frequency of SMS meals increased their intake of energy, Na, K, Ca, Fe, vitamin B₁, vitamin B₂ and vitamin C. In addition, subjects who consumed SMS meals <1.5 times per day were less likely to meet the DRI for Ca, vitamin A, vitamin B₁, vitamin B₂ and vitamin C. Okubo *et al.*⁽³⁰⁾ examined optimal food intake patterns using a linear programming model. The authors suggested that Fe, vitamin B₁, retinol activity equivalent and Mg made it difficult to analyse the optimal food intake pattern. In addition, the ideal food intake pattern predicted by the model was not that different to the actual intake, except for the intake of fruits and vegetables for younger age groups, and whole grains for all age groups. One possible reason for the difficulty in achieving an adequate intake of some nutrients is that only some foods are rich in vitamin A, vitamin B₁, Ca and Fe. In addition, an SMS meal as defined here does not include an intake of milk and dairy products. Thus, it would be difficult to consume sufficient milk and dairy products by only consuming SMS meals. We therefore propose that more information relating to foodstuff choice in main and side dishes is needed to meet the DRI for these nutrients when dish-based recommendations are used. In addition, the use of refined or whole grains in a staple dish is also recommended.

We also found that high proportions of participants consumed excessive salt, fat and SFA compared with those recommended in the DG. The proportion of participants who consumed excessive fat and SFA decreased with the frequency of SMS meals, but was still high among participants aged 18–40 years who consumed SMS meals ≥ 3 times a day. On the other hand, the proportion of excessive salt intake increased with the frequency of SMS meals. This excessive salt intake is partly caused by an increased use of seasoning with an increase in the frequency of SMS meals. Takimoto *et al.*⁽³¹⁾ showed that the main salt source in Japanese food is salt-containing seasoning and Chinese

Table 2a Proportion of subjects who consumed less than the lower reference values by the frequency of meals including staple, main and side dishes by age group*

	Ref. †	Unit	6–11 years					12–17 years					18–40 years					41–64 years					≥65 years				
			Ref. †	≤1	2	≥3	P	Ref. †	≤1	2	≥3	P	Ref. †	≤1	2	≥3	P	Ref. †	≤1	2	≥3	P	Ref. †	≤1	2	≥3	P
Male participants (n)																											
Protein	DG	%E	13	25.8	9.8	6.0	<0.001	13	29.1	19.9	16.3	0.002	13	36.8	26.7	15.3	<0.001	13	28.8	24.9	14.3	<0.001	13	27.0	14.7	7.4	<0.001
Fat	DG	%E	20	2.0	0.0	0.0	<0.001	20	4.6	1.2	0.0	<0.001	20	7.4	4.0	2.7	<0.001	20	15.7	11.3	11.5	<0.001	20	39.8	27.4	27.6	<0.001
Carbohydrate	DG	%E	50	10.7	10.9	4.2	0.031	50	11.9	9.4	7.7	<0.001	50	21.7	16.7	9.5	<0.001	50	32.9	19.3	13.1	<0.001	50	14.5	9.2	4.6	<0.001
Dietary fiber	DG	g	12	63.6	37.8	15.6	<0.001	18	87.8	76.1	46.4	<0.001	20	82.9	75.8	53.0	<0.001	20	75.9	70.1	46.3	<0.001	20	81.0	66.3	48.2	<0.001
Vitamin A	EAR	µg RAE	369	16.0	1.8	0.0	<0.001	599	59.1	34.1	18.0	<0.001	630	73.7	53.3	29.4	<0.001	616	72.9	54.9	28.0	<0.001	565	65.0	43.2	23.5	<0.001
Vitamin B ₁	EAR	mg/4184 kJ	0.45	58.7	42.4	35.9	<0.001	0.45	60.9	59.9	54.1	0.253	0.45	54.9	57.4	50.9	0.076	0.45	61.3	63.3	56.0	<0.001	0.45	64.9	55.8	54.0	<0.001
Vitamin B ₂	EAR	mg/4184 kJ	0.50	5.8	1.1	3.6	0.001	0.50	28.6	16.1	17.1	<0.001	0.50	36.5	43.2	25.8	<0.001	0.50	24.9	30.2	18.9	<0.001	0.50	19.1	12.0	7.9	<0.001
Niacin	EAR	mg NE/ 4184 kJ	4.8	21.4	7.4	3.6	<0.001	4.8	10.9	8.4	4.5	0.044	4.8	3.6	1.4	0.4	<0.001	4.8	1.5	0.4	0.2	<0.001	4.8	2.6	0.3	0.2	<0.001
Vitamin B ₆	EAR	mg	0.8	36.4	7.7	1.2	<0.001	1.2	42.7	22.1	4.5	<0.001	1.2	18.4	5.2	0.9	<0.001	1.2	14.2	4.5	1.6	<0.001	1.2	29.2	10.7	5.4	<0.001
Vitamin B ₁₂	EAR	µg	1.2	0.0	0.0	0.0	—	2.0	0.0	0.0	0.0	—	2.0	0.4	0.0	0.0	0.0	2.0	0.4	0.0	0.0	<0.001	2.0	1.1	0.0	0.1	<0.001
Folate	EAR	µg	124	2.4	0.2	0.0	0.002	200	15.5	4.9	1.4	<0.001	200	5.4	2.3	0.0	<0.001	200	3.7	0.8	0.0	<0.001	200	4.5	1.3	0.0	<0.001
Vitamin C	EAR	mg	52	53.4	26.2	10.8	<0.001	82	66.8	48.7	23.4	<0.001	85	61.4	46.2	19.1	<0.001	85	44.4	30.5	11.3	<0.001	85	29.4	15.5	6.3	<0.001
Ca	EAR	mg	551	31.6	18.2	13.8	<0.001	752	70.9	63.3	55.4	0.003	590	58.8	59.8	47.3	<0.001	584	53.7	53.9	39.9	<0.001	600	64.9	53.8	44.9	<0.001
Mg	EAR	mg	144	6.8	0.4	0.0	<0.001	274	60.9	49.6	27.9	<0.001	298	54.7	46.2	25.8	<0.001	296	37.0	34.7	17.7	<0.001	276	48.4	33.9	20.7	<0.001
Fe	EAR	mg	6	69.4	41.6	20.4	<0.001	8	46.4	30.3	14.9	<0.001	6	0.9	0.3	0.0	0.022	6	1.1	0.2	0.1	<0.001	6	3.9	1.0	0.0	<0.001
Zn	EAR	mg	5	1.5	0.0	0.0	0.006	8	1.8	0.2	0.0	0.008	8	2.6	0.3	0.0	<0.001	8	4.8	1.2	0.1	<0.001	8	29.0	13.9	5.9	<0.001
Cu	EAR	mg	0.4	0.0	0.0	0.0	—	0.7	0.0	0.0	0.0	—	0.7	0.2	0.0	0.0	0.125	0.7	0.2	0.0	0.0	0.048	0.7	0.2	0.0	0.0	0.073
Female participants (n)																											
Protein	DG	%E	13	26.8	9.2	5.5	<0.001	13	25.3	8.5	6.7	<0.001	13	26.5	14.5	5.5	<0.001	13	21.0	9.5	4.5	<0.001	13	20.2	8.5	2.2	<0.001
Fat	DG	%E	20	0	0.2	0.7	0.361	20	1.0	0.0	0.6	0.148	20	2.1	0.8	0.6	0.005	20	6.3	2.6	2.2	<0.001	20	25.9	15.3	14.0	<0.001
Carbohydrate	DG	%E	50	13.4	10.6	6.9	0.130	50	20.1	14.7	7.8	0.003	50	22.3	15.6	12.8	<0.001	50	16.4	10.4	7.2	<0.001	50	4.0	2.3	2.2	0.002
Dietary fiber	DG	g	12	70.3	48.1	15.0	<0.001	16	76.7	56.0	35.0	<0.001	18	85.2	80.3	65.5	<0.001	18	70.5	61.7	44.0	<0.001	17	60.2	47.7	31.2	<0.001
Vitamin A	EAR	µgRAE	351	5.6	0.5	1.4	<0.001	500	6.1	0.5	7.2	<0.001	480	44.6	27.8	10.6	<0.001	500	48.7	28.4	18.1	<0.001	472	38.5	25.2	11.9	<0.001
Vitamin B ₁	EAR	mg/4184 kJ	0.45	65.1	47.7	36.1	<0.001	0.45	55.1	51.2	51.1	0.607	0.45	45.2	50.0	48.1	0.015	0.45	47.6	46.8	40.7	0.001	0.45	50.7	44.5	30.1	<0.001
Vitamin B ₂	EAR	mg/4184 kJ	0.50	16.8	2.5	6.8	<0.001	0.50	24.8	15.4	12.8	0.002	0.50	25.9	28.5	16.3	<0.001	0.50	23.1	17.0	12.4	<0.001	0.50	15.8	11.8	6.5	<0.001
Niacin	EAR	mg NE/ 4184 kJ	4.8	19.4	10.0	2.7	<0.001	4.8	11.7	6.0	3.9	0.005	4.8	2.8	0.6	0.3	<0.001	4.8	0.7	0.4	0.0	0.027	4.8	1.2	0.2	0.1	<0.001
Vitamin B ₆	EAR	mg	0.8	44.0	17.3	4.8	<0.001	1.1	40.7	25.6	16.7	<0.001	1.0	28.7	18.1	4.3	0.027	1.0	19.8	9.2	3.8	<0.001	1.0	24.9	14.5	5.8	<0.001
Vitamin B ₁₂	EAR	µg	1.5	0.0	0.0	0.0	—	2.0	0.0	0.0	0.0	—	2.0	0.9	0.1	0.0	0.003	2.0	0.7	0.2	0.1	0.009	2.0	0.9	0.1	0.0	<0.001
Folate	EAR	µg	124	4.7	0.0	0.0	<0.001	199	15.0	2.9	1.7	<0.001	200	15.6	7.8	0.8	<0.001	200	5.5	1.7	0.6	<0.001	200	4.9	1.9	0.2	<0.001
Vitamin C	EAR	mg	52	44.0	25.5	10.2	<0.001	82	55.6	49.4	39.4	0.006	85	56.6	48.8	26.9	<0.001	85	35.6	25.5	12.9	<0.001	85	22.0	15.0	7.3	<0.001
Ca	EAR	mg	554	47.8	24.3	23.8	<0.001	632	57.9	37.4	28.9	<0.001	550	66.1	65.3	59.0	0.035	550	54.2	50.3	42.6	<0.001	522	52.3	43.1	33.9	<0.001
Mg	EAR	mg	144	12.9	2.1	0.7	<0.001	249	56.5	36.3	21.7	<0.001	236	50.1	44.2	23.4	<0.001	240	27.3	24.3	13.7	<0.001	220	28.0	17.4	9.3	<0.001
Fe	EAR	mg	8	96.6	93.0	78.9	<0.001	9	83.2	76.2	56.1	<0.001	9	80.5	76.4	56.5	<0.001	9	70.8	60.6	42.4	<0.001	7	30.4	18.8	6.9	<0.001
Zn	EAR	mg	5	0.9	0.0	0.0	0.047	7	1.9	0.0	0.0	0.003	6	2.5	0.2	0.0	<0.001	6	2.3	0.3	0.1	<0.001	6	8.9	2.8	0.7	<0.001
Cu	EAR	mg	0.4	0.0	0.0	0.0	—	0.6	0.0	0.0	0.0	—	0.6	0.2	0.0	0.0	0.212	0.6	0.0	0.0	—	0.6	0.1	0.0	0.0	0.123	

DG, dietary goal; EAR, estimated average requirement; %E, proportion of energy from each nutrient; RAE, retinol equivalent; NE, niacin equivalent.

*The proportion of subjects who consumed less than the DG or EAR was estimated using the best power method.

†Reference values (Ref.) were calculated as the weighted mean values of subjects, including each sex and age group. The P-value was estimated using the χ^2 test. ≤1, 2 and ≥3 indicate the frequency of SMS meals per day. An SMS meal was defined as a combination of staple, main and side dishes.

Table 2b Proportion of subjects who consumed more than the upper reference values by the frequency of meals including staple, main and side dishes by age group*

	Ref.	Unit	6–11 years					12–17 years					18–40 years					41–64 years					≥65 years				
			Ref.	≤1	2	≥3	P	Ref.	≤1	2	≥3	P	Ref.	≤1	2	≥3	P	Ref.	≤1	2	≥3	P	Ref.	≤1	2	≥3	P
Male participants (n)																											
Protein	DG	%E	20	0.0	0.2	0.0	0.709	20	0.5	0.2	0.5	0.758	20	0.9	0.1	0.3	0.011	20	1.2	0.3	0.5	0.001	20	0.8	0.4	0.7	0.352
Fat	DG	%E	30	35.5	39.5	19.8	<0.001	30	32.3	31.1	24.8	0.057	30	21.9	21.5	17.1	<0.001	30	11.8	8.8	5.9	<0.001	30	2.7	2.4	1.7	0.220
SFA	DG	%E											7	51.4	51.8	44.2	0.037	7	33.5	30.2	19.8	<0.001	7	17.4	16.3	11.1	<0.001
Carbohydrate	DG	%E	65	3.4	0.4	0	<0.001	65	6.9	1.7	0.5	<0.001	65	4.8	2.5	2.1	0.002	65	3.8	3.0	1.4	0.002	65	17.4	10.4	10.2	<0.001
Vitamin A	UL	µg RAE	1208	0.0	0.4	0.0	0.502	2344	0.0	0.0	0.0	–	2700	0.1	0.3	0.0	0.323	2700	0.2	0.2	0.4	0.458	2700	0.2	0.1	0.1	0.796
Ca	UL	mg											2500	0.0	0.0	0.0	–	2500	0.0	0.0	0.0	–	2500	0.0	0.0	0.0	–
Fe	UL	mg	33	0.0	0.0	0.0	–	50	0.0	0.0	0.0	–	53	0.0	0.0	0.0	–	52	0.0	0.0	0.0	–	50	0.0	0.0	0.0	–
Zn	UL	mg											43	0.0	0.0	0.0	–	45	0.0	0.0	0.0	–	41	0.0	0.0	0.0	–
Cu	UL	mg											10	0.0	0.0	0.0	–	10	0.0	0.0	0.0	–	10	0.0	0.0	0.0	–
Salt	DG	g	6	83.5	97.3	99.5	<0.001	8	96.0	96.7	98.7	0.209	8	99.4	99.8	99.8	0.214	8	99.2	99.5	99.9	0.041	8	94.9	98.3	99.0	<0.001
Female participants (n)																											
Protein	DG	%E	20	0.0	0.4	0.0	0.508	20	0.5	0.0	0.0	0.230	20	1.7	0.6	0.6	0.014	20	1.2	0.7	1.0	0.193	20	1.6	1.0	1.9	0.074
Fat	DG	%E	30	35	41.1	29.9	0.027	30	51.9	42.4	26.7	<0.001	30	38.9	32.7	22.9	<0.001	30	21.5	17.9	12.0	<0.001	30	4.0	3.0	2.1	0.013
SFA	DG	%E											7	72.5	71.6	57.7	<0.001	7	54.8	54.9	36.8	<0.001	7	23.6	20.8	14.0	<0.001
Carbohydrate	DG	%E	65	1.3	0.2	0.0	0.062	65	1.5	0.3	0.6	0.181	65	3.5	1.3	0.9	<0.001	65	6.6	3.1	1.7	<0.001	65	26.6	15.5	10.3	<0.001
Vitamin A	UL	µgRAE	1208	0.5	0.4	0.0	0.746	2326	0.0	0.0	0.0	0.212	2700	0.2	0.0	0.0	0.212	2700	0.2	0.1	0.0	0.211	2700	0.2	0.1	0.0	0.294
Ca	UL	mg											2500	0.0	0.0	0.0	–	2500	0.0	0.0	0.0	–	2500	0.0	0.0	0.0	–
Fe	UL	mg	33	0.0	0.0	0.0	–	45	0.0	0.0	0.0	–	40	0.0	0.0	0.0	–	40	0.0	0.0	0.0	–	40	0.0	0.0	0.0	–
Zn	UL	mg											35	0.0	0.0	0.0	–	35	0.0	0.0	0.0	–	35	0.0	0.0	0.0	–
Cu	UL	mg											10	0.0	0.0	0.0	–	10	0.0	0.0	0.0	–	10	0.0	0.0	0.0	–
Salt	DG	g	6	85.4	91.8	98.0	<0.001	7	95.8	99.6	100	<0.001	7	96.6	98.3	99.0	0.002	7	97.4	99.0	99.7	<0.001	7	94.8	97.0	98.2	<0.001

RAE, retinol equivalent; NE, niacin equivalent; DG, dietary goal; RDA, reference dietary intake; %E, proportion of energy from each nutrient.

*The proportion of subjects who consumed more than the UL (upper limit of DG) was estimated using the best power method. Reference values (Ref.) were calculated as weighted mean values of subjects, including each sex and age group. ≤1, 2 and ≥3 represent the frequency of SMS meals per day. An SMS meal was defined as a combination of staple, main and side dishes.

noodles. Okubo *et al.*⁽³⁰⁾ also suggested that salt and SFA intake were the limiting factors to make an optimal food intake pattern. Their model suggested that the intake of reduced-fat dairy products should be increased, and that of full-fat dairy products and salt-containing seasoning should be reduced from the current food intake pattern to meet the DRI for Japan.

Dish-based recommendations could have a positive impact on dietary intake. However, the relationship between nutrient- and dish-based recommendations depends on the dietary pattern of the time. Thus, the evaluation of these educational tools should be ongoing. A similar simple tool for recommending a combination of food intake patterns is MyPlate, which was developed in 2011 in the USA. MyPlate is a foodstuffs-based concept rather than dish-based one, and includes a recommended daily intake of grain, protein, vegetables and fruits on one plate, in addition to dairy products. While one review evaluating the use of MyPlate over 8 years found a lack of evidence to support its efficacy⁽³²⁾, another study that examined its efficacy in enforcing healthy eating habits suggested that the awareness of MyPlate increased the Healthy Eating Index⁽³³⁾. Although MyPlate includes five food groups, it only recommends checking whether each food group is represented in one meal, without showing its quantity. One intervention study used MyPlate, but they only reported an improvement of the knowledge of DG rather than any actual changes in nutrient intake or health⁽³⁴⁾.

A most serious limitation of the present study is that the dietary survey in JNHNS was only completed on one day, from which we calculated the frequency of SMS meal intake. Thus, we could not assess the habitual frequency of SMS meal consumption and the associated nutrient intake. Nonetheless, our study provides an indication of such an association. In addition, to calculate the proportion of subjects who did not meet the DG, EAR or UL, we used the best power method to estimate the habitual distribution of nutrient intakes. Second, dietary surveys are known to underestimate food intakes⁽²²⁾. In this study, we applied energy adjustment for nutrients intake and intakes of each food group based on the estimated energy requirement. To calculate the estimated energy requirement, physical activity level was categorised according to the subjects' occupations and exercise habits; however, this may still result in some errors in the estimation of subjects who did not meet the DG, EAR or UL. It is also possible that the use of energy adjustment resulted in an over-adjustment, especially for subjects who did not eat an SMS meal often. However, although the subjects with a higher frequency of SMS meals consumed more energy, the energy-adjusted intake of nutrients and food was still higher among those subjects compared with subjects who consumed an SMS meal less often. Third, the distribution of Fe requirement is known to be skewed, especially for menstruating women⁽³⁵⁾. However, we could not estimate the population distribution of intake from a one-day dietary survey and skewed

requirement distribution simultaneously. We used the same statistical method to estimate the proportion of subjects who did not meet the DG with other nutrients, which could have caused errors in estimating the subjects who did not meet Fe requirements. Fourth, the definition of an SMS meal did not follow the basic concept of SMS dishes proposed by Adachi⁽¹⁰⁾. In the present study, we used food-stuff- and nutrient-based definitions of the Japanese Food Guide Spinning Top to analyse the dietary survey data from JNHNS⁽³⁾. Fifth, we only used age and region as covariates. Given the limited data in JNHNS, we could not examine other variables that may have affected the relationship between the frequency of SMS meals and nutrient intake. Sixth, this study obtained data from the JNHNS database, and so we could not control the dietary assessment method. In this survey, the staff dietitian met the representative person in each household, and verified information about foods eaten outside the home with each individual household member. However, if this was not possible, the representative person asked their household members about details of foods eaten outside the home and confirmed these with the dietitian. To increase the accuracy of dietary survey outside of the home, the staff dietitian collected the school lunch menus of all child participants.

In conclusion, the consumption of SMS meals at least twice a day is associated with a higher nutrient intake and a higher proportion of participants meeting the DRI across all age groups. Thus, the consumption of SMS meals at least twice a day could be a useful and straightforward nutrition goal in daily life. However, more information on foodstuff choice in main and side dishes is needed to meet the DRI for vitamin A, vitamin B₁, vitamin B₂, Ca and Fe, and to avoid an excessive intake of SFA and salt.

Acknowledgements

Acknowledgements: We thank Marion Barnett, BSc, from Edanz Group (www.edanzediting.com/ac) for editing a draft of this manuscript. **Financial support:** This work was supported by the Health and Labour Sciences Research Grants (Special Research Project H26-tokubetsu-shitei-033) from the Ministry of Health, Labour and Welfare of Japan, and Health Japan 21 (the second term), analysis and assessment project. **Conflict of interest:** There are no conflicts of interest. **Authorship:** K.K.I.-T. analysed the data and drafted and critically revised the manuscript. K.K and H.T. drafted and critically revised the manuscript. M.A. joined the discussion to revise the manuscript. All authors contributed to the interpretation of the results and read and approved the final manuscript. **Ethics of human subject participation:** This study was conducted in accordance with the Ethical Guidelines of Epidemiological Research. This study was exempt from the application of these guidelines because anonymised

data was only used in this study. The protocol of the present study was approved by the Ethics Committee of the National Institutes of Biomedical Innovation, Health and Nutrition (20141225-01, 25 December 2014).

Supplementary material

For supplementary material accompanying this paper visit
<https://doi.org/10.1017/S1368980020001202>

References

- United Nations, Educational, Scientific, and Cultural Organization & UNESCO Intangible Cultural Heritage. Palestine continues efforts to safeguard living heritage during the COVID-19 pandemic. <https://ich.unesco.org/en/home> (accessed August 2018).
- Ministry of Health, Labour and Welfare, Japan (2014) *Dietary Reference Intake for Japanese 2015*. Tokyo: Ministry of Health, Labour and Welfare, Japan.
- Yoshiike N, Hayashi F, Takemi Y *et al.* (2007) A new food guide in Japan: the Japanese food guide spinning top. *Nutr Rev* **65**, 149–154.
- Ministry of Health, Labour and Welfare, Japan (2012) Ministerial Notification No. 430 of the Ministry of Health, Labour and Welfare. <https://www.mhlw.go.jp/file/06-Seisaku-jouhou-10900000-Kenkoukyoku/0000047330.pdf> (accessed August 2018) (in Japanese).
- Ministry of Agriculture, Forestry, and Fisheries (2016) Shokuiku Promotion Policies: FY2016 (White Paper on Shokuiku) Summary; available at <http://www.maff.go.jp/e/data/publish/attach/pdf/index-62.pdf> (accessed August 2018) (in Japanese).
- Kurotani K, Akter S, Kashino I *et al.* (2016) Quality of diet and mortality among Japanese men and women: Japan Public Health Center based prospective study. *BMJ* **352**, i1209.
- Nishimura T, Murakami K, Livingstone MB *et al.* (2015) Adherence to the food-based Japanese dietary guidelines in relation to metabolic risk factors in young Japanese women. *Br J Nutr* **114**, 645–653.
- Adachi M (2017) Dish-based nutrition education/promotion with “combination of Shushoku, Shusai and Fukusai method” and “3,1,2 lunch box magic method”: linking research, theory and practice based upon a historical review of literature in Japan since 1970s. An Annual Report of Institute of Health and Nutrition, Nagoya University of Arts and Science 9, 49–83 (in Japanese).
- Harigai J & Adachi M (2018) *Shushoku, Shusai and Fukusai, Dish Composition Table for Diet Coordinate*. Tokyo: Gunyosha (in Japanese).
- Adachi M (1984) A study on the core-dishes and these combination as a framework of nutrition education on the basis of dish-selecting method. *J Health Human Ecol* **50**, 70–107 (in Japanese).
- Harigai J & Adachi M (2014) Validation of the “3,1,2 meal box magic” method for improving the proper intake of energy and nutrients. An Annual Report of Institute of Health and Nutrition, Nagoya University of Arts and Science 6, 33–55 (in Japanese).
- Saito A, Imai S, Htun NC *et al.* (2018) The trends in total energy, macronutrients and sodium intake among Japanese: findings from the 1995–2016 National Health and Nutrition Survey. *Br J Nutr* **120**, 598.
- Ministry of Health, Labour and Welfare, Japan (2014) The National Health and Nutrition Survey in Japan 2012. <https://www.mhlw.go.jp/bunya/kenkou/eiouy/dl/h24-houkoku.pdf> (accessed June 2020) (in Japanese).
- Nishi N, Nakade M, Sarukura N *et al.* (2012) Response rate of National Health and Nutrition Survey and related factors. *Indexes for Public Welfare* **59**, 10–15.
- Katanoda K & Matsumura Y (2002) National Nutrition Survey in Japan – its methodological transition and current findings. *J Nutr Sci Vitaminol* **48**, 423–432.
- Ministry of Health, Labour and Welfare, Japan The National Health and Nutrition Survey in Japan. <https://www.mhlw.go.jp/toukei/itiran/guiyo/k-eisei.html> (accessed June 2020) (in Japanese).
- Iwaoka F, Yoshiike N, Date C *et al.* (2001) A validation study on a method to estimate nutrient intake by family members through a household-based food-weighing survey. *J Nutr Sci Vitaminol* **47**, 222–227.
- National Institute of Health and Nutrition (2018) Manual for Shokuji Shirabe 2018 (Limited for the Registered Members). <http://www.nibiohn.go.jp/eiken/chosa/kenkoeiyo.html> (accessed June 2020) (in Japanese).
- Ministry of Education, Culture, Sports, Science and Technology, Japan (2010) Standard Tables of Food Composition in Japan Fifth Revised and Enlarged Edition. https://www.mext.go.jp/b_menu/shingi/gijyutu/gijyutu3/houkoku/1298713.htm (accessed June 2020) (in Japanese).
- Koyama T, Yoshita K, Arai Y *et al.* (2014) Relationship between daily intake frequency of meals including a staple food, a main dish and a side dish and nutrient intake in independent elderly individuals. *J Jpn Soc Nutr Food Sci* **67**, 299–305 (in Japanese).
- Koyama T, Yoshita K & Arai Y (2016) Effect of food diversity on dietary intakes in independent elderly people. *J Jpn Diet Assoc* **59**, 28–37 (in Japanese).
- Willett WC, Howe GR & Kushi LH (1997) Adjustment for total energy intake in epidemiologic studies. *Am J Clin Nutr* **65**, 1220S–1228S; discussion 1229S–1231S.
- Asakura K & Sasaki S (2017) School lunches in Japan: their contribution to healthier nutrient intake among elementary-school and junior high-school children. *Public Health Nutr* **20**, 1523–1533.
- Nusser SM, Carriquiry KW, Dodd KW *et al.* (1996) A semi-parametric transformation approach to estimating usual daily intake distributions. *J Am Stat Assoc* **91**, 1440–1449.
- Yokoyama T (2013) Theory and application of statistical methods to estimate the distribution of usual intakes of a nutrient in a population: for the appropriate use of dietary reference intakes. *Jpn J Nutr Diet* **71**, S7–S14 (in Japanese).
- Yokoyama T (2012) Estimation of the Distribution of Habitual Intakes of a Nutrient Ver 1.2. <https://www.niph.go.jp/soshiki/gijutsu/download/habitdist/setsumei.pdf> (accessed June 2020) (in Japanese).
- Ministry of Agriculture, Forestry and Fisheries (2018) The Consciousness Survey for Syokuiku. http://www.maff.go.jp/syokuiku/ishiki/h30/pdf_index.html (accessed June 2020) (in Japanese).
- Kakutani Y, Kamiya S & Omi N (2015) Association between the frequency of meals combining “Shushoku, Shusai, and Fukusai” (staple food, main dish, and side dish) and intake of nutrients and food groups among Japanese young adults aged 18–24 years: a cross-sectional study. *J Nutr Sci Vitaminol* **61**, 55–63.
- Koyama T, Yoshita K, Sakurai M *et al.* (2016) Relationship of consumption of meals including grain, fish and meat, and vegetable dishes to the prevention of nutrient deficiency: the INTERMAP Toyama Study. *J Nutr Sci Vitaminol (Tokyo)* **62**, 101–107.
- Okubo H, Sasaki S, Murakami K *et al.* (2015) Designing optimal food intake patterns to achieve nutritional goals for Japanese adults through the use of linear programming optimization models. *Nutr J* **14**, 57.

31. Takimoto H, Saito A, Htun NC *et al.* (2018) Food items contributing to high dietary salt intake among Japanese adults in the 2012 National Health and Nutrition Survey. *Hypertens Res* **41**, 209–212.
32. Chrisman M & Diaz Rios LK (2019) Evaluating MyPlate after 8 years: a perspective. *J Nutr Educ Behav* **51**, 899–903.
33. Jahns L, Conrad Z, Johnson LK *et al.* (2018) Recognition of federal dietary guidance icons is associated with greater diet quality. *J Acad Nutr Diet* **118**, 2120–2127.
34. Brown ON, O'Connor LE & Savaiano D (2014) Mobile MyPlate: a pilot study using text messaging to provide nutrition education and promote better dietary choices in college students. *J Am Coll Health* **62**, 320–327.
35. Subcommittee on Interpretation and Uses of Dietary Reference Intakes and the Standing Committee on the Scientific Evaluation of Dietary Reference Intake, Food and Nutrition Board (2000) Dietary Reference Intakes: Applications in Dietary Assessment. <https://www.ncbi.nlm.nih.gov/books/NBK222873/> (accessed June 2020).