

THE LAW OF THE ITERATED LOGARITHM ON SUBSEQUENCES-CHARACTERIZATIONS

MICHEL WEBER

Let \mathfrak{s} be any increasing sequence of integers and $M > 1$; we connect to them in a very simply way, an increasing unbounded function $\varphi: \mathfrak{s} \rightarrow \mathbf{R}^+$. Let also X_1, X_2, \dots be a sequence of i.i.d. random vectors with value in euclidian space \mathbf{R}^m . We prove that the cluster set of the sequence $\{(X_1 + \dots + X_n)/\sqrt{n} \varphi(n), n \in \mathfrak{s}\}$ almost surely coincides with the unit ball of \mathbf{R}^m , if, and only if, the covariance matrix of X_1 is the identity matrix of \mathbf{R}^m and EX_1 is the zero vector of \mathbf{R}^m . We define a functional A on the set of increasing sequences of integers as follows:

$$A(\mathfrak{s}) = \limsup_{j \rightarrow \infty} \left\{ \frac{\log \#(i \leq j: \mathfrak{s} \cap [M^i, M^{i+1}[\neq \emptyset)}{\log j} \right\}^{1/2}.$$

We prove that $P\{\limsup_{n \rightarrow \infty} (X_1 + \dots + X_n)/\sqrt{2n \log \log n} > 0\} > 0$, for at least one sequence X_1, X_2, \dots of i.i.d. real r.v.'s with $EX_1 = 0$ and $E(X_1)^2 < \infty$, if, and only if $A(\mathfrak{s}) > 0$; further the definition of $A(\cdot)$ does not depend of the value of M . Different characterizations are also established. Further, the law of the iterated logarithm in the sense of Strassen is considered. We finally show a functional law of the iterated logarithm on subsequences for lipschitzian random functions.

§ 1. Introduction

This work is a natural continuation of our previous results obtained in [16]. Let X_1, X_2, \dots be a sequence of independent, identically distributed (i.i.d), real random variables (r.v.'s). Set

$$\forall n \geq 1, \quad S_n = X_1 + \dots + X_n.$$

We investigate the asymptotic behavior of the sequence $\{S_n\}$ when n runs on arbitrary subsequences of integers. This is a quite natural

Received November 28, 1988.

question since it is well-known that, when $EX_1 = 0$, $E(X_1)^2 = 1$,

$$(1.1) \quad P\left\{\limsup_{k \rightarrow \infty} \frac{S_{2^k}}{\sqrt{2 \cdot 2^k \log \log 2^k}} = 1\right\} = 1.$$

In fact, it turns out that for any $\delta \geq 1$, letting

$$(1.1') \quad \mathfrak{s}_\delta = \{2^{\lfloor n^\delta \rfloor}, n \geq 1\},$$

where we write $\lfloor x \rfloor$ the integer part of x ,

$$(1.2) \quad P\left\{\limsup_{\mathfrak{s}_\delta \ni n \rightarrow \infty} \frac{S_n}{\sqrt{2n \log \log n}} = \delta^{-1/2}\right\} = 1,$$

so that (1.1) appears as a particular case ($\delta = 1$) of a much more general phenomenon. From this, we can also deduce the classical Hartmann-Wintner law of the iterated logarithm [7] under a stronger form.

Let now \mathfrak{s} be any strictly increasing sequence of integers and $M > 1$. In a previous work, we connected to \mathfrak{s} and M , in a very simple way a function $\varphi: \mathfrak{s} \rightarrow \mathbf{R}^+$ such that for every sequence X_1, X_2, \dots of i.i.d. real r.v.'s satisfying $EX_1 = 0$, $E(X_1)^2 = 1$,

$$(1.3) \quad P\left\{C_1 \leq \limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{S_n}{\sqrt{n} \varphi(n)} \leq C_2\right\} = 1,$$

where $0 < C_1 \leq C_2 < \infty$ depends on M only. One of our first results in this work (see Theorem 2.1) is that $C_1 = C_2 = 1$ without any modification nor restriction. Conversely, if X_1, X_2, \dots is a sequence of i.i.d. real r.v.'s such that (1.3) holds ($C_1 = C_2 = 1$), for some triple $(\mathfrak{s}, M, \varphi)$ defined above, then

$$EX_1 = 0 \quad \text{and} \quad E(X_1)^2 = 1.$$

Further, this characterization extends to the case of any sequence of i.i.d. random vectors with values in euclidian spaces. This is the Theorem 2.1. The cluster set of the sequence $\left\{\frac{S_n}{\sqrt{n} \varphi(n)}, n \in \mathfrak{s}\right\}$ is also identified, so that the law of the iterated logarithm on any subsequence, in euclidian spaces is characterized. Besides, we characterize all the subsequences \mathfrak{s} such that

$$(1.4) \quad P\left\{\limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{X_1 + \dots + X_n}{\sqrt{2n \log \log n}} > 0\right\} > 0,$$

for at least one sequence X_1, X_2, \dots of i.i.d. real r.v.'s such that $EX_1 = 0$

and $E(X_i)^2 < \infty$. This is the Theorem 2.3. Moreover, we generalize this result in Theorem 2.8. These results are proved in Section 3. Fix now any strictly increasing sequence of integers $\mathfrak{s} = \{n_k, k \geq 1\}$ and $M > 1$. Let $\eta_N \in C([0, 1])$ be obtained by linearly interpolating the partials sums S_{n_k} at $n_k/n_N, 1 \leq k \leq N, N \geq 1$. In Theorem 2.9, we determine the cluster set of the sequence

$$\left\{ \frac{\eta_N}{\sqrt{n_N} \varphi(n_N)}, N \geq 1 \right\},$$

in two particular cases, and therefore we show that this cluster set changes very much with the sequence \mathfrak{s} . This is proved in Section 4. The main feature of the law of the iterated logarithm on subsequences in euclidian spaces, is contained in the fact that the behavior of the partial sums, when indexed on subsequences, can be as small as we want, (see (2.2')). In Theorem 2.10, we show that this feature is preserved in infinite dimension. This result is proved in Section 5.

§ 2. Notations, main results

Let $\mathfrak{s} = \{n_k, k \geq 1\}$ be any strictly increasing sequence of integers and $M > 1$. We associate to them, (as in [16]),

$$(2.1) \quad \left\{ \begin{array}{l} I_0 = I_0(M) = [0, M[\text{ and for each integer } k \geq 1, I_k = I_k(M) = \\ [M^k, M^{k+1}], \text{ for every } p \geq 0, \delta_p = \delta_p(\mathfrak{s}, M) = \begin{cases} 1 & \text{if } \mathfrak{s} \cap I_p(M) \neq \emptyset, \\ 0 & \text{unless,} \end{cases} \\ k_1 = k_1(\mathfrak{s}, M) = \inf \{n \geq 0: \delta_n(\mathfrak{s}, M) = 1\}, \text{ and for every } p > 1, \\ k_p = k_p(\mathfrak{s}, M) = \inf \{n \geq k_{p-1}: \delta_n(\mathfrak{s}, M) = 1\}, \\ \text{for each } p \geq 1, n_p^* \text{ is some point of } \mathfrak{s} \cap I_{k_p}(M) \text{ and } \mathfrak{s}^* = \{n_p^*, p \geq 1\}, \\ \text{for every } n \in \mathfrak{s}, \varphi(n) = \varphi(\mathfrak{s}, M, n) = \sqrt{2 \log(p+2)} \text{ iff } n \in I_{k_p}(M). \end{array} \right.$$

Afterwards, a triple $(\mathfrak{s}, M, \varphi)$ will be always composed of a strictly increasing sequence of integers, a number M strictly greater than 1 and a map $\varphi: \mathfrak{s} \rightarrow \mathbf{R}^+$, defined in accordance with the notations (2.1). Note that φ is depending on \mathfrak{s} and M . Let $\psi: N \rightarrow N$ be strictly increasing and set

$$(2.2) \quad \mathfrak{s}(\psi) = \{2^{\psi(k)}, k \geq 1\},$$

Then, for every $M \geq 1$,

$$(2.2') \quad \lim_{\mathfrak{s}(\psi) \ni n \rightarrow \infty} \frac{\varphi(\mathfrak{s}(\psi), M, n)}{\sqrt{2 \log \psi^{-1}(\log \log n)}} = 1,$$

so that, when ψ grows very fast, the corresponding function φ grows very slowly. Our first result characterizes the law of the iterated logarithm on subsequences in euclidian spaces.

THEOREM 2.1. *Let $(\mathfrak{s}, M, \varphi)$ be any triple defined in accordance with (2.1), and a sequence $X = \{X_i, i \geq 1\}$ of i.i.d. random vectors with value in m -dimensional euclidian space \mathbf{R}^m . Let B_m be the unit ball of \mathbf{R}^m and set*

$$\forall n \geq 1, \quad S_n(X) = X_1 + \dots + X_n.$$

Then,

$$(2.3) \quad P\left\{C\left(\left\{\frac{S_n(X)}{\sqrt{n} \varphi(n)}, n \in \mathfrak{s}\right\}\right) = B_m\right\} = 1,$$

if, and only if,

$$(2.4) \quad \text{Cov}(X_i) \text{ is the identity matrix of } \mathbf{R}^m \text{ and } EX_1 \text{ the zero vector of } \mathbf{R}^m.$$

In particular, when $m = 1$.

$$(2.5) \quad P\left\{\limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{S_n(X)}{\sqrt{n} \varphi(n)} = 1\right\} = 1,$$

if, and only if,

$$(2.6) \quad EX_1 = 0 \quad \text{and} \quad E(X_1)^2 = 1.$$

Further

$$(2.7) \quad P\left\{C\left(\left\{\frac{S_n(X)}{\sqrt{n} \varphi(n)}, n \in \mathfrak{s}\right\}\right) = [-1, 1]\right\} = 1.$$

This characterization is proved in Section 3. Once it is proved for $m = 1$, the general case is easily deduced from the work of H. Finkelstein [5], for the sufficiency part. The necessity part is obtained by choosing suitable linear forms $f_{i,j} \in (\mathbf{R}^m)$, $i, j = 1, \dots, m$, and applying the result obtained for $m = 1$ to the sequences $\{f_{i,j}(X_n), n \in \mathfrak{s}\}$. The proof of Theorem 2.1, when $m = 1$, depends essentially of the following intermediate result:

PROPOSITION 2.2. *Let $(\mathfrak{s}, M, \varphi)$ be any triple defined in accordance with (2.1) and $\delta \geq 1$. Set,*

$$(2.8) \quad \mathfrak{s}_\delta^* = \{n_{[p^\delta]}^*, p \geq 1\},$$

where $[x]$ is the integer part of x . Then, for any sequence X_1, X_2, \dots of i.i.d. real r.v.'s satisfying $EX_1 = 0$ and $E(X_1)^2 = 1$,

$$(2.9) \quad P \left\{ \limsup_{\mathfrak{s}_j^* \ni n \rightarrow \infty} \frac{S_n(X)}{\sqrt{n} \varphi(n)} = (\delta)^{-1/2} \right\} = 1.$$

This is precisely the intermediate step (3.59) in the proof of lemma 3.4; further (2.9) easily implies (2.7). One sees in particular, when $\delta = 1$, that the lower bound in (2.5) is reached on \mathfrak{s}^* .

Next, we define on the set of strictly increasing sequences of integers, a functional as follows: let $M > 1$, then we put:

$$(2.10) \quad A(\mathfrak{s}) = \limsup_{j \rightarrow \infty} \left\{ \frac{\log \#(i \leq j: \mathfrak{s} \cap [M^i, M^{i+1}[\neq \emptyset)}{\log j} \right\}^{1/2}.$$

It is easily seen that

$$(2.11) \quad A(\mathfrak{s}) = \limsup_{j \rightarrow \infty} \left\{ \frac{\log j}{\log k_j(\mathfrak{s}, M)} \right\}^{1/2},$$

where we use the notations (2.1).

Note that $0 \leq A(\mathfrak{s}) \leq 1$ and $A(\mathfrak{s})$, a priori, depends on the value of $M > 1$. This functional will give us the possibility to characterize in a very simple way the sequences of integers which "support" the law of the iterated logarithm.

THEOREM 2.3. *Let X_1, X_2, \dots be any sequence of i.i.d. real r.v.'s satisfying $EX_1 = 0$ and $E(X_1)^2 = 1$. Then, for any strictly increasing sequence of integers \mathfrak{s} ,*

$$(2.12) \quad P \left\{ \limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{X_1 + \dots + X_n}{\sqrt{2n \log \log n}} > 0 \right\} > 0,$$

if, and only if,

$$(2.13) \quad A(\mathfrak{s}) > 0.$$

Then, we have,

$$(2.14) \quad P \left\{ \limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{X_1 + \dots + X_n}{\sqrt{2n \log \log n}} = A(\mathfrak{s}) \right\} = 1.$$

From (2.14), follows that $A(\mathfrak{s})$ does actually not depend on the value of $M > 1$. This result takes a very simple form when the sequence \mathfrak{s} is of the type defined in (2.2).

COROLLARY 2.4. *Let X_1, X_2, \dots be a sequence of i.i.d. r.v.'s satisfying $EX_1 = 0$ and $E(X_1)^2 = 1$. Then, for any strictly increasing map $\psi: N \rightarrow N$,*

$$(2.15) \quad P \left\{ \limsup_{\mathfrak{s}(\psi) \ni n \rightarrow \infty} \frac{X_1 + \dots + X_n}{\sqrt{2n \log \log n}} = \limsup_{m \rightarrow \infty} \left[\frac{\log m}{\log \psi(m)} \right]^{1/2} \right\} = 1,$$

where the sequence $\mathfrak{s}(\psi)$ is defined in (2.2).

In our next approach, we collect some easy properties of the functional $A(\cdot)$. We set for any integer $r \geq 1$, $\alpha > 0$ and any increasing sequence of integers \mathfrak{s} ,

$$(2.16) \quad \begin{cases} P_\alpha(\mathfrak{s}) = \{[n^\alpha], n \in \mathfrak{s}\} \text{ and } P_\alpha = P_\alpha(N), \\ 2^\mathfrak{s} = \{2^n, n \in \mathfrak{s}\}, \\ \mathfrak{s}^{(r)} = \{n_1 + \dots + n_r, \forall i = 1, \dots, r, n_i \in \mathfrak{s}\}, \\ \alpha\mathfrak{s} = \{[\alpha n], n \in \mathfrak{s}\}. \end{cases}$$

LEMMA 2.5. *The following properties hold for any $\alpha > 0$ and any increasing sequence of integers \mathfrak{s} ,*

- a) $A(\alpha\mathfrak{s}) = A(P_\alpha(\mathfrak{s})) = A(\mathfrak{s})$,
- b) $A(2^\mathfrak{s}) = A(2^{\alpha\mathfrak{s}})$,
- c) *if $\mathfrak{s}_i = \{n_k^i, k \geq 1\}$, $i = 1, 2$ satisfy*

$$(2.16') \quad 0 < \liminf_{k \rightarrow \infty} n_k^1/n_k^2 \leq \limsup_{k \rightarrow \infty} n_k^1/n_k^2 < \infty,$$

then, $A(\mathfrak{s}_1) = A(\mathfrak{s}_2)$.

We notice that b) expresses the fact that $A(\cdot)$ does actually not depend of the number $M > 1$ used in its definition; further a) follows easily from b). As for c), since there are constants $0 < C_1 \leq C_2 < \infty$, such that for all k ,

$$C_1 \leq n_k^1/n_k^2 \leq C_2,$$

we therefore have $\#\{i \leq n: \mathfrak{s}_1 \cap [M^i, M^{i+1}] \neq \emptyset\} \leq \inf\{p: M^n \leq C_1 M^{k_p(\mathfrak{s}_2, M)}\}$, for all n , and this easily leads to $A(\mathfrak{s}_1) \leq A(\mathfrak{s}_2)$, which implies c) by symmetry.

Let now $\xi = \{\xi_n, n \geq 1\}$ be an increasing sequence of integer valued r.v.'s such that

$$(2.17) \quad P \left\{ \lim_{k \rightarrow \infty} \frac{\xi_k}{n_k} = 1 \right\} = 1,$$

for some increasing sequence $\mathfrak{s} = \{n_k, k \geq 1\}$.

From the previous lemma, one deduces

$$(2.18) \quad P\{A(\{\xi_n, n \geq 1\}) = A(\xi)\} = 1.$$

Therefore, putting together Theorem 2.3 and (2.18) leads to

COROLLARY 2.6. *Let $\mathfrak{s} = \{n_k, k \geq 1\}$ be an increasing sequence of integers and an increasing sequence $\xi = \{\xi_k, k \geq 1\}$ of integer valued r.v.'s related to \mathfrak{s} by (2.17). Then, for any sequence X_1, X_2, \dots of i.i.d. real r.v.'s with $EX_1 = 0, E(X_1)^2 = 1$, which is independent of the sequence ξ , one has*

$$(2.19) \quad P\left\{\limsup_{k \rightarrow \infty} \frac{\sum_{i=1}^{\mathfrak{s}_k} X_i}{\sqrt{2n_k \log \log n_k}} = A(\mathfrak{s})\right\} = 1.$$

The next corollary establishes some relation between the law of the iterated logarithm and the classical Waring's problem in additive number theory, (see e.g. [15], Chapter I).

COROLLARY 2.7. *Let r, q be two positive integers. For any sequence X_1, X_2, \dots of i.i.d. real r.v.'s satisfying $EX_1 = 0$ and $E(X_1)^2 = 1$,*

$$(2.20) \quad P\left\{\limsup_{2^{P_q^{(r)}} \ni n \rightarrow \infty} \frac{X_1 + \dots + X_n}{\sqrt{2n \log \log n}} = \limsup_{N \rightarrow \infty} \left[\frac{\log \#\{n = i_1^q + \dots + i_r^q \leq N\}}{\log N} \right]^{1/2}\right\} = 1,$$

where we use the notation (2.16).

This raises the following question: to determine the asymptotic behavior of

$$\{(X_1 + \dots + X_n)/\sqrt{2n \log \log n}, n \in 2^{s(r)}\}$$

with respect to those of

$$\{(X_1 + \dots + X_n)/\sqrt{2n \log \log n}, n \in 2^s\}.$$

Indeed, the fundamental aspect of the Waring's problem being solved (see again [15]), we know that

$$\forall q \text{ integer, } \exists r = r(q) < \infty \text{ such that } P_q^{(r)} = N,$$

in other words each sequence P_q defines a natural basis of the integers. For instance, if $q = 2$

$$P_2^{(4)} = N,$$

so that,

$$(2.20a) \quad P\left\{\limsup_{P_2^{(1)} \ni n \rightarrow \infty} \frac{X_1 + \dots + X_n}{\sqrt{2n \log \log n}} = (2)^{-1/2}\right\} = 1,$$

whereas,

$$(2.20b) \quad P\left\{\limsup_{P_2^{(4)} \ni n \rightarrow \infty} \frac{X_1 + \dots + X_n}{\sqrt{2n \log \log n}} = 1\right\} = 1.$$

We thus can wonder what becomes (2-20) when indexing the partials sums on $P_2^{(2)}$ or $P_2^{(3)}$. We have not been able to answer this question. The problem solved by Theorem 2.3 extends immediately: let $(\mathfrak{s}, M, \varphi)$ be a triple defined in accordance with (2.1); characterize all the subsequences, $\mathfrak{s}_1 \subset \mathfrak{s}$, such that

$$(2.21) \quad P\left\{0 < \limsup_{\mathfrak{s}_1 \ni n \rightarrow \infty} \frac{X_1 + \dots + X_n}{\sqrt{n} \varphi(n)} \leq 1\right\} > 0,$$

for at least one sequence X_1, X_2, \dots of i.i.d. real r.v.'s satisfying $EX_1 = 0, E(X_1)^2 = 1$. Put,

$$(2.22) \quad A(\mathfrak{s}, M, \mathfrak{s}_1) = \limsup_{j \rightarrow \infty} \left[\frac{\log \#\{i \leq j: \mathfrak{s}_1 \cap [M^i, M^{i+1}[\neq \emptyset\}}{\log \#\{i \leq j: \mathfrak{s} \cap [M^i, M^{i+1}[\neq \emptyset\}} \right]^{1/2}.$$

We have the following characterization

THEOREM 2.8. *Let $(\mathfrak{s}, M, \varphi)$ be any triple defined in accordance with the notation (2.1). Then, for any sequence X_1, X_2, \dots of i.i.d. real r.v.'s $EX_1 = 0, E(X_1)^2 = 1$, and any subsequence $\mathfrak{s}_1 \subset \mathfrak{s}$,*

$$(2.23) \quad P\left\{\limsup_{\mathfrak{s}_1 \ni n \rightarrow \infty} \frac{X_1 + \dots + X_n}{\sqrt{n} \varphi(n)} = A(\mathfrak{s}, M, \mathfrak{s}_1)\right\} = 1.$$

This theorem is proved in Section 3. Its proof as well as proof of Theorem 2.3, depends essentially of the Theorem 3.3.

Fix now a triple $(\mathfrak{s}, M, \varphi)$ according to the notation (2.1) with $\mathfrak{s} = \{n_k, k \geq 1\}$, as well as a sequence X_1, X_2, \dots of i.i.d. real r.v.'s satisfying $EX_1 = 0, E(X_1)^2 = 1$. Let $\eta_N \in \mathcal{C}([0, 1]), N \geq 1$, be obtained by linearly interpolating the partial sums

$$S_{n_k} = X_1 + \dots + X_{n_k},$$

at points $n_k/n_N, 1 \leq k \leq N$. Thus, (with the convention $S_0 = 0$)

$$(2.24) \quad \begin{cases} \forall t \in [0, 1], \eta_N(t) = S_{n_k} \left[1 + \frac{n_k - tn_N}{n_{k+1} - n_k} \right] + S_{n_{k+1}} \left[\frac{tn_N - n_k}{n_{k+1} - n_k} \right], \\ \text{if } n_k \leq tn_N \leq n_{k+1}. \end{cases}$$

We put,

$$(2.25) \quad \begin{cases} K_1 = \left\{ f(t) = \int_0^t g(u)du : \int_0^1 g^2(u)du \leq 1 \right\}, \\ K_2 = \{f(t) = \varepsilon t \quad \varepsilon \in [-1, 1]\}. \end{cases}$$

The next result states a law of the iterated logarithm in the sense of V. Strassen [12], for subsequences.

THEOREM 2.9. *Let $(\mathfrak{s}, M, \varphi)$ be any triple defined in accordance with (2.1), and a sequence X_1, X_2, \dots of i.i.d. real r.v.'s satisfying $EX_1 = 0$ and $E(X_1)^2 = 1$.*

a) *if $\lim_{k \rightarrow \infty} n_k/n_{k+1} = 1$, then*

$$(2.26) \quad P\{\lim_{N \rightarrow \infty} \text{dist}(\eta_N/\sqrt{n_N} \varphi(n_N), K_1) = 0\} = 1,$$

and

$$(2.27) \quad P\left\{C\left(\left\{\frac{\eta_N}{\sqrt{n_N} \varphi(n_N)}, N \geq 1\right\}\right) = K_1\right\} = 1.$$

b) *if $\lim_{k \rightarrow \infty} n_k/n_{k+1} = 0$, then (2.26) and (2.27) hold with K_2 instead of K_1 .*

This theorem is proved in Section 4. It is also to be connected with Theorem 4.3. Theorem 2.1 together with (2.2') show that in euclidian spaces, the partial sums of i.i.d. r.v.'s, when indexed on subsequences, can grow as slow as we want, it is enough to choose a subsequence of type defined in (2.2) with a function ψ that grows very fast. The aim of our next statement is to show that the same property can happen in infinite dimensional spaces.

THEOREM 2.10. *Let (T, Θ) be a compact topological space. Assume there exists a sample continuous gaussian process $G = \{G(\omega, t), \omega \in \Omega, t \in T\}$ and set*

$$\forall s, t \in T, \quad \rho(s, t) = [E\{G(s) - G(t)\}^2]^{1/2}.$$

Let $Y = \{Y(\omega, t), \omega \in \Omega, t \in T\}$ be a Θ -separable random function satisfying

$$(2.28) \quad E \sup \left\{ \left| \frac{Y(s) - Y(t)}{\rho(s, t)} \right|^2, (s, t) \in T \otimes T \right\} < \infty,$$

$$(2.29) \quad \forall t \in T, \quad EY(t) = 0 \quad \text{and} \quad EY^2(t) < \infty.$$

Let $(\mathfrak{s}, M, \varphi)$ be any triple defined in accordance with (2.1) and a sequence Y_1, Y_2, \dots of independent copies of Y . Then,

$$(2.30) \quad P \left\{ \left\{ \frac{Y_1 + \dots + Y_n}{\sqrt{n} \varphi(n)}, n \in \mathfrak{s} \right\} \text{ is relatively compact in } \underline{C}(T) \right\} = 1.$$

In other words, the random function Y satisfies the compact law of the iterated logarithm in $\underline{C}(T)$ for any subsequence \mathfrak{s} .

This result is proved in Section 5.

§ 3. Characterization of the law of the iterated logarithm on subsequences

Let us denote $W = \{W(t), 0 \leq t < \infty\}$ the usual brownian motion. In accordance with the notations (2.1), let also $(\mathfrak{s}, M, \varphi)$ be a triple in which \mathfrak{s} is a strictly increasing sequence of integers, $M > 1$ and φ defined by (2.1). Set,

$$(3.1) \quad C(M) = \text{median} (\sup \{W(t)/\sqrt{t}, 1 \leq t \leq M\}).$$

LEMMA 3.1. For any triple $(\mathfrak{s}, M, \varphi)$,

$$(3.2) \quad P\{\limsup_{\mathfrak{s} \ni n \rightarrow \infty} W(n)/\sqrt{n} - \varphi(n) \leq C(M)\} = 1.$$

Furthermore,

$$(3.2') \quad P\{\limsup_{\mathfrak{s} \ni n \rightarrow \infty} W(n)/\sqrt{n} \varphi(n) \leq 1\} = 1.$$

Proof. Fix $\varepsilon > 0$ and set,

$$(3.3) \quad \forall p \geq 1, \quad A_p = \{\sup (W(t)/\sqrt{t}, t \in \mathfrak{s} \cap I_{k_p}) > \sqrt{2 \log(p+2)} + C(M) + \varepsilon\}.$$

By applying Borell's inequality [1], and letting $\psi(x) = P\{N(0, 1) > x\}$,

$$(3.4) \quad \forall p \geq 1, \quad P\{A_p\} \leq \psi(\sqrt{2 \log(p+2)} + \varepsilon),$$

so that, $P\{A_p, p \text{ i.o.}\} = 0$. Q.E.D.

LEMMA 3.2. For any triple $(\mathfrak{s}, M, \varphi)$ and any sequence Y_1, Y_2, \dots of i.i.d. real r.v.'s satisfying $EY_1 = 0$ and $E(Y_1)^2 = 1$,

$$(3.5) \quad P\left\{\limsup_{\delta \ni n \rightarrow \infty} \frac{Y_1 + \dots + Y_n}{\sqrt{n} \varphi(n)} \leq 1\right\} = 1.$$

Proof. By virtue of the Skohorod embedding scheme of partial sums (see e.g. [2], Theorem 13.6, p. 276), it is enough to prove

$$(3.5') \quad P\left\{\limsup_{\delta \ni n \rightarrow \infty} \frac{W(T_1 + \dots + T_n)}{\sqrt{n} \varphi(n)} \leq 1\right\} = 1,$$

where T_1, T_2, \dots is a sequence of nonnegative, independent, identically distributed r.v.'s with $ET_1 = 1$. Given $h > 0$, we can choose q large enough, so that, by virtue of the strong law of large numbers, the event

$$(3.6) \quad A = \left\{\sup\left(\left|\frac{\sum_{i=1}^n T_i}{n} - 1\right|, n \in \delta, n \geq q\right) \leq h\right\},$$

has a probability greater than $1 - h$. Fix $\varepsilon > 0$ and set,

$$(3.7) \quad \begin{cases} D = \text{median}\{(\sup(W(\theta t)/\sqrt{\theta}, 1 \leq \theta \leq M, 1 - h \leq t \leq 1 + h)), \\ \forall p \geq 1, C_p = \{\sup(W(\sum_{i=1}^n T_i)/\sqrt{n}, n \in \delta \cap I_{k_p}) \\ > \sqrt{2(1+h)\log(p+2)} + D + \varepsilon\}, \\ \forall p \geq 1, C'_p = C_p \cap A. \end{cases}$$

On C'_p , one has $\sum_{i=1}^n T_i = \delta n$ for some $\delta \in [1 - h, 1 + h]$, and $n = \theta M^{k_p}$ for some $\theta \in [1, M]$. Thus,

$$(3.8) \quad C'_p \subset C''_p = \left\{\sup\left(\frac{W(\theta \delta M^{k_p})}{\sqrt{\theta M^{k_p}}}, 1 \leq \theta \leq M, 1 - h \leq \delta \leq 1 + h\right) > \sqrt{2(1+h)\log(p+2)} + D + \varepsilon\right\},$$

and

$$P\{C''_p\} = P\{\sup(W(\theta \delta)/\sqrt{\theta}, 1 \leq \theta \leq M, 1 - h \leq \delta \leq 1 + h) > \sqrt{2(1+h)\log(p+2)} + D + \varepsilon\}.$$

Since $\sup\{E(W(\theta \delta)/\sqrt{\theta})^2, 1 \leq \theta \leq M, 1 - h \leq \delta \leq 1 + h\} = 1 + h$, Borell's inequality again implies,

$$(3.9) \quad P\{C''_p\} \leq \psi(\sqrt{2\log(p+2)} + \varepsilon/\sqrt{1+h}).$$

Therefore,

$$(3.10) \quad P\{C''_p, p \text{ i.o.}\} = 0,$$

and thus,

$$P\left\{\limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{W(\sum_{i=1}^n T_i)}{\sqrt{n} \varphi(n)} \leq \sqrt{1+h}\right\} \geq 1-h.$$

Since h is arbitrary, we finally obtain,

$$(3.11) \quad P\left\{\limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{W(\sum_{i=1}^n T_i)}{\sqrt{n} \varphi(n)} \leq 1\right\} = 1. \quad \text{Q.E.D.}$$

Unlike the previous results, the proof of the next theorem involves a far more important work.

THEOREM 3.3. *Let $(\mathfrak{s}, M, \varphi)$ be any triple defined in accordance with (2.1). Let also $X = \{X_i, i \geq 1\}$ be any sequence of i.i.d. real r.v.'s satisfying $EX_1 = 0$ and $E(X_1)^2 = 1$. Fix any $0 < \rho < 1$ and $0 < \eta < 1 - \rho^2$ and let*

$$(3.12) \quad \begin{cases} \mathfrak{s}(\rho, X) = \{n \in \mathfrak{s} : X_1 + \dots + X_n \geq \rho\sqrt{n} \varphi(n)\}, \\ \mathfrak{s}(\rho, \eta, X) = \{n \in \mathfrak{s} : X_1 + \dots + X_n \geq \rho\sqrt{n} \varphi(n) \text{ and} \\ \varphi(n) \geq \Lambda(\mathfrak{s})\sqrt{2\eta \log \log n}\}. \end{cases}$$

Then

$$(3.13) \quad P\{\#\mathfrak{s}(\rho, X) = \infty\} = 1,$$

and

$$(3.13') \quad P\left\{\limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{X_1 + \dots + X_n}{\sqrt{n} \varphi(n)} = 1\right\} = 1.$$

Further

$$(3.14) \quad P\{\Lambda(\mathfrak{s}(\rho, \eta, X)) \geq \Lambda(\mathfrak{s})\sqrt{1-\rho^2}\} = 1,$$

and

$$(3.15) \quad P\left\{\limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{X_1 + \dots + X_n}{\sqrt{2n \log \log n}} = \Lambda(\mathfrak{s})\right\} = 1.$$

Proof.

Step 1. By virtue of the Skohorod embedding scheme for partial sums, it is enough to prove the theorem when we replace $X_1 + \dots + X_n$ by $W(T_1 + \dots + T_n)$, $n \geq 1$, where T_1, T_2, \dots is a sequence of i.i.d. non-negative r.v.'s with $ET_1 = 1$. Let h, μ, ρ and ρ_0 be fixed in $]0, 1[$ with $\rho_0 + \sqrt{h} < \rho$,

$$(3.16) \quad \varepsilon(h) = \text{median} \{ \sup (|W(\theta) - W(1)|, 1 - h \leq \theta \leq 1 + h) \},$$

and for each integer $p \geq 1$,

$$(3.17) \quad \begin{cases} \varepsilon_p = \varepsilon(h) + \mu\sqrt{h} + \sqrt{2h \log(p+2)}, \\ A_p = \{ \exists n \in [M^p, M^{p+1}[\cap \mathfrak{B} : X_1 + \dots + X_n \geq \rho_0\sqrt{n} \varphi(n) \}, \\ A_p^0 = \{ X_1 + \dots + X_{n_p^*} \geq \rho_0\sqrt{2n_p^* \log(p+2)} \}, \\ A'_p = \{ \inf (W(\theta n_p^*)\sqrt{n_p^*}, 1 - h \leq \theta \leq 1 + h) \geq \rho_0\sqrt{2 \log(p+2)} \}, \\ A''_p = \{ W(n_p^*)/\sqrt{n_p^*} > \rho\sqrt{2 \log(p+2)} \}, \\ A'''_p = \{ \sup (|W(\theta n_p^*) - W(n_p^*)|/\sqrt{n_p^*}, 1 - h \leq \theta \leq 1 + h) \leq \varepsilon_p \}, \end{cases}$$

and choose q large enough so that, by virtue of the strong law of large numbers, the event

$$(3.18) \quad A = \left\{ \sup \left(\left| \frac{\sum_{i=1}^n T_i}{n} - 1 \right|, n \geq q \right) \leq h \right\},$$

has a probability greater than $1 - h$.

Observe now, for any m_0 large enough and any $m \geq m_0$,

$$(3.18') \quad \begin{aligned} \sum_{p=m_0}^m I_{A_{k_p}} &\geq \sum_{p=m_0}^m I_{A_p^0}, \\ &\geq \sum_{p=m_0}^m I_{A'_p} \cdot I_A, \\ &\geq \sum_{p=m_0}^m I_{A''_p} \cdot I_{A'''_p} \cdot I_A, \\ (3.19) \quad &\geq [\sum_{p=m_0}^m (I_{A''_p} - I_{(A''_p)^c})] \cdot I_A. \end{aligned}$$

Suppose now,

$$(3.20) \quad \sum_{p \geq 1} P\{(A''_p)^c\} < \infty.$$

Then,

$$(3.21) \quad P\{\exists m_0 < \infty : \forall m \geq m_0, \sum_{p=m_0}^m I_{A_p^0} \geq \sum_{p=m_0}^m I_{A'_p} I_A\} = 1.$$

By applying the classical Paley-Zygmund inequality for nonnegative square integrable r.v.'s: $P\{X \geq \lambda EX\} \geq [1 - \lambda]^2 [EX]^2 / E(X)^2$, $0 \leq \lambda \leq 1$, one has

$$(3.22) \quad \begin{aligned} P\{\sum_{p=m_0}^m I_{A''_p} \geq \lambda \sum_{p=m_0}^m P\{A''_p\}\} \\ \geq [1 - \lambda]^2 \frac{[\sum_{p=m_0}^m P\{A''_p\}]^2}{\sum_{p=m_0}^m P\{A''_p\} + \sum_{\substack{p,q=m_0 \\ p \neq q}}^m P\{A''_p \cap A''_q\}}. \end{aligned}$$

Further, suppose,

$$(3.23) \quad \text{for any } 0 < h < 1, \text{ there exists } m_0 = m_0(h) < \infty, \text{ such that for every } m \geq m_0,$$

$$\sum_{\substack{m_0 \leq p, q \leq m \\ p \neq q}} P\{A''_p \cap A''_q\} \leq (1 + h)[\sum_{p=m_0}^m P\{A''_p\} + (\sum_{p=m_0}^m P\{A''_p\})^2],$$

and

(3.24) for any $\rho_1 \in]\rho, 1[$, there exists $m_0 = m_0(\rho) < \infty$, and $m_1 = m_1(m_0, \rho_1) < \infty$, such that for every $m > \sup(m_0, m_1)$,

$$\sum_{p=m_0}^m P\{A''_p\} \geq m^{1-\rho_1}.$$

Then, putting together (3.19), (3.20), (3.21), (3.22), (3.23) and (3.24) leads to

(3.25) for any strictly increasing sequence of integers \bar{E} and any $0 < \rho_0 < \rho_1 < 1$,

$$P\{\sum_{p=1}^m I_{A_p^0} \geq m^{1-\rho_1}, m \in \bar{E}, m \text{ i.o.}\} = 1.$$

This is our first step. We therefore must show (3.20), (3.23) and (3.24). We first prove (3.20). By using Borell's inequality,

$$\begin{aligned} (3.26) \quad \forall p \geq 1, P\{(A''_p)^c\} & \leq P\{\sup(|W(\theta n_p^*) - W(n_p^*)|/\sqrt{n_p^*}, 1 - h \leq \theta \leq 1) > \varepsilon_p\} \\ & \quad + P\{\sup(|W(\theta n_p^*) - W(n_p^*)|/\sqrt{n_p^*}, 1 \leq \theta \leq 1 + h) > \varepsilon_p\}, \\ & \leq 2\psi([\varepsilon_p - \varepsilon(h)]/\sqrt{h}), \\ & \leq 2\psi(\sqrt{2 \log(p+2)} + \mu). \end{aligned}$$

Therefore, by applying Borel-Cantelli lemma, leads to (3.20). In order to obtain (3.23), we need the following classical estimate on gaussian distributions, (see e.g. [3], p. 269–270).

LEMMA 3.4. *Let U, V be jointly gaussian real r.v.'s satisfying $EU^2 = EV^2 = 1$, $EU = EV = 0$, $EUV = r$ and let $\varepsilon > 0$.*

a) *for any $x > 0$, $y > 0$ such that $rx y \leq \varepsilon$,*

$$P\{U > x, V > y\} \leq c(\varepsilon)P\{U > x\}P\{V > y\},$$

where $\lim_{\varepsilon \rightarrow 0} c(\varepsilon) = 1$;

b) *for any $a \geq 0$, if $r \geq 0$,*

$$P\{\inf(U, V) > a\} \leq P\{U > a\}\psi\left(a\sqrt{\frac{1-r}{1+r}}\right).$$

Let $0 < \alpha < \rho^2(1 - M^{-1/2})/2$ be fixed and suppose first

a) $m_0 \leq p < q \leq p + p^\alpha$,

then by Lemma 3.4,

$$(3.27) \quad P\{A''_p \cap A''_q\} \leq P\{A''_p\} \psi(\rho \sqrt{(1 - M^{-1/2}) \log(p + 2)}),$$

since $E \left[\frac{W(n_p^*) W(n_q^*)}{\sqrt{n_p^*} \sqrt{n_q^*}} \right] \leq M^{1 - (q-p)/2} \leq M^{-1/2}, \quad \text{if } q > p + 1.$

Thus,

$$P\{A''_p \cap A''_q\} \leq P\{A''_p\} (p + 2)^{-\rho^2(1 - M^{-1/2})/2},$$

so that,

$$(3.28) \quad \sum_{p < q \leq p + p^\alpha} P\{A''_p \cap A''_q\} \leq P\{A''_p\} \{ [p + 2]^{\alpha - \rho^2(1 - M^{-1/2})/2} + 1 \}, \\ \leq (1 + h) P\{A''_p\},$$

once m_0 is sufficiently large.

b) $m_0 \leq p \leq p + p^\alpha < q.$

Fix $\varepsilon \in]\alpha, 1[.$ Then assuming m_0 large enough, one has $q - p \geq q^\varepsilon$, so that

$$(3.29) \quad \sup \left[\varphi(n_p^*) \varphi(n_q^*) E \left\{ \frac{W(n_p^*) W(n_q^*)}{\sqrt{n_p^*} \sqrt{n_q^*}} \right\}, p > m_0, q > p + p^\alpha \right] \\ \leq 2 \sup [(\log p + 2)(\log q + 2) M^{2 - (q-p)/2}], p > m_0, q > p + p^\alpha, \\ \leq 2 \sup [(\log q + 2) M^{(1 - q^\varepsilon)/2}], q \geq m_0.$$

By virtue of Lemma 3.4, this one leads to

$$(3.30) \quad P\{A''_p \cap A''_q\} \leq (1 + h) P\{A''_p\} P\{A''_q\},$$

once m_0 is sufficiently large, that we suppose; thus (3.23) is now established. As for (3.24), fix $1 > \rho'' > \rho' > \rho.$ For m_0 large enough,

$$(3.31) \quad \sum_{p=m_0}^m P\{A''_p\} \geq \frac{1}{\sqrt{2\pi}} \sum_{p=m_0}^m \frac{1}{1 + \rho \sqrt{2 \log(p + 2)}} [p + 2]^{-\rho^2}, \\ \geq \sum_{p=m_0}^m [p + 2]^{-(\rho')^2}, \\ \geq \{ [m + 3]^{1 - (\rho')^2} - [m_0 + 2]^{1 - (\rho')^2} \} / [1 - (\rho')^2], \\ \geq m^{1 - (\rho'')^2},$$

once $m > m_1 = m_1(m_0, \rho', \rho'').$ Hence (3.24) holds and consequently our first step, which is (3.25) is now established. Letting $\mathcal{E} = N$ and ρ_0 tending to 1 in (3.25) leads to (3.13').

Step 2. In this step we prove (3.14). Obviously, there is no loss when assuming $\Lambda(\mathfrak{s}) > 0.$ Fix $0 < \delta' < \Lambda(\mathfrak{s})^2,$ and set

$$(3.32) \quad \mathcal{E} = \{m \in N: k_m \leq m^{1/\delta'}\}.$$

Since
$$A(\mathfrak{s})^2 = \limsup_{m \rightarrow \infty} [\log m] / [\log k_m] > \delta',$$
 (3.33)
$$\#(\mathfrak{E}) = \infty.$$

Fix also $0 < \eta < 1 - \rho_1^2$. Then, according to (3.25), with probability one, there are infinitely many integers $m \in \mathfrak{E}$, such that,

(3.34) A_{k_p} occurs for at least $m^{1-\rho_1^2} - m^\eta$ integers p in the interval $[m^\eta, m]$, since, one has

$$\sum_{1 \leq p \leq m} I_{A_{k_p}} \leq m^\eta + \sum_{m^\eta < p \leq m} I_{A_{k_p}}.$$

For such integers, by definition of A_{k_p} , one has

$$S_n(X) \geq \rho_0 \sqrt{n} \varphi(n) \quad \text{for some } n \in \mathfrak{s} \cap I_{k_p},$$

and

$$\varphi(n) = \sqrt{2 \log(p + 2)}.$$

But $p \geq m^\eta$, $n \leq M^{k_{p+1}} \leq M^{k_m+1}$ and $k_m \leq m^{1/\delta'}$; hence

$$\begin{aligned} \log(p + 2) &\geq \log \left[2 + \left(\frac{\log(n/M)}{\log M} \right)^{\delta' \eta} \right], \\ &\geq [\log \log n] \delta' \eta', \end{aligned}$$

for any $0 < \eta' < \eta$, once n is sufficiently large, namely, once m is large enough. Thus, we have

(3.35)
$$\varphi(n) \geq \sqrt{2 \delta' \eta' \log \log n}.$$

On the whole, with probability one, there are infinitely many integers m such that

(3.36)
$$\left\{ \begin{array}{l} \text{there exists at least } m^{1-(\rho_1)^2} - m^\eta \text{ integers } p \in [m^\eta, m] \text{ such that} \\ S_n(X) \geq \rho_0 \sqrt{n} \varphi(n) \text{ and } \varphi(n) \geq \sqrt{2 \delta' \eta' \log \log n} \text{ for some } n \in \mathfrak{s} \cap I_{k_p}. \end{array} \right.$$

Letting $\delta' \eta' = A^2(\mathfrak{s}) \eta''$, gives

(3.37)
$$P \left\{ \sum_{1 \leq p \leq m} I_{\{\mathfrak{s}(\rho_0, \eta'', X) \cap I_{k_p}\}} \geq m^{1-(\rho_1)^2} - m^\eta, m \in \mathfrak{E}, m \text{ i.o.} \right\} = 1.$$

Since

$$\sum_{1 \leq p \leq m} I_{\{\mathfrak{s}(\rho_0, \eta'', X) \cap I_{k_p}\}} = \sum_{1 \leq i \leq k_m} I_{\{\mathfrak{s}(\rho_0, \eta'', X) \cap [M^i, M^{i+1}[}},$$

(3.37) together with the definition of $A(\mathfrak{s})$ imply

(3.38)
$$P\{A(\mathfrak{s}(\rho_0, \eta'', X)) \geq \sqrt{1 - (\rho_1)^2} A(\mathfrak{s})\} = 1.$$

This one easily establishes (3.14).

Step 3. We now prove (3.15). Let $X^i = \{X_n^i, n \geq 1\}$, $i = 1, 2$, be two independent copies of the sequence $X = \{X_n, n \geq 1\}$, and define

$$(3.39) \quad c_1 = \max \{ \rho \sqrt{1 - \rho^2}, 0 < \rho < 1 \} = 1/2.$$

From (3.14) and (3.39),

$$(3.40) \quad \limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{S_n(X)}{\sqrt{2n \log \log n}} \geq \max_{\substack{0 < \rho < 1 \\ 0 < \eta < 1 - \rho^2}} \left[\liminf_{\mathfrak{s}(\rho, \eta, X) \ni n \rightarrow \infty} \frac{S_n(X)}{\sqrt{2n \log \log n}} \right] \geq (1/2)A(\mathfrak{s}),$$

almost surely. This is the first step. We now apply (3.40) and (3.14) simultaneously. We first observe, for every $0 < \alpha < 1$,

$$(3.41) \quad \begin{aligned} \limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{S_n(\alpha X^1 + X^2 \sqrt{1 - \alpha^2})}{\sqrt{2n \log \log n}} &\geq \limsup_{\mathfrak{s}(\rho, \eta, X^2) \ni n \rightarrow \infty} \frac{S_n(\alpha X^1 + X^2 \sqrt{1 - \alpha^2})}{\sqrt{2n \log \log n}}, \\ &\geq \left[\liminf_{\mathfrak{s}(\rho, \eta, X^2) \ni n \rightarrow \infty} \frac{S_n(X^2)}{\sqrt{2n \log \log n}} \right] \sqrt{1 - \alpha^2} \\ &\quad + \alpha \left[\limsup_{\mathfrak{s}(\rho, \eta, X^2) \ni n \rightarrow \infty} \frac{S_n(X^1)}{\sqrt{2n \log \log n}} \right], \end{aligned}$$

almost surely, and thus

$$(3.42) \quad \geq \left[\sqrt{1 - \alpha^2}(\rho \sqrt{\eta}) + \frac{\alpha}{2} \sqrt{1 - \rho^2} \right] A(\mathfrak{s}),$$

by virtue of (3.40) (and 3.14) since X^1 and X^2 are independent. We put

$$(3.43) \quad \forall n \geq 1, \quad c_n = \max \{ [\rho \sqrt{1 - \alpha^2} + \alpha c_{n-1}] \sqrt{1 - \rho^2}, 0 < \rho < 1, 0 < \alpha < 1 \}.$$

From (3.42), assuming that X is gaussian,

$$(3.44) \quad P \left\{ \limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{S_n(X)}{\sqrt{2n \log \log n}} \geq c_2 A(\mathfrak{s}) \right\} = 1,$$

and, by repeating the same argument,

$$(3.45) \quad P \left\{ \limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{S_n(X)}{\sqrt{2n \log \log n}} \geq \sup_p (c_p) A(\mathfrak{s}) \right\} = 1.$$

Now, the lower bound in (3.25) will be deduced from the study of the sequence $\{c_n, n \geq 1\}$. It is easy to check that the maximum of

$$\varphi_{\rho, n}(\alpha) = \{ \rho \sqrt{1 - \alpha^2} + \alpha c_{n-1} \} \sqrt{1 - \rho^2}$$

is reached at the value

$$\alpha(\rho, n) = c_{n-1} / \sqrt{\rho^2 + (c_{n-1})^2}$$

and

$$f_n(\rho) = \varphi_{\rho, n}(\alpha(\rho, n)) = \sqrt{[1 - \rho^2][(c_{n-1})^2 + \rho^2]}.$$

Further, $f_n(\cdot)$ has unique maximum on $[0, 1]$ at the value $\rho(n) = \sqrt{[1 - (c_{n-1})^2]/2}$. Its corresponding value defines c_n and finally, one has

$$(3.46) \quad \forall n \geq 1, \quad c_n = [1 + (c_{n-1})^2]/2.$$

It is now easy to see that c_n increases to 1 when n tends to infinity. This gives the lower bound in (3.15) when X is gaussian.

We now must prove this in general. First, it is quite clear that $A(\varepsilon) = A(\varepsilon^*)$. Replacing ε by ε^* , our result becomes

$$(3.47) \quad P\left\{\limsup_{p \rightarrow \infty} \frac{W(n_p^*)}{\sqrt{2n_p^* \log \log n_p^*}} \geq A(\varepsilon)\right\} = 1.$$

By virtue of the Skohorod embedding scheme and (3.18), for any integer p and $0 < h < 1$,

$$(3.48) \quad \begin{aligned} P\{A \cap \{|W(\sum_{i=1}^{n_p^*} T_i) - W(n_p^*)| / \sqrt{n_p^*} > \sqrt{2h \log \log n_p^*} + \varepsilon(h) + \mu\sqrt{h}\}\} \\ \leq P\{\sup(|W(\theta) - W(1)|, 1 - h \leq \theta \leq 1 + h) \\ > \sqrt{2h \log \log n_p^*} + \varepsilon(h) + \mu\sqrt{h}\}, \\ \leq 2\psi(\sqrt{2 \log \log n_p^*} + \mu), \quad (\text{by applying Borell's inequality}), \\ \leq 2\psi(\sqrt{2 \log [p(\log M)]} + \mu), \quad (\text{since } n_p^* \geq M^{k_p} \geq M^p). \end{aligned}$$

Therefore, by applying Borel-Cantelli lemma and letting h tend to 0,

$$(3.49) \quad P\left\{\lim_{p \rightarrow \infty} \frac{|W(\sum_{i=1}^{n_p^*} T_i) - W(n_p^*)|}{\sqrt{2n_p^* \log \log n_p^*}} = 0\right\} = 1.$$

Putting together (3.47) and (3.49) leads to

$$(3.50) \quad P\left\{\limsup_{p \rightarrow \infty} \frac{S_{n_p^*}(X)}{\sqrt{2n_p^* \log \log n_p^*}} \geq A(\varepsilon)\right\} = 1,$$

which produces the lower bound in (3.15).

We now turn to the upper bound of (3.15). Set,

$$(3.51) \quad L(X) = \limsup_{\varepsilon \in n \rightarrow \infty} \frac{S_n(X)}{\sqrt{2n \log \log n}}.$$

By 0-1 law, $L(X)$ is a number and there is no loss when assuming $L(X)$

> 0 . Let $0 < L' < L(X)$ be fixed. Then for some random subsequence $\{n_j, j \geq 1\}$ of \mathfrak{s} ,

$$(3.52) \quad \forall j \geq 1, \quad L' \leq \frac{S_{n_j}(X)}{\sqrt{2n_j \log \log n_j}}.$$

Define the random sequence of integers $\{p_j, j \geq 1\}$ such that

$$(3.53) \quad \forall j \geq 1, \quad n_j \in I_{k_{p_j}}.$$

Fix $h > 0$ arbitrary. By Lemma 3.2, for all large enough j ,

$$(3.54) \quad S_{n_j}(X) \leq \sqrt{(1+h)n_j \varphi(n_j)} = \sqrt{(1+h)n_j 2 \log(p_j + 2)},$$

so that,

$$(3.55) \quad \begin{aligned} L' &\leq \left[\frac{\log(p_j + 2)}{\log \log n_j} (1+h) \right]^{1/2}, \\ &\leq \left[\frac{(1+h) \log(p_j + 2)}{\log[(\log M)k_{p_j}]} \right]^{1/2}, \\ &\leq \left[\frac{(1+h) \log \{2 + \sum_{i=1}^{k_{p_j}} \delta_i(\mathfrak{s}, M)\}}{\log[(\log M)k_{p_j}]} \right]^{1/2}. \end{aligned}$$

Therefore,

$$(3.56) \quad L' \leq \sqrt{1+h} \limsup_{p \rightarrow \infty} \left[\frac{\log \sum_{i=1}^p \delta_i(\mathfrak{s}, M)}{\log p} \right]^{1/2}.$$

Letting L' tend to L and h tend to 0 gives the conclusion. Thus (3.15) is proved, and the proof of the Theorem 3.3 is now complete. Q.E.D.

The first part of the above proof is very classical in the study of upper or lower classes of gaussian sequences; part 2 and part 3 are the original parts of the proof.

LEMMA 3.5. *For any triple $(\mathfrak{s}, M, \varphi)$ defined in accordance with (2.1) and any sequence $X = \{X_i, i \geq 1\}$ of i.i.d. real r.v.'s satisfying $EX_1 = 0$ and $E(X_1)^2 < \infty$,*

$$(3.57) \quad P\{C(\{S_n(X)/\sqrt{n} \varphi(n), n \in \mathfrak{s}\}) = [-\sqrt{E(X_1)^2}, \sqrt{E(X_1)^2}]\} = 1.$$

Proof. The proof is very simple. First, by homogeneity there is no loss when assuming $E(X_1)^2 = 1$. Then, let $\delta \geq 1$ be fixed. We associate to it the following subsequence of \mathfrak{s}^* ,

$$(3.58) \quad \mathfrak{s}_\delta^* = \{n_{[p^\delta]}^*, p \geq 1\}.$$

The corresponding sequence $k_p^{\delta} = k_p(\mathfrak{S}_\delta^*)$, $p \geq 1$, trivially satisfies,

$$(3.58') \quad \forall p \geq 1, \quad k_p^{\delta} = k_{\lfloor p^{\delta} \rfloor},$$

so that the corresponding function φ_δ also satisfies

$$(3.58'') \quad \forall n \in \mathfrak{S}_\delta^*, \quad \varphi_\delta(n) = \sqrt{2 \log(p+2)} \quad \text{iff } n \in \mathfrak{S}_\delta^* \cap I_{k_{\lfloor p^{\delta} \rfloor}},$$

and therefore $\varphi_\delta(n) \approx \varphi(n)\sqrt{\delta}$ as n tends to infinity along the sequence \mathfrak{S}_δ^* . By applying (3.13'), one obtains

$$(3.59) \quad P \left\{ \limsup_{\mathfrak{S}_\delta^* \ni n \rightarrow \infty} \frac{S_n(X)}{\sqrt{n} \varphi(n)} = \delta^{-1/2} \right\} = 1.$$

This one easily implies (3.57). Q.E.D.

The next lemma shows the necessity of the condition $E(X_1)^2 < \infty$. The proof uses a classical argument on truncated r.v.'s.

LEMMA 3.6. *Let $(\mathfrak{S}, M, \varphi)$ be any triple defined in accordance with (2.1). Let also $X = \{X_i, i \geq 1\}$ be a sequence of i.i.d. real r.v.'s. Then,*

$$(3.60) \quad P \left\{ \limsup_{\mathfrak{S} \ni n \rightarrow \infty} \frac{S_n(X)}{\sqrt{n} \varphi(n)} = 1 \right\} = 1,$$

if, and only if,

$$(3.61) \quad EX_1 = 0 \quad \text{and} \quad E(X_1)^2 = 1.$$

Proof. There is just the "only if" part of the assertion to prove. First, assume that X_1 is a symmetric r.v., and let $c > 0$ fixed. Then,

$$X'_1 = X_1 I_{\{|X_1| \leq c\}} - X_1 I_{\{|X_1| > c\}},$$

has same distribution as X_1 . Let $X' = \{X'_i, i \geq 1\}$ be a sequence of independent copies of X'_1 . By writing (3.60) for X and X' , then using triangular inequality, one obtains,

$$(3.62) \quad P \left\{ \limsup_{\mathfrak{S} \ni n \rightarrow \infty} \frac{S_n(X I_{\{|X_1| \leq c\}})}{\sqrt{n} \varphi(n)} \leq 1 \right\} = 1.$$

By (3.13'),

$$(3.62') \quad P \left\{ \limsup_{\mathfrak{S} \ni n \rightarrow \infty} \frac{S_n(X I_{\{|X_1| \leq c\}})}{\sqrt{n} \varphi(n)} = \sqrt{E\{X_1^2 I_{\{|X_1| \leq c\}}\}} \right\} = 1,$$

so that,

$$(3.62'') \quad E\{X_1^2 I_{\{|X_1| \leq c\}}\} \leq 1,$$

and finally $EX_1^2 \leq 1$. But, by using again (3.13') and (3.60), this, necessarily implies $EX_1^2 = 1$. If X_1 is not symmetric, let X' be an independent copy of X , and set $Y = X - X'$. Writing (3.60) for X and X' , and using triangular inequality, leads to

$$(3.63) \quad P\left\{\limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{S_n(Y)}{\sqrt{n} \varphi(n)} \leq 2\right\} = 1.$$

Therefore, $E|X_1 - X_1'|^2 \leq 4$, and by Corollary 2 (M. Loève, Probability theory, 3^e ed., p. 246),

$$(3.64) \quad E|X_1 - \mu X_1|^2 \leq 8,$$

where μX_1 denotes a median of X_1 . This implies $E|X_1|^2 < \infty$. By applying (3.13') and (3.60), one obtains

$$(3.65) \quad E(X_1)^2 = 1.$$

Further, $EX_1 = m < \infty$. The centering of X_1 follows from the strong law of large numbers, since

$$\text{a.s.} \quad |m| = \lim_{\mathfrak{s} \ni n \rightarrow \infty} \left| \frac{S_n(X)}{n} \right| \leq \limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{|S_n(X)|}{\sqrt{n} \varphi(n)} \cdot \limsup_{\mathfrak{s} \ni n \rightarrow \infty} \frac{\varphi(n)}{\sqrt{n}} = 0. \quad \text{Q.E.D.}$$

Proof of Theorem 2.1.

a) if $m = 1$. This is given by (3.13') in Theorem 3.3, Lemmas 3.5 and 3.6.

b) if $1 \leq m < \infty$. The sufficiency results from the proof of Lemma 2 in [5], in which the sequence of normalization constants $\{\sqrt{2n \log \log n}, n \geq 3\}$ does not matter except the fact that it is needed to control the case $m = 1$.

As for the necessity, let $f_i, f_{i,j}, i, j = 1, \dots, m$, be linear forms on \mathbf{R}^m defined by $f_i(x) = x^i$, and $f_{i,j}(x) = x^i + x^j$ for every $x = (x^1, \dots, x^m) \in \mathbf{R}^m$. By applying Lemma 3.5 to the sequences $\{f_{i,j}(S_n(X)), n \in \mathfrak{s}\}, i, j = 1, \dots, m$, one obtains

$$(3.66) \quad \forall i, j = 1, \dots, m, \quad \begin{aligned} Ef_{i,j}(X_1)^2 &= E[X_1^i + X_1^j]^2 = 2, \\ Ef_i(X_1)^2 &= E[X_1^i]^2 = 1, \\ Ef_i(X_1) &= EX_1^i = 0, \end{aligned}$$

which easily allows to conclude. Q.E.D.

Proof of Theorem 2.3. As far as this is just the assertion (3.15) of Theorem 3.3, it is already proved. Q.E.D.

Proof of Theorem 2.8.

a) Lower bound. Fix $0 < \rho_0 < \rho_1 \leq 1$, and set,

$$(3.67) \quad \begin{cases} \forall p \geq 1, & k_p^1 = k_p^1(\mathfrak{s}_1, M), & (k_p = k_p(\mathfrak{s}, M)), \\ \forall n \in \mathfrak{s}_1, & \varphi_1(n) = \varphi(\mathfrak{s}_1, M, n), & \text{(using notations (2.1))}, \\ \forall p \geq 1, & A_p = \{\exists n \in \mathfrak{s}_1 \cap [M^p, M^{p+1}[: S_n(X) \geq \rho_0 \sqrt{n} \varphi_1(n)\}. \end{cases}$$

Since \mathfrak{s}_1 is a subsequence of \mathfrak{s} , then $\{k_p^1, p \geq 1\}$ is a subsequence of $\{k_p, p \geq 1\}$, and

$$(3.68) \quad \forall p \geq 1, \quad k_p^1 = k_{c(p)},$$

where $c: N \rightarrow N$ is strictly increasing ($c(p) \geq p$, for every p).

There is no loss when assuming $\Lambda(\mathfrak{s}, M, \mathfrak{s}_1) > 0$. Observe that

$$(3.69) \quad \begin{aligned} \Lambda(\mathfrak{s}, M, \mathfrak{s}_1) &= \limsup_{m \rightarrow \infty} \left[\frac{\log m}{\log \#\{i \leq k_m^1 : \mathfrak{s} \cap [M^i, M^{i+1}[\neq \emptyset\}} \right]^{1/2}, \\ &= \limsup_{m \rightarrow \infty} \left[\frac{\log m}{\log c(m)} \right]^{1/2}. \end{aligned}$$

Fix $0 < \delta < \Lambda^2(\mathfrak{s}, M, \mathfrak{s}_1)$, and set

$$\mathcal{E} = \{m \in N : c(m) \leq m^{1/\delta}\}.$$

Then $\#\mathcal{E} = \infty$. Further, by (3.25)

$$(3.70) \quad P\{\sum_{i=1}^m I_{A_{k_p^1}} \geq m^{1-(\rho_1)^2}, m \in \mathcal{E}, m \text{ i.o.}\} = 1.$$

Fix $0 < \eta < 1 - (\rho_1)^2$. With probability one, there are infinitely many $m \in \mathcal{E}$ such that

$$(3.71) \quad A_{k_p^1} \text{ occurs for at least } m^{1-(\rho_1)^2} - m^\eta \text{ integers } p \in [m^\eta, m].$$

For these integers one has

$$S_n(X) \geq \rho_0 \sqrt{n} \varphi_1(n) \quad \text{for some } n \in \mathfrak{s}_1 \cap I_{k_p^1}(M),$$

and
$$\varphi_1(n) = \sqrt{2 \log(p+2)}.$$

But $m^\eta \leq p \leq m$, and $M^{k_p^1} = M^{k_{c(p)}} \leq n < M^{k_p^1+1} = M^{k_{c(p)+1}}$, so that

$$\varphi(n) = \sqrt{2 \log(c(p) + 2)} \leq \sqrt{2 \log(c(m) + 2)}.$$

Thus,

$$(3.72) \quad \begin{aligned} \varphi_1(n) &\geq \sqrt{2 \log(2 + m^\eta)}, \\ &\geq \sqrt{2 \log(2 + c(m)^{\delta\eta})}, \\ &\geq \sqrt{\delta\eta} \varphi(n), \end{aligned}$$

for any $0 < \eta' < \eta$, once n is large enough. Therefore, with probability one, there are infinitely many $m \in \mathcal{E}$ such that

there exist at least $m^{1-(\rho_1)^2} - m^\nu$ integers $p \in [m^\nu, m]$ for which $S_n(X) \geq \rho_0 \sqrt{n} \varphi_1(n)$ and $\varphi_1(n) \geq \sqrt{\delta \eta'} \varphi(n)$ for some $n \in \mathcal{E}_1 \cap I_{k_p}^1$.

By arguing along the lines (3.36)–(3.38), one obtains for,

$$(3.73) \quad \mathcal{E}_1(\rho_0, \eta'', X) = \{n \in \mathcal{E}_1: S_n(X) \geq \rho_0 \sqrt{n} \varphi_1(n) \text{ and } \varphi_1(n) \geq \sqrt{\eta''} \Lambda(\mathcal{E}, M, \mathcal{E}_1) \varphi(n)\},$$

$$P\{\Lambda(\mathcal{E}, M, \mathcal{E}_1(\rho_0, \eta'', X)) \geq \sqrt{1 - (\rho_1)^2} \Lambda(\mathcal{E}, M, \mathcal{E}_1)\} = 1,$$

where we put $\delta \eta' = \eta'' \Lambda^2(\mathcal{E}, M, \mathcal{E}_1)$. Next, we conclude by following the same scheme of proof as in step 3 of the proof of Theorem 3.3.

b) Upper bound. Again there is no loss when assuming that

$$L(X) = \limsup_{\mathcal{E}_1 \ni n \rightarrow \infty} \frac{S_n(X)}{\sqrt{n} \varphi(n)} > 0.$$

Fix $0 < L' < L(X)$, and let $h > 0$. Then for some random subsequence $\{n_j, j \geq 1\}$ of \mathcal{E}_1 ,

$$\forall j \geq 1, \quad L' \leq S_{n_j}(X) / \sqrt{n_j} \varphi(n_j).$$

One defines two random sequences of integers $\{q_j, j \geq 1\}$ and $\{p_j, j \geq 1\}$ such that

$$\forall j \geq 1, \quad n_j \in I_{k_p}^1(M) \cap I_{k_q}^1(M).$$

By Lemma 3.2,

$$S_{n_j}(X) \leq (1 + h) \sqrt{2n_j \log(p_j + 2)},$$

for all j large enough, and $\varphi(n_j) = \sqrt{2 \log(q_j + 2)}$. Thus,

$$L' \leq (1 + h) \left[\frac{\log(p_j + 2)}{\log(q_j + 1)} \right]^{1/2},$$

for all j large enough. This one easily leads to the result by letting L' tend to $L(X)$ and h tend to 0. Q.E.D.

§ 4. Strassen's laws of the iterated on subsequences

First, we recall the following lemma due to J. Kuelbs ([9], p. 247–248).

LEMMA 4.1. 1) *Let B a separable Banach space. Let $\{Y_k, k \geq 1\}$ be a sequence of B -valued random variables and assume μ is a mean zero*

Gaussian measure on B . Let K denote the unit ball of the reproducing Hilbert space H_μ of μ . If,

$$(4.1) \quad L((Y_k), \mu) = b_k \quad (k \geq 1),$$

where $\sum_{k=1}^\infty b_k < \infty$, and L is the Prohorov metric for probability measures on $(B, \|\cdot\|)$. Then,

$$(4.2) \quad P\left\{\omega: \lim_{n \rightarrow \infty} d\left(\frac{Y_n(\omega)}{\sqrt{2 \log n}}, K\right) = 0\right\} = 1,$$

where $d(x, A) = \inf \{\|x - y\|, y \in A\}$.

2) If $(Y_k) = \mu$ for every k , and

$$(4.3) \quad \lim_{\substack{m \rightarrow \infty \\ k-m \rightarrow \infty}} E\{E[f(Y_k) | \underline{F}_m]^2\} = 0,$$

for every $f \in B^*$ where $\underline{F}_m = \underline{F}\{Y_k, k \leq m\}$; then

$$(4.4) \quad P\left\{\omega: C\left(\left\{\frac{Y_n(\omega)}{\sqrt{2 \log n}}, n \geq 1\right\}\right) = K\right\} = 1.$$

Let W be a 1-dimensional brownian motion and define for every integer n

$$(4.5) \quad \zeta_n(t) = \frac{W(nt)}{\sqrt{n}} \quad (0 \leq t \leq 1).$$

Clearly each $\{\zeta_n(t), 0 \leq t \leq 1\}$ is a brownian motion on $[0, 1]$ and it induces a Wiener measure μ on $\underline{C}([0, 1])$. Further, it is well known that the reproducing kernel Hilbert space of μ is

$$H_\mu = \left\{f \in \underline{C}([0, 1]): f(t) = \int_0^t g(u)du \text{ where } \int_0^1 g^2(s)ds < \infty\right\},$$

with inner product $\langle f_1, f_2 \rangle = \int_0^1 f_1'(u)f_2'(u)du$ and hence

$$K = \left\{f \in \underline{C}([0, 1]): f(t) = \int_0^t g(u)du \text{ where } \int_0^1 g^2(u)du \leq 1\right\}.$$

Let now $(\mathfrak{s}, M, \varphi)$ be defined in accordance with (2.1) and set

$$(4.5') \quad \forall p \geq 1, \quad \zeta_p^* = \zeta_{n_p^*}.$$

LEMMA 4.2. For any triple $(\mathfrak{s}, M, \varphi)$

$$a) \quad P\left\{\lim_{n \rightarrow \infty} d\left(\frac{\zeta_{n_p^*}}{\sqrt{2 \log(p+2)}}, K\right) = 0\right\} = 1,$$

$$b) P\left\{C\left\{\frac{\zeta_{n_p^*}}{\sqrt{2\log(p+2)}}, p \geq 1\right\} = K\right\} = 1.$$

Proof. We mimic Kuelbs’s proof in [9] p. 249–251. Since $L((\zeta_p^*), \mu) = 0$, a) is easily deduced from the first part of Lemma 4.1. To prove b) if $f \in C^*([0, 1])$ with $f(x) = \int_0^1 x(t)dF(t)$, for $x \in C([0, 1])$, then

$$\begin{aligned} (4.6) \quad E\{f(\zeta_{p+q}^*) | \underline{E}_p\} &= E\left\{\int_0^1 \frac{W(n_{p+q}^*t)}{\sqrt{n_{p+q}^*}} dF(t) | \underline{E}_p\right\}, \\ &= E\left\{\int_0^{n_p^*/n_{p+q}^*} \frac{W(n_{p+q}^*t)}{\sqrt{n_{p+q}^*}} dF(t) + \int_{n_p^*/n_{p+q}^*}^1 \frac{W(n_{p+q}^*t)}{\sqrt{n_{p+q}^*}} dF(t) | \underline{E}_p\right\} \\ &= \int_0^{n_p^*/n_{p+q}^*} \frac{W(n_{p+q}^*t)}{\sqrt{n_{p+q}^*}} dF(t) + \int_{n_p^*/n_{p+q}^*}^1 \frac{W(n_p^*)}{\sqrt{n_{p+q}^*}} dF(t). \end{aligned}$$

Hence,

$$\begin{aligned} E\{E[f(\zeta_p^*) | \underline{E}_p]^2\} &= \frac{1}{n_{p+q}^*} \int_0^{n_p^*/n_{p+q}^*} \int_0^{n_p^*/n_{p+q}^*} \min[n_{p+q}^*s, n_{p+q}^*t] dF(s)dF(t) \\ &\quad + \frac{2}{n_{p+q}^*} \int_0^{n_p^*/n_{p+q}^*} \int_{n_p^*/n_{p+q}^*}^1 \min[n_p^*, n_{p+q}^*t] dF(s)dF(t) \\ &\quad + \frac{n_p^*}{n_{p+q}^*} \left(\int_{n_p^*/n_{p+q}^*}^1 dF(t)\right)^2, \end{aligned}$$

and (4.3) holds since $\lim_{q \rightarrow \infty} \sup_{p \geq 1} n_p^*/n_{p+q}^* = 0$. Thus, the second part of Lemma 4.1 gives b). Q.E.D.

(4.6’) Lemma 4.2 then holds for any subsequence $\mathfrak{s}' \subset \mathfrak{s}$ such that $\#(\mathfrak{s}' \cap I_{k_p}) = 1$, for every p . This simple observation will be afterwards convenient.

THEOREM 4.3. *For any triple $(\mathfrak{s}, M, \varphi)$ defined in accordance with (2.1),*

- a) $P\left\{\lim_{\mathfrak{s} \ni n \rightarrow \infty} d\left(\frac{\zeta_n}{\varphi(n)}, K\right) = 0\right\} = 1,$
- b) $P\left\{C\left(\left\{\frac{\zeta_n}{\varphi(n)}, n \in \mathfrak{s}\right\}\right) = K\right\} = 1.$

Proof. Fix $h > 0$ and define

$$(4.7) \quad \begin{cases} \forall p \geq 1, & J(\mathfrak{s}, k_p, k) = \mathfrak{s} \cap [M^{k_p}(1 + (k - 1)h), M^{k_p}(1 + kh)[, \\ & \text{where } k \in \mathcal{A}(\mathfrak{s}, p) \text{ and} \\ \forall p \geq 1, & \mathcal{A}(\mathfrak{s}, p) = \{1 \leq j \leq (M - 1)/h: J(\mathfrak{s}, k_p, j) \neq \emptyset\}, \end{cases}$$

$$\left\{ \begin{array}{l} \beta = 4E\{\sup\{|W(r)|, 1 \leq r \leq M\}\}, \\ \forall p \geq 1, \quad \forall k \in \mathcal{A}(\mathfrak{s}, p), \quad n_{p,k}^* \text{ is the first point of } J(\mathfrak{s}, k_p, k), \\ \forall p \geq 1, \quad \forall k \in \mathcal{A}(\mathfrak{s}, p), \quad \zeta_{p,k}^* = \zeta_{n_{p,k}^*}^*. \end{array} \right.$$

Fix $k \in [1, (M-1)/h]$ and set for each p such that $k \in \mathcal{A}(\mathfrak{s}, p)$,

$$(4.8) \quad A_p^k = \{\exists n \in J(\mathfrak{s}, k_p, k): \|\zeta_n - \zeta_{p,k}^*\| \geq \sqrt{2h \log(p+2)} + 2\beta\}.$$

Observe now, for $n \in J(\mathfrak{s}, k_p, k)$ and $0 \leq \theta \leq 1$,

$$(4.9) \quad E \left| \frac{W(n\theta)}{\sqrt{n}} - \frac{W(n_{p,k}^*\theta)}{\sqrt{n_{p,k}^*}} \right|^2 = 2\theta(1 - n_{p,k}^*/n), \\ \leq 2h,$$

and,

$$(4.10) \quad E\{\sup\{\|\zeta_n - \zeta_{p,k}^*\|, n \in J(\mathfrak{s}, k_p, k)\}\} \\ \leq E\{\sup\{\|\zeta_n\|, n = M^{k_p}\lambda, 1 \leq \lambda \leq M\}\} + E\{\|\zeta_{p,k}^*\|\}, \\ \leq \beta/2.$$

By applying Borell's inequality,

$$(4.11) \quad \forall p \geq 1, \quad P\{A_p^k\} \leq 2\nu(\sqrt{2 \log(p+2)} + \beta/\sqrt{h}),$$

so that, for each $k \in [1, (M-1)/h]$,

$$(4.12) \quad P\{\{\exists n \in J(\mathfrak{s}, k_p, k): \|\zeta_n - \zeta_{p,k}^*\| > \sqrt{2h \log(p+2)} + 2\beta\}, p \text{ i.o.}\} = 0.$$

Therefore,

$$(4.13) \quad P\left\{\limsup_{p \rightarrow \infty} \sup\left\{\frac{\|\zeta_n - \zeta_{p,k}^*\|}{\varphi(n)}, n \in J(\mathfrak{s}, k_p, k)\right\} \leq \sqrt{h}\right\} = 1.$$

But Lemma 4.2 and remark (4.6') imply,

$$(4.14) \quad \forall k \in [1, (M-1)/h], \quad P\{\lim_{p \rightarrow \infty} d(\zeta_{p,k}^*/\sqrt{2 \log(p+2)}, K) = 0\} = 1,$$

and,

$$(4.15) \quad \forall k \in [1, (M-1)/h], \quad P\{C(\{\zeta_{p,k}^*/\sqrt{2 \log(p+2)}, p \geq 1\}) = K\} = 1.$$

Combining (4.13) with (4.14), then letting h tend to 0

$$(4.16) \quad P\{\lim_{\mathfrak{s} \ni n \rightarrow \infty} d(\zeta_n/\varphi(n), K) = 0\} = 1.$$

Combining (4.15) with (4.16),

$$(4.17) \quad P\{C(\{\zeta_n/\varphi(n), n \in \mathfrak{s}\}) = K\} = 1. \quad \text{Q.E.D.}$$

LEMMA 4.4. For any triple $(\mathfrak{s}, M, \varphi)$ defined in accordance with (2.1) such that

$$(4.19) \quad \lim_{k \rightarrow \infty} n_k/n_{k+1} = 1, \quad (\mathfrak{s} = \{n_k, k \geq 1\}),$$

and for any sequence $X = \{X_i, i \geq 1\}$ of i.i.d. real r.v.'s satisfying $EX_1 = 0$ and $E(X_1)^2 = 1$, one has

$$(4.20) \quad P\left\{\lim_{j \rightarrow \infty} \left\| \frac{\eta_j}{\sqrt{n_j}} - \zeta_{n_j} \right\| / \varphi(n_j) = 0\right\} = 1,$$

where $\eta_j, j \geq 1$, is defined in (2.24) relatively to X and \mathfrak{s} .

Proof. By virtue of the Skohorod embedding scheme for partial sums, it is enough to prove the Lemma 4.4 when replacing X_i ($i \geq 1$), by

$$\tilde{X}_i = W(T_1 + \cdots + T_i) - W(T_1 + \cdots + T_{i-1}) \quad (i \geq 1),$$

where W is a 1-dimensional brownian motion and T_1, T_2, \dots a sequence of nonnegative i.i.d. r.v.'s satisfying $ET_1 = 1$. Define, with the convention $n_0 = 0$,

$$(4.21) \quad \forall \theta \geq 0, \text{ if } n_k \leq \theta \leq n_{k+1} \text{ for some } k \geq 0, \\ \tilde{\gamma}(\theta) = \tilde{S}_{n_k}[1 + (n_k - \theta)/(n_{k+1} - n_k)] + \tilde{S}_{n_{k+1}}[(\theta - n_k)/(n_{k+1} - n_k)], \\ \text{where } \tilde{S}_{n_k} = \sum_{i=1}^{n_k} \tilde{X}_i \quad (k \geq 1).$$

Then,

$$(4.22) \quad |\tilde{\gamma}(t) - W(t)| \leq \max\{|\tilde{S}_{n_k} - W(t)|, |\tilde{S}_{n_{k+1}} - W(t)|\},$$

if $n_k \leq t \leq n_{k+1}$. Fix $h > 0$. By the strong law of large numbers as well as assumption (4.19), we can choose an integer q large enough in order that

$$A = \{\sup\{|\langle \sum_{i=1}^{n_k} T_i \rangle / n_k - 1|, k \geq q\} \leq h\},$$

satisfies

$$(4.23) \quad P\{A\} \geq 1 - h,$$

and further,

$$(4.24) \quad \sup\{[n_{k+1} - n_k]/n_k, k \geq q\} \leq h.$$

For $\omega \in A$, $k \geq q$ and $n_k \leq t \leq n_{k+1}$,

$$(4.25) \quad \left| \sum_{i=1}^{n_k} T_i - t \right| \leq 2hn_k.$$

Set

$$(4.26) \quad \forall \theta \in [0, 1], \quad \forall j \geq 1, \quad \tilde{\eta}_j(\theta) = \tilde{\eta}(\theta n_j) / \sqrt{n_j}.$$

If $\theta \geq n_q/n_j, j \geq q,$

$$\begin{aligned} \left| \tilde{\eta}_j(\theta) - \frac{W(\theta n_j)}{\sqrt{n_j}} \right| &\leq \max \left\{ \left| \frac{W(s) - W(\theta n_j)}{\sqrt{n_j}} \right|, |\theta n_j - s| \leq 2hn_j, 0 \leq \theta \leq 1 \right\}, \\ &\leq \max \left\{ \left| \frac{W(un_j) - W(\theta n_j)}{\sqrt{n_j}} \right|, 0 \leq \theta \leq 1, |u - \theta| \leq 2h \right\}. \end{aligned}$$

Let p be fixed and $n_j \in I_{k_p} \cap \mathfrak{B}$; one has on $A,$

$$(4.27) \quad \begin{aligned} \sup \{ |\tilde{\eta}_j(\theta) - \zeta_{n_j}(\theta)|, n_q/n_j \leq \theta \leq 1 \} \\ \leq \sup \{ |W(M^{k_p} \alpha \gamma) - W(M^{k_p} \theta \gamma)| / M^{k_p/2}, \\ 1 \leq \gamma \leq M, 0 \leq \theta \leq 1, |\theta - \gamma| \leq 2h \}, \end{aligned}$$

and on $\Omega,$

$$(4.28) \quad \begin{aligned} \sup \{ |\tilde{\eta}_j(\theta) - \zeta_{n_j}(\theta)|, 0 \leq \theta \leq n_q/n_j \} \\ \leq M^{-k_p/2} [\max \{ \tilde{S}_j, j \leq q + 1 \} + \max \{ |W(s)|, 0 \leq s \leq n_q \}], \end{aligned}$$

so that

$$(4.30) \quad P \{ \limsup_{j \rightarrow \infty} \{ |\tilde{\eta}_j(\theta) - \zeta_{n_j}(\theta)|, 0 \leq \theta \leq n_q/n_j \} = 0 \} = 1.$$

Set

$$(4.31) \quad \begin{cases} m = \text{median} \{ \sup \{ |W(uv) - W(\theta v)|, 1 \leq v \leq M, 0 \leq \theta \leq 1, \\ |\theta - u| \leq 2h \} \}, \\ \forall p \geq 1, \quad \varphi_p = \sqrt{2h \log(p + 2)} + 2m, \\ \forall p \geq 1, \quad A_p = \{ \sup \{ |\tilde{\eta}_j(\theta) - \zeta_{n_j}(\theta)|, n_j \in I_{k_p}, n_q/n_j \leq \theta \leq 1 \} > \varphi_p \}. \end{cases}$$

By using Borell's inequality again,

$$(4.32) \quad P\{A_p \cap A\} \leq 2\psi(\sqrt{2 \log(p + 2)} + m/\sqrt{2h}),$$

so that, $P\{A_p \cap A, p \text{ i.o.}\} = 0,$ and combining this with (4.23),

$$(4.33) \quad P\{ \limsup_{j \rightarrow \infty} \sup \{ |\tilde{\eta}_j(\theta) - \zeta_{n_j}(\theta)| / \varphi(n_j), n_q/n_j \leq \theta \leq 1 \} \leq \sqrt{2h} \} = 1.$$

From (4.30) and (4.33),

$$(4.34) \quad P\{ \limsup_{j \rightarrow \infty} \{ |\tilde{\eta}_j - \zeta_{n_j}| / \varphi(n_j) \leq \sqrt{2h} \} \geq 1 - h \}.$$

This achieves the proof by letting h tend to 0.

Q.E.D.

As an easy consequence, one has, by putting together Lemma 4.4 and Theorem 4.3

COROLLARY 4.5. *For any triple $(\mathfrak{s}, M, \varphi)$ be defined in accordance with (2.1), such that (4.19) holds, and for any sequence $X = \{X_i, i \geq 1\}$ of i.i.d. real r.v.'s satisfying $EX_1 = 0$ and $E(X_1)^2 = 1$, one has*

$$(4.35) \quad P\{\lim_{j \rightarrow \infty} d(\eta_j/\varphi(n_j), K) = 0\} = 1,$$

and

$$(4.36) \quad P\{C(\{\eta_j/\varphi(n_j), j \geq 1\}) = K\} = 1.$$

Proof of Theorem 2.9.

a) if $\lim_{k \rightarrow \infty} n_k/n_{k+1} = 1$. This is already proved by Corollary 4.5.

b) if $\lim_{k \rightarrow \infty} n_k/n_{k+1} = 0$. This is easily deduced from

$$P\left\{\limsup_{N \rightarrow \infty} \left(\frac{|S_{n_k}|}{\sqrt{n_N} \varphi(n_N)}, 1 \leq k \leq N-1\right) = 0\right\} = 1,$$

and the fact that the sequence of subdivisions $\{n_k/n_N, 1 \leq k \leq N\}$ tends to $\{0, 1\}$. Q.E.D.

§ 5. The law of the iterated logarithm on subsequences for random functions

As explained in Section 2, the main goal of this section is to prove that the behavior of partial sums of i.i.d. r.v.'s taking value in some infinite dimensional space B , can be as small as we want, like in euclidian spaces, when indexed on subsequences. This is the aim of the Theorem 2.10, that we are going to prove. We will use the following classical exponential bound for martingales.

LEMMA 5.1. *Let $M_n = \sum_{i=1}^n d_i$, ($n \geq 1$), be a real-valued martingale satisfying*

$$(5.1) \quad \forall i \geq 1, \quad |d_i| \leq \alpha_i \quad a.s.$$

Then,

$$(5.2) \quad \forall t \geq 0, \quad P\{|M_n| \geq t\} \leq 2 \exp\{-t^2/2[\sum_{i=1}^n \alpha_i^2]\}.$$

Proof of Theorem 2.10. We first assume that Y is a symmetric r.v. For the clarity we denote $(\Omega_Y, \mathcal{A}_Y, P_Y)$ the basic probability space of the sequence Y_1, Y_2, \dots . Let g_1, g_2, \dots be independent $\mathcal{N}(0, 1)$ r.v.'s defined on another probability space $(\Omega_g, \mathcal{A}_g, P_g)$. Let also $\varepsilon_1, \varepsilon_2, \dots$ be independent Rademacher r.v.'s defined on a third probability space $(\Omega_\varepsilon, \mathcal{A}_\varepsilon, P_\varepsilon)$. The corresponding symbols of integration are denoted E_Y, E_g and E_ε respectively. We now use an argument due to V.V. Yurinskii [19], which will be the first tool of the proof. Let $\underline{E}_i = \sigma\{\varepsilon_1, \dots, \varepsilon_i\}$ for every $i \geq 1$. Then

$$\|\sum_{k=1}^n Y_k \varepsilon_k\| - E_\varepsilon \|\sum_{k=1}^n Y_k \varepsilon_k\| = \sum_{k=1}^n d_i,$$

where

$$(5.3) \quad \forall i \geq 1, \quad d_i = E^{E_i} \{\|\sum_{k=1}^n Y_k \varepsilon_k\|\} - E^{E_{i-1}} \{\|\sum_{k=1}^n Y_k \varepsilon_k\|\},$$

and thus, $|d_i| \leq 2\|Y_i\|$, for every $i \geq 1$.

By applying Lemma 5.1 conditionally,

$$(5.4) \quad \forall t \geq 0, \quad P_\varepsilon \{ \|\sum_{i=1}^n Y_i \varepsilon_i\| - E_\varepsilon \|\sum_{i=1}^n Y_i \varepsilon_i\| > t \}, \\ \leq 2 \exp \left\{ -\frac{t^2}{8 \sum_{i=1}^n \|Y_i\|^2} \right\}.$$

We now need to control $E_\varepsilon \{\|\sum_{i=1}^n Y_i \varepsilon_i\|\}$. By using the easy fact that $\{g_i, i \geq 1\}$ and $\{g_i | \varepsilon_i, i \geq 1\}$ are identically distributed, one has by Jensen inequality,

$$(5.5) \quad E_\varepsilon \{\|\sum_{i=1}^n Y_i \varepsilon_i\|\} \leq \sqrt{\frac{\pi}{2}} E_g \{\|\sum_{i=1}^n Y_i g_i\|\}.$$

Further

$$(5.6) \quad \forall (s, t) \in T \otimes T, \quad [E_g |\sum_{k=1}^n g_k (Y_k(s) - Y_k(t))^{1/2}|^2], \\ \leq [\sum_{k=1}^n (Y_k(s) - Y_k(t))^2]^{1/2}, \\ \leq \rho(s, t) [\sum_{i=1}^n D_i^2]^{1/2},$$

where D_1, D_2, \dots is a sequence of independent copies of $\sup \{ |Y(s) - Y(t)| / \rho(s, t), (s, t) \in T \otimes T \}$. Since we have assumed that G is sample continuous, the classical integrability properties of gaussian processes (see e.g. [4]), together with Slepian's lemma (see again [4], Theorem 2.1.2) and results in [13], imply

$$(5.7) \quad \forall n \geq 1, \quad E_\varepsilon \{\|\sum_{i=1}^n Y_i \varepsilon_i\|\} \leq B [\sum_{i=1}^n D_i^2]^{1/2},$$

where the constant $B < \infty$ only depends on T and ρ and tends to 0 with $\text{diam}(T, \rho)$, Let $c > \max(1, ED_1^2)$ be fixed and

$$(5.8) \quad A = \sup \{ \sum_{i=1}^n D_i^2/n, n \geq q \} \leq c,$$

where we choose q large enough so that, by virtue of the strong law of large numbers.

$$(5.9) \quad P\{A\} \geq 1/2.$$

On A , one has, (using (5.7)),

$$(5.10) \quad \sup \{ [E_\varepsilon\{\|\sum_{i=1}^n Y_i \varepsilon_i\|\}]/\sqrt{n}, n \geq q \} \leq Bc.$$

Let now $(\mathfrak{s}, M, \varphi)$ be any triple defined according to (2.1). Choose $s > 0$ such that,

$$(5.11) \quad \forall p \geq 1, \quad s\sqrt{[2n_p^* \log(p+2)]/M} > Bc\sqrt{n_p^*} + 3\sqrt{n_p^* c \log(p+2)}.$$

Then, applying Levy's inequality conditionally to P_ε ,

$$\begin{aligned} P_Y \otimes P_\varepsilon \{ \{ \sup [\|\sum_{i=1}^n Y_i \varepsilon_i\|/\sqrt{2n \log(p+2)}, n \in \mathfrak{s} \cap I_{k_p}] > s \} \cap A \} \\ \leq 2E_Y \{ I_A \cdot P_\varepsilon \{ \|\sum_{i=1}^{n_p^*} Y_i \varepsilon_i\| > s\sqrt{[2n_p^* \log(p+2)]/M} \} \}, \end{aligned}$$

and by (5.4),

$$(5.12) \quad \begin{aligned} &\leq 4E_Y \left\{ I_A \cdot \exp \left(-\frac{9cn_p^* \log(p+2)}{8 \sum_{1 \leq i \leq n_p^*} D_i^2} \right) \right\}, \\ &\leq 4P_Y\{A\} \exp \left\{ -\frac{9}{8} \log(p+2) \right\}, \end{aligned}$$

so that, by using Borel-Cantelli lemma and 0-1 law,

$$(5.13) \quad P_Y \left\{ \sup \left\{ \frac{\|S_n(Y)\|}{\sqrt{n} \varphi(n)}, n \in \mathfrak{s} \right\} < \infty \right\} = 1,$$

then, by (5.7) and usual conclusion drawn from the inequality of J. Hoffman-Jørgensen [8], p. 164–165.

$$(5.14) \quad E_Y \left\{ \sup \left\{ \frac{\|S_n(Y)\|}{\sqrt{n} \varphi(n)}, n \in \mathfrak{s} \right\} \right\} < \infty.$$

We now can drop the assumption of symmetry, by using classical inequality of symmetrization. The conclusion is obtained by applying, as usual, the closed graph theorem and arguing as along the lines following (4.13) in [10]. Q.E.D.

§ 6. Conclusion

In this work, several problems are solved, and in the same time, some others are raised. One can summarize them as follows:

PROBLEM 1 (CONJECTURE): is any unbounded sequence of integers \mathfrak{s} a natural basis of the integers (i.e. $\mathfrak{s}^{(r)} = N$ for some finite r), if, and only if, $A(2^{\mathfrak{s}}) > 0?$, (assuming $1 \in \mathfrak{s}$).

PROBLEM 2: identify the set of cluster points $C(\{\gamma_N / [\sqrt{n_N} \varphi(n_N)], N \geq 1\})$ in full generality. A partial answer is brought by the Theorem 2.9.

PROBLEM 3: extend Theorem 2.1 in any Banach space. The recent characterization of this property when $\mathfrak{s} = N$ [10], reducing it to check the same property in probability, is certainly a good basis. Nevertheless the classical condition

$$E\{\|X\|^p / \log \log \|X\|\} < \infty,$$

which is needed to satisfy this property, is no longer necessary when indexing partial sums on subsequences. This brings a serious complication in order to truncate the r.v.'s, that is a necessary step in the proof given in [10], since the corresponding condition cannot be expressed in term of moment of X .

ACKNOWLEDGEMENT. This work was partially done during my stay, from October 1986 to March 1987, as visiting member, at the Courant Institute of Mathematical Sciences of New-York University, and I am indebted to Professor S.R.S. Varadhan for several useful and stimulating comments.

REFERENCES

- [1] Borell, C., The Brunn-Minkowski inequality in Gauss space, *Invent. Math.*, **30** (1975), 207–216.
- [2] Breiman, L., *Probability*. Addison Wesley, 1968.
- [3] Chung, K. L., Erdos, P. and Sirao, T., On the Lipschitz's conditions for Brownian motion, *J. Math. Soc. Japan*, **11** (1959), 263–274.
- [4] Fernique, X., Régularité des trajectoires des fonctions aléatoires gaussiennes, *Ecole d'été de Probabilités de Saint-Flour 1974*, *Lecture Notes in Math.*, **480** (1975), 1–96. Springer, Berlin.
- [5] Finkelstein, H., The law of the iterated logarithm for empirical distributions, *Ann. of Math. Stat.*, **42** (1971), 607–615.
- [6] Gut, A., *The law of the iterated on subsequences*, Publ. Uppsala University, 1983.

- [7] Hartman, P. and Wintner, A., On the law of the iterated logarithm, *Amer. J. Math.*, **63** (1941), 169–176.
- [8] Hoffman-Jørgensen, J., Sums of independent Banach space valued random variables, *Studia Math.*, **52** (1974), 159–186.
- [9] Kuelbs, J., The law of the iterated logarithm and related strong convergence theorems for Banach space valued random variables, *Ecole d'été de Probabilité de Saint-Flour 1975. Lecture Notes in Math.*, **539** (1976), 224–314. Springer, Berlin.
- [10] Ledoux, M. and Talagrand, M., Characterization of the law of the iterated logarithm in Banach spaces, *Ann. of Prob.*, **16**, (1988), 1242–1264.
- [11] Strassen, V., An invariance principle for the law of the iterated logarithm, *Z. Wahrsch. verw. Geb.*, **3** (1964), 211–226.
- [12] Strassen, V., A converse to the law of the iterated logarithm, *Z. Wahrsch. verw. Geb.*, **4** (1966), 265–268.
- [13] Talagrand, M., Regularity of gaussian processes, *Acta Math.*, **159** (1987), 99–149.
- [14] Torrang, I., The law of the iterated logarithm — cluster points of deterministic and random subsequences, *Prob. Math. Stat.*, **8** (1987), 133–141.
- [15] Vaughan, R. C., *The Hardy-Littlewood method*, Cambridge University Press, 1981.
- [16] Weber, M., La loi du logarithme itéré sur les sous-suites, *Comptes Rendus Acad. Sci. Paris*, **303** Sér. 1 (1986), 77–80.
- [17] Weber, M., La loi du logarithme itéré sur toute sous-suite, caractérisations, *C.R. Acad. Paris*, **305** (1987), 835–840.
- [18] Weber, M., The law of the iterated logarithm for subsequences in Banach spaces, *Prob. in Banach spaces VII*, (1988), to appear in Birkhäuser ed., *Progress in Prob. and Stat.*
- [19] Yurinskii, V. V., Exponential bound for large deviations, *Theor. Prob., Appl.*, **19** (1974), 154–155.

Université Louis Pasteur
Uer de Mathématiques et Informatique
7, rue René Descartes
67084 Strasbourg Cedex
France