

Medical News

EDITED BY GINA PUGLIESE

Enterovirus and Diabetes Mellitus

Because of the abrupt dependence on insulin among juvenile-onset diabetics, researchers have attempted to link juvenile diabetes with an infectious agent. Circumstantial evidence exists that the etiology might be an enterovirus, such as coxsackievirus or echovirus.

Researchers recently studied 128 young patients with recent onset of insulin-dependent diabetes who were admitted to the Children's Hospital in Pittsburgh, Pennsylvania. The sera from the 128 affected children were studied for IgM positivity to three enteroviruses and compared with the sera of a control group. For age groups 1 to 8 years, only enterovirus 41 serology was significantly different between the study and control cases. It was noted that the IgM positivity was unlikely to represent persistent infection because of the lack of clinical symptoms in affected children. The biggest differences were found in the 13- to 18 year age group, with IgM responses to six types of coxsackie and three types of echovirus significantly more often positive in cases than in controls.

FROM: Helfand RF Serologic evidence of an association between enterovirus and the onset of type I diabetes mellitus. *J Infect Dis* 1995;172:1206-1211.

Joint Commission to Streamline Surveys

Five national groups have agreed to work with the Joint Commission on Accreditation of Healthcare Organizations to streamline the hospital accreditation process. The goal of the initiative is to reduce the amount of survey duplication in accredited facilities.

Resolving the problem was high on the list of issues cited last year by the board of the American Hospital Association, which expressed a concern for the Joint Commission's overall performance in accrediting hospitals. The accreditors who will work with the Joint Commission to reduce survey duplication are the American Association for Ambulatory Healthcare (medical groups and ambulatory surgery centers); American Society of Histocompatibility and Immunogenetics (tissue-matching labs for organ transplants); Commission on Accreditation of Rehabilitation Facilities (rehabilitation and mental health facilities); Community Health Accreditation Program (home care programs); and the Commission on Office Laboratory Accreditation (laboratories).

By joining this effort, the accreditors' survey results are being accepted in lieu of the Joint Commission as part of the commission's accreditation of healthcare networks. Following a review and approval of the accreditors program, the Joint Commission will begin to accept other groups' survey results in lieu of its own for specific areas and programs of hospitals.

The Joint Commission is at different stages with each participant in comparing standards, survey decisions and processes, qualifications, training, and supervision of sur-

veyors. Accrediting groups still weighing a decision to participate in the collaborative effort include the American Osteopathic Association and the Unified Medical Quality Commission (capitated HMOs, medical groups, and independent practice associations). Only one group, the National Committee for Quality Assurance, which accredits HMOs, has declined to participate.

FROM: JCAHO joins project to streamline surveys. *Health Facilities Management*. 1995;8:7-8.

EPA Would Close Many Incinerators

The Environmental Protection Agency (EPA) proposed incinerator regulations that could have a devastating effect on how hospitals treat medical waste and dramatically increase their compliance cost, with little benefit to public health or the environment. Opponents to these regulations, to be issued in final form in April 1996, estimate that hospitals could spend up to \$1 billion to comply. The regulations would in effect signal the end of on-site medical waste incineration, causing all waste to be shipped to landfills.

Healthcare officials say the average cost to upgrade one hospital-based incinerator could run as high as \$1 million. The EPA also estimates that there are 3,700 existing incinerators, and approximately 700 new ones are expected to be built by 2000. The number of hospital-based incinerators is 2,200. The proposed regulations broadly define medical waste so that it would include virtually all waste generated in hospitals, healthcare, and research facilities. The rule also would require compliance testing every year for 3 years.

The recently installed state-of-the-art incinerator at the Mayo Clinic in Rochester, Minnesota, at a cost of \$14 million, would have to be shut down if the EPA implements its proposed rules. In addition, under the current EPA proposal, the one million pounds of noninfectious materials, such as office waste, that the clinic recycles each year would be treated as medical waste.

EPA officials have said they currently are reviewing some of the concerns raised by hospitals in preparation for the final regulations, which are due to be released in early 1996.

FROM: Incinerator plan provokes complaints. *Health Facilities Management*. 1995;8(10):11-12.

Infection Control Indicators in 1996

The Joint Commission has announced that the infection control indicators approved by the Joint Commission's Board of Commissioners in 1995 will become available to Indicator Measurement System (IMSystem) participants in 1996. The three infection control indicators focus on surgical site infections, pneumonia in ventilator patients, and bloodstream infections in patients with central or umbilical