

A SCHWARZ LEMMA FOR COMPLETE RIEMANNIAN MANIFOLDS

LEUNG-FU CHEUNG AND PUI-FAI LEUNG

We prove a Schwarz Lemma for conformal mappings between two complete Riemannian manifolds when the domain manifold has Ricci curvature bounded below in terms of its distance function. This gives a partial result to a conjecture of Chua.

1. INTRODUCTION

In recent paper [2], Chua has given an interesting generalisation of the classical Ahlfors-Schwarz Lemma to conformal mappings from the hyperbolic space into a Riemannian manifold. His result is the following:

THEOREM. [2] *Let (M^n, g) be an n -dimensional Riemannian manifold and (H^n, g_0) be the n -dimensional hyperbolic space of constant sectional curvature -1 . Let $f: (H^n, g_0) \rightarrow (M^n, g)$ be a conformal map. If $\text{scal}(M^n) \leq -n(n-1)$ where $\text{scal}(M^n)$ denotes the scalar curvature of M^n , then f is distance decreasing, that is, $f^*g \leq g_0$.*

In the same paper [2], Chua proposed the following interesting generalisation.

CONJECTURE. [2] *Let (M_0^n, g_0) and (M^n, g) be two complete n -dimensional Riemannian manifolds. Let $f: (M_0^n, g_0) \rightarrow (M^n, g)$ be a conformal map. If there exist positive constants c and k such that $f^*(\text{scal}(M^n)) \leq k^2 \text{scal}(M_0^n) \leq -c^2 < 0$, then $k^2(f^*g) \leq g_0$.*

Our purpose in this note is to prove the following partial result on Chua's Conjecture.

MAIN THEOREM. *Let (M_0^n, g_0) and (M^n, g) be two complete n -dimensional Riemannian manifolds. Let $f: (M_0^n, g_0) \rightarrow (M^n, g)$ be a conformal map. Assume that $\text{Ric}(M_0^n) \geq -a^2(1 + y^2 \log^2(y + 2))$, where $\text{Ric}(M_0^n)$ denotes the Ricci curvature of M_0^n , a denotes a constant and y is the distance function from a fixed origin $x_0 \in M_0^n$. If there exist positive constants c and k such that $f^*(\text{scal}(M^n)) \leq k^2 \text{scal}(M_0^n) \leq -c^2 < 0$, then $k^2(f^*g) \leq g_0$.*

Received 30 July 1996

We would like to thank K.S. Chua for introducing us to his conjecture and for many helpful discussions.

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9729/97 \$A2.00+0.00.

2. PROOF OF THE MAIN THEOREM

Since f is conformal, there exists a smooth function ρ on M_0^n such that

$$f^*g = e^{2\rho}g_0.$$

(See [3, p.53].) Following [3, p.62] we put $u = e^{-\rho}$.

Since $u \geq 0$, therefore $r := \inf u$ exists.

The assumption on $\text{Ric}(M_0^n)$ allows us to apply the Generalised Maximum Principle of Chen and Xin [1, Theorem 2.2, p.359] to the function $-u$ to obtain the following:

For each positive integer m , there exists a point $x_m \in M_0^n$ such that

$$0 \leq r \leq u(x_m) < r + \frac{1}{m},$$

$$|\nabla u(x_m)|^2 < \frac{1}{m},$$

and

$$\Delta u(x_m) > -\frac{1}{m}.$$

Here ∇u and Δu denote the gradient and the Laplacian of u respectively.

Let $K_0 = \text{scal}(M_0^n)$ and $K = f^*(\text{scal}(M^n))$. Then by [3, p.62, (3.12)] we have

$$K = u^2K_0 + 2(n-1)u\Delta u - n(n-1)|\nabla u|^2.$$

Therefore

$$k^2K_0 \geq u^2K_0 + 2(n-1)u\Delta u - n(n-1)|\nabla u|^2.$$

(since $K \leq k^2K_0$) and hence

$$\begin{aligned} (k^2 - u(x_m)^2)K_0(x_m) &\geq (n-1) \left[2u(x_m)\Delta u(x_m) - n|\nabla u(x_m)|^2 \right] \\ (2.1) \qquad \qquad \qquad &\geq (n-1) \left[2 \left(r + \frac{1}{m} \right) \left(-\frac{1}{m} \right) - \frac{n}{m} \right] \\ &= -\frac{n-1}{m} \left(2r + \frac{2}{m} + n \right). \end{aligned}$$

Our Main Theorem will be proven if we can show that $r \geq k$. Suppose on the contrary that we have $r < k$. Then for m large enough, we have

$$u(x_m) < r + \frac{1}{m} < k.$$

Since $K \leq k^2K_0 \leq -c^2$, it follows from (2.1) that for m large enough, we have

$$(k^2 - u(x_m)^2)(-c^2) \geq -\frac{n-1}{m} \left(2r + \frac{2}{m} + n \right) k^2.$$

Letting $m \rightarrow \infty$, we obtain

$$(k^2 - r^2)(-c^2) \geq 0$$

and this contradicts the assumption that $r < k$. Therefore we must have $r \geq k$. □

REFERENCES

- [1] Q. Chen and Y.L. Xin, 'A generalized maximum principle and its applications in geometry', *Amer. J. Math.* **114** (1992), 355–366.
- [2] K.S. Chua, 'A generalisation of Ahlfors-Schwarz lemma to Riemannian geometry', *Bull. Austral. Math. Soc.* **51** (1995), 517–520.
- [3] K. Yano and M. Obata, 'Conformal changes of Riemannian metrics', *J. Differential Geom.* **4** (1970), 53–72.

Department of Mathematics
Hong Kong Polytechnic University
Hung Hom
Kowloon
Hong Kong
e-mail: malfcheu@hkpu02.polyu.edu.hk

Department of Mathematics
National University of Singapore
Kent Ridge
Singapore 0511
e-mail: matfredl@leonis.nus.sg