

rate of the algorithm of 39% (12 of 31) is somewhat misleading. This rate includes both patients for whom isolation was delayed for >24 hours (n=7) or not implemented (n=5) and does not differentiate by level of infectiousness. Of the 12 patients with pulmonary TB who were not placed in negative-pressure isolation rooms within 24 hours of admission to the MGH, 5 were AFB smear-positive, 6 were smear-negative, and 1 patient had no smear obtained (TB diagnosed on autopsy). If only high-risk patients (ie, smear-positive, more infectious) are prioritized for immediate isolation, then only 5 of 31 patients with pulmonary TB were not isolated appropriately (algorithm failure rate=16%). The timing and duration of isolation of smear-negative patients is more uncertain, given the knowledge that only a small minority of such patients will be found to be culture-positive. The use of more rapid and sensitive diagnostic tests that currently are undergoing evaluation at our hospital and others may assist in the assessment of these patients.

Finally, we agree that clinical algorithms are subject to limitations and cannot substitute for careful clinical judgment. However, use of the TB algorithm at the MGH has improved the awareness of TB among clinicians and other health-care workers and has assisted infection control personnel in the ongoing evaluation of TB control program needs and priorities throughout the hospital. We agree that new, as well as feasible, approaches to the management of this problem are needed.

**Clare F. Pegues, RN, MPH, CIC**  
**David A. Pegues, MD**  
**Cyrus C. Hopkins, MD**  
 Massachusetts General Hospital  
 Boston, Massachusetts

#### REFERENCES

1. Wurtz R. Administrative controls for TB: 'keep doing what you've always done, and you'll get what you always got.' *Infect Control Hosp Epidemiol* 1996;17:409-411.
2. Pegues CF, Johnson DC, Pegues DA, Spencer M, Hopkins CC. Implementation and evaluation of an algorithm for isolation of patients with suspected pulmonary tuber-

culosis. *Infect Control Hosp Epidemiol* 1996;17:412-418.

### Triple Combination Antiretroviral Prophylaxis for Needlestick Exposure to HIV

#### To the Editor:

Zidovudine has been used widely for prophylactic treatment of persons exposed to human immunodeficiency virus (HIV) through needlestick exposures. There is limited evidence of efficacy due to the relatively small risk of infection after needlestick and the difficulty of implementing randomized, placebo-controlled trials.<sup>1</sup> There are, however, clearly documented failures, despite high doses of zidovudine given soon after HIV exposure.<sup>2</sup> There is also increasing frequency of zidovudine resistance in persons with HIV who have been taking zidovudine. Recent work has demonstrated that triple antiretroviral therapy with protease inhibitors is extremely effective at decreasing viral load among patients with established HIV infection.<sup>3</sup> Although the Centers for Disease Control and Prevention (CDC) recently recommended triple combination antiretroviral prophylaxis for needlestick exposure to HIV, there is no data on the tolerability or effectiveness of this therapy.<sup>4</sup>

A healthy, 38-year-old health-care worker was performing femoral vein phlebotomy on a patient with acquired immunodeficiency syndrome. The healthcare worker accidentally sustained a deep intramuscular index-finger needlestick with an 18-gauge needle that had just come out of the femoral vein with obvious blood on it. The wound was bled, and triple therapy with D4T, 3TC, and indinavir was begun within 2 hours of the needlestick. Triple therapy was continued for 2 weeks without any side effects. Human immunodeficiency virus serology at baseline and at 3- and 6-month follow-up was negative. The healthcare worker remained in excellent health.

This is the first reported case of triple combination antiretroviral therapy, including a protease inhibitor, to prevent HIV infection after a signifi-

cant exposure. The absence of HIV infection after this needlestick exposure is not surprising, given the low likelihood of developing subsequent HIV infection. As noted in the recent CDC recommendations, triple combination therapy is likely to be more effective than zidovudine; however, it is certainly more expensive, and there is potential for increased toxicity. Zidovudine was not used in this case because the index patient had been on zidovudine for many years. Consideration of a patient's prior antiretroviral treatment may be useful in guiding appropriate prophylactic strategies in the event of occupational exposure.<sup>5</sup> We are encouraged that a registry of prophylactic treatment has been established (telephone: 1-888-737-4448). It will be important to describe the regimen taken and its tolerability, the extent of needlestick injury, and the rate of seroconversion. These data should be interpreted in the context of index patient viral load and previous antiretroviral therapy.

**Josiah D. Rich, MD, MPH**  
**Bharat Ramratnam, MD**  
**Timothy P. Flanigan, MD**

The Miriam Hospital  
 Brown University School of Medicine  
 Providence, Rhode Island

#### REFERENCES

1. Centers for Disease Control and Prevention. Case control study of HIV seroconversion in health care workers after percutaneous exposure to HIV infected blood—France, United Kingdom and United States, January 1988-August 1994. *MMWR* 1995;44:929-933.
2. Tokars JI, Marcus R, Culver DH, et al. Surveillance of HIV infection and zidovudine use among health care workers after occupational exposure to HIV infected blood. The CDC Cooperative Needlestick Surveillance Group. *Ann Intern Med* 1993;118:913-919.
3. Collier AC, Coombs R, Schoenfeld DA, et al. Treatment for human immunodeficiency virus infection with saquinavir, zidovudine and zalcitabine. *N Engl J Med* 1996;334:1011-1017.
4. Centers for Disease Control and Prevention. Update: provisional Public Health Service recommendations for chemoprophylaxis after occupational exposure to HIV. *MMWR* 1996;45:468-472.
5. Gerberding J. Prophylaxis for occupational exposure to HIV. *Ann Intern Med* 1996; 125:497-501.