

## A NOTE ON HOLOMORPHIC VECTOR BUNDLES OVER COMPLEX TORI

HISASI MORIKAWA

1. Let  $\omega : \mathbf{Z}^{2r} \rightarrow \mathbf{C}^r$  be an isomorphism of the free additive group of rank  $2r$  into the complex vector  $n$ -space such that the quotient group  $\mathbf{T} = \mathbf{C}^r / \omega(\mathbf{Z}^{2r})$  is compact, i.e.,  $\mathbf{T}_\omega$  is a complex torus.

We mean by a matrix multiplier of rank  $n$  with respect to  $\omega$  a family of complex holomorphic  $n \times n$ -matrix functions  $\{\mu_\alpha(z)\}_{\alpha \in \mathbf{Z}^{2r}}$  on  $\mathbf{C}^r$  such that

- 1)  $\det \mu_\alpha(z) \neq 0 \quad (z \in \mathbf{C}^r),$
- 2)  $\mu_\alpha(z) \mu_\beta(z + \omega(\alpha)) = \mu_{\alpha+\beta}(z), \quad (\alpha, \beta \in \mathbf{Z}^{2r}).$

By virtue of the conditions 1) and 2) we may define an action of  $\mathbf{Z}^{2r}$  on the product  $\mathbf{C}^r \times \mathbf{C}^n$  as follows:

$$(z, u) \rightarrow (z + \omega(\alpha), v\mu_\alpha(z)), \quad (\alpha \in \mathbf{Z}^{2r}).$$

The quotient  $\mathbf{V}_\mu$  of  $\mathbf{C}^r \times \mathbf{C}^n$  by this action of  $\mathbf{Z}^{2r}$  is a holomorphic vector  $n$ -bundle over the complex torus  $\mathbf{T}_\omega$ , and conversely every holomorphic vector bundle over  $\mathbf{T}_\omega$  is constructed by this method with a matrix multiplier, since holomorphic vector bundles over a vector space are always trivial.<sup>1)</sup>

2. We shall recall the definition of *finite Heisenberg groups* and their canonical representations.

Let  $G$  be an additive group of order  $n$  and of exponent  $d$ , and  $\zeta$  be a primitive  $d$ -th root of unity. Let  $\hat{G}$  be the dual group of  $G$  defined by a pairing  $(\hat{a}, a) \rightarrow \langle \hat{a}, a \rangle$  of  $\hat{G} \times G$  into the multiplicative group  $\{1, \zeta, \dots, \zeta^{d-1}\}$ . We mean by the finite Heisenberg group  $H(G)$  associated with  $G$  the group consisting of triples  $\{(\hat{a}, a, \zeta^l) \mid \hat{a} \in \hat{G}, a \in G, 0 \leq l \leq d-1\}$  with the composition law

$$(\hat{a}, a, \zeta^l)(\hat{b}, b, \zeta^h) = (\hat{a} + \hat{b}, a + b, \langle \hat{a}, b \rangle \zeta^{l+h}).$$

---

Received March 17, 1970.

The Heisenberg group  $H(G)$  has a faithful irreducible  $n \times n$ -matrix representation  $\{U_{(\hat{a}, a, \zeta^t)}\}$  characterized by its character

$$\text{tr}U_{(\hat{a}, a, \zeta^t)} = \begin{cases} 0 & \text{for } \hat{a} + a \neq 0, \\ n\zeta^t, & \text{for } \hat{a} + a = 0, \end{cases}$$

We call such a representation  $U$  the canonical representation of  $H(G)$ . Actually  $U$  is given by

$$U_{(\hat{a}, a, \zeta^t)} = (\zeta^t u_{b,c}(\hat{a} + a))_{b,c \in G},$$

where

$$u_{b,c}(\hat{a} + a) = \langle \hat{a}, b \rangle \delta_{b,c+a},$$

$$\delta_{b,a} = \begin{cases} 1 & \text{for } b = a \\ 0 & \text{otherwise,} \end{cases}$$

In these terminologies we shall show the following result:

**THEOREM 1.** *Let  $\omega : \mathbf{Z}^{2r} \rightarrow \mathbf{C}^r$  be an isomorphism such that  $\mathbf{T}_\omega = \mathbf{C}^r / \omega(\mathbf{Z}^{2r})$  is a complex torus. If a matrix multiplier  $\{\mu_\alpha(z)\}$  of rank  $n$  with respect to  $\omega$  satisfies the conditions*

- i)  $\mu_\alpha(z) = \mu_\alpha(0)\chi_\alpha(z)$  with scalar functions  $\chi_\alpha(z)$  ( $\alpha \in \mathbf{Z}^{2r}$ ),
- ii) the commutators of  $\{\mu_\alpha(0)\}_{\alpha \in \mathbf{Z}^{2r}}$  are scalar matrices,

then there exist an additive group  $G$  of order  $n$ , a surjective homomorphism  $\hat{\sigma} \oplus \sigma$  of  $\mathbf{Z}^{2r}$  onto  $\hat{G} \oplus G$  and a family of holomorphic functions  $\{\xi_\alpha(z)\}_{\alpha \in \mathbf{Z}^{2r}}$  such that

$$\mu_\alpha(z) = U_{(\hat{\sigma}(\alpha), \sigma(\alpha), 1)} \xi_\alpha(z), \quad (\alpha \in \mathbf{Z}^{2r}),$$

where  $U$  is the canonical representation of the Heisenberg group  $H(G)$ .

*Proof.* Putting  $z = 0$  in  $\mu_\alpha(z)\mu_\beta(z + \omega(\alpha)) = \mu_{\alpha+\beta}(z)$ , we have

$$\begin{aligned} \mu_\alpha(0)\mu_\beta(0)\chi_\beta(\omega(\alpha)) &= \mu_{\alpha+\beta}(0), \\ \chi_\beta(\omega(\alpha))^n &= \frac{\det \mu_{\alpha+\beta}(0)}{\det \mu_\alpha(0) \det \mu_\beta(0)}, \quad (\alpha, \beta \in \mathbf{Z}^{2r}). \end{aligned}$$

Since the left hand side of the last equation is symmetric with respect to  $\alpha$  and  $\beta$ , denoting

$$\zeta_{\alpha,\beta} = \chi_\beta(\omega(\alpha))^{-1} \chi_\alpha(\omega(\beta)),$$

we have

$$\zeta_{\alpha,\beta}^n = 1, \quad \zeta_{\alpha,\beta}\zeta_{\beta,\alpha} = 1,$$

and the commutation relation

$$\mu_\alpha(0)\mu_\beta(0) = \zeta_{\alpha,\beta}\mu_\beta(0)\mu_\alpha(0).$$

Moreover  $\zeta_{\alpha,\beta}$  is bimultiplicative, namely

$$\zeta_{\alpha,\beta+\gamma} = \zeta_{\alpha,\beta}\zeta_{\alpha,\gamma}.$$

because

$$\begin{aligned} \mu_\alpha(0)\mu_\beta(0)\mu_\gamma(0) &= \mu_\alpha(0)\mu_{\beta+\gamma}(0)\chi_\gamma(\omega(\beta))^{-1} \\ &= \zeta_{\alpha,\beta+\gamma}\mu_{\beta+\gamma}(0)\mu_\alpha(0)\chi_\gamma(\omega(\beta))^{-1} \\ &= \zeta_{\alpha,\beta+\gamma}\mu_\beta(0)\mu_\gamma(0)\mu_\alpha(0) \\ &= \zeta_{\alpha,\beta+\gamma}\zeta_{\beta,\alpha}\zeta_{\gamma,\alpha}\mu_\alpha(0)\mu_\beta(0)\mu_\gamma(0) \\ &= \zeta_{\alpha,\beta+\gamma}\zeta_{\alpha,\beta}^{-1}\zeta_{\alpha,\gamma}^{-1}\mu_\alpha(0)\mu_\beta(0)\mu_\gamma(0). \end{aligned}$$

Therefore  $\mu_\alpha(0)^n\mu_\beta(0) = \zeta_{\alpha,\beta}^n\mu_\beta(0)\mu_\alpha(0)^n = \mu_\beta(0)\mu_\alpha(0)^n$ , ( $\alpha, \beta \in \mathbf{Z}^{2r}$ ) and these  $\mu_\alpha(0)^n(\alpha \in \mathbf{Z}^{2r})$  are scalar matrices by the condition (ii). On the other hand

$$\mu_{n\alpha}(0) = \chi_\alpha(\omega(\alpha))\chi_\alpha(\omega(2\alpha)) \cdots \chi_\alpha(\omega(n-1)\alpha)\mu_\alpha(0)^n,$$

hence  $\mu_{n\alpha}(0)$ , ( $\alpha \in \mathbf{Z}^{2r}$ ), are scalar matrices. Let  $N$  be the subgroup consisting of all the elements  $\alpha$  such that  $\mu_\alpha(0)$  is a scalar matrix. Then  $N \supset n\mathbf{Z}^{2r}$  and the element  $\alpha$  is characterized by  $\zeta_{\alpha,\beta} = 1$  for every  $\beta$  in  $\mathbf{Z}^{2r}$ ,

Since  $\zeta_{\alpha,\beta}$  is bimultiplicative and  $\zeta_{\alpha,\beta} = \zeta_{\beta,\alpha}^{-1}$ , there exists a skew symmetric rational matrix  $A$  such that

$$\zeta_{\alpha,\beta} = e^{2\pi\sqrt{-1}\alpha A\beta}$$

and  $nA$  is an integral matrix. Therefore we can choose a base  $\{a_1, \dots, a_s, \hat{a}_1, \dots, \hat{a}_s\}$  of the quotient additive group  $\mathbf{Z}^{2r}/N$  and their representatives  $\alpha_1, \dots, \alpha_s, \hat{\alpha}_1, \dots, \hat{\alpha}_r$  in  $\mathbf{Z}^{2r}$  such that

$$\begin{aligned} \zeta_{\alpha_i, \alpha_j} &= \zeta_{\hat{\alpha}_i, \hat{\alpha}_j} = 1, & (1 \leq i, j \leq s) \\ \zeta_{\hat{\alpha}_i, \alpha_h} &= 1, & (i \neq h) \\ \zeta_{\hat{\alpha}_i, \alpha_i}^{d_i} &= 1, & (1 \leq i \leq s), \end{aligned}$$

where  $d_i$  is the common order of  $\alpha_i$  and  $\hat{\alpha}_i$ . We denote by  $G$  and  $\hat{G}$  the subgroups of  $\mathbf{Z}^{2r}/N$  generated by  $\{a_1, \dots, a_s\}$  and  $\{\hat{a}_1, \dots, \hat{a}_s\}$ , respectively. Then  $\mathbf{Z}^{2r}/N = G \oplus \hat{G}$  and the map

$$\left(\sum_{i=1}^s l_i \hat{a}_i, \sum_{i=1}^s h_i a_i\right) \rightarrow \zeta^{\sum_{i=1}^s l_i \hat{a}_i, \sum_{i=1}^s h_i a_i} = \prod_{i=1}^s \zeta^{l_i h_i}$$

is the pairing of the dual pair  $(G, \hat{G})$ . We denote by  $\langle \sum_{i=1}^s l_i \hat{a}_i, \sum_{i=1}^s h_i a_i \rangle$  its value. Denote by  $G^*$  and  $\hat{G}^*$  the inverse images of  $G$  and  $\hat{G}$  in  $\mathbb{Z}^{2r}$ , respectively. Then from the definition of  $G$  and  $\hat{G}$  it follows

$$\begin{aligned} \mu_\alpha(0)\mu_\beta(0) &= \mu_\beta(0)\mu_\alpha(0), & (\alpha, \beta \in G^*) \\ \mu_{\hat{\alpha}}(0)\mu_{\hat{\beta}}(0) &= \mu_{\hat{\beta}}(0)\mu_{\hat{\alpha}}(0), & (\hat{\alpha}, \hat{\beta} \in \hat{G}^*), \end{aligned}$$

Since  $d_i \hat{a}_i$  and  $d_i a_i$  are elements in  $N$ , the matrices  $\mu_{d_i \hat{a}_i}(0)$  and  $\mu_{d_i a_i}(0)$  are scalar matrices and we can choose scalar matrices  $\nu_{\hat{a}_i}$  and  $\nu_{a_i}$  such that

$$\nu_{\hat{a}_i}^{d_i} = \mu_{d_i \hat{a}_i}(0) \quad \nu_{a_i}^{d_i} = \mu_{d_i a_i}(0) \quad (1 \leq i \leq s).$$

Let us construct an irreducible  $n \times n$ -representation of the Heisenberg group  $H(G)$ . From the definition follows the commutation relations

$$\begin{aligned} (\nu_{\hat{a}_i}^{-1} \mu_{\hat{a}_i}(0)) (\nu_{\hat{a}_j}^{-1} \mu_{\hat{a}_j}(0)) &= (\nu_{\hat{a}_j}^{-1} \mu_{\hat{a}_j}(0)) (\nu_{\hat{a}_i}^{-1} \mu_{\hat{a}_i}(0)), \\ (\nu_{\hat{a}_i}^{-1} \mu_{a_i}(0)) (\nu_{a_j}^{-1} \mu_{a_j}(0)) &= (\nu_{a_j}^{-1} \mu_{a_j}(0)) (\nu_{\hat{a}_i}^{-1} \mu_{a_i}(0)), \\ (\nu_{\hat{a}_l}^{-1} \mu_{\hat{a}_l}(0)) (\nu_{\hat{a}_h}^{-1} \mu_{a_h}(0)) &= (\nu_{\hat{a}_h}^{-1} \mu_{a_h}(0)) (\nu_{\hat{a}_l}^{-1} \mu_{\hat{a}_l}(0)), \quad (l \neq h), \\ (\nu_{\hat{a}_i}^{-1} \mu_{\hat{a}_i}(0)) (\nu_{a_i}^{-1} \mu_{a_i}(0)) &= \langle \hat{a}_i, a_i \rangle (\nu_{\hat{a}_i}^{-1} \mu_{a_i}(0)) (\nu_{\hat{a}_i}^{-1} \mu_{\hat{a}_i}(0)). \end{aligned}$$

This shows that we may define a map

$$\sum_{i=1}^s l_i a_i + \sum_{i=1}^s h_i \hat{a}_i \rightarrow M\left(\sum_{i=1}^s l_i a_i + \sum_{i=1}^s h_i \hat{a}_i\right) = \prod_{i=1}^s (\nu_{a_i}^{-1} \mu_{a_i}(0))^{l_i} \prod_{i=1}^s (\nu_{\hat{a}_i}^{-1} \mu_{\hat{a}_i}(0))^{h_i}$$

such that

$$\begin{aligned} M\left(\sum_{i=1}^s l_i a_i\right) M\left(\sum_{i=1}^s h_i \hat{a}_i\right) &= M\left(\sum_{i=1}^s l_i a_i + \sum_{i=1}^s h_i \hat{a}_i\right), \\ M\left(\sum_{i=1}^s h_i \hat{a}_i\right) M\left(\sum_{i=1}^s l_i a_i\right) &= \left\langle \sum_{i=1}^s h_i \hat{a}_i, \sum_{i=1}^s l_i a_i \right\rangle M\left(\sum_{i=1}^s l_i a_i + \sum_{i=1}^s h_i \hat{a}_i\right). \end{aligned}$$

Let  $d$  be the exponent of  $G$  and  $\zeta$  be the primitive  $d$ -th root of unity. Then the map

$$(\hat{a}, a, \zeta^l) \rightarrow U_{(a, \hat{a}, \zeta^l)} = \zeta^l M(\hat{a} + a)$$

is a representation of the Heisenberg group of  $H(G)$ .

From the commutation relation

$$U_{(\hat{a}, a, \zeta^l)} U_{(\hat{b}, b, \zeta^h)} = \langle \hat{a}, b \rangle U_{(\hat{b}, b, \zeta^h)} U_{(\hat{a}, a, \zeta^l)}$$

follows

$$\begin{aligned} \text{tr } U_{(\hat{a}, a, \zeta^l)} &= \langle \hat{a}, b \rangle \text{tr } U_{(\hat{a}, a, \zeta^l)}, \\ \text{tr } U_{(\hat{b}, b, \zeta^h)} &= \langle \hat{a}, b \rangle \text{tr } U_{(\hat{b}, b, \zeta^h)}. \end{aligned}$$

This mean that

$$\text{tr } U_{(\hat{a}, a, \zeta^l)} = \begin{cases} 0 & \text{for } a + \hat{a} \neq 0 \\ n\zeta^l & \text{for } a + \hat{a} = 0, \end{cases}$$

Since the commutators of  $\{U_{(\hat{a}, a, \zeta^l)}\}$  are scalar matrices,  $U$  is the canonical representation of  $H(G)$ . Denote by  $\sigma(\alpha) + \hat{\sigma}(\alpha)$  the direct sum decomposition of the image of  $\alpha$  in  $\mathbf{Z}^{2r}/N$  with respect to the decomposition  $\mathbf{Z}^{2r}/N = G \oplus \hat{G}$  and put

$$\begin{aligned} \rho(\alpha) &= (\hat{\sigma}(\alpha), \sigma(\alpha), 1), \\ \xi_\alpha(z) &= U_{\rho(\alpha)}^{-1} \mu_\alpha(z), \quad (\alpha \in \mathbf{Z}^{2r}). \end{aligned}$$

Then  $\xi_\alpha(z) (\alpha \in \mathbf{Z}^{2r})$  are scalar functions satisfying

$$\mu_\alpha(z) = U_{\rho(\alpha)} \xi_\alpha(z), \quad (\alpha \in \mathbf{Z}^{2r}).$$

This completes the proof of Theorem 1.

3. In the notations in the proof of Theorem 1, we denote by  $\lambda$  the natural isogeny of the complex tori.

$$\lambda : \mathbf{C}^r / \omega(\hat{G}^*) \rightarrow \mathbf{T}_\omega = \mathbf{C}^r / \omega(\mathbf{Z}^{2r}).$$

After changing the base, we may assume that

$$U_{(a, \hat{a}, 1)} = \langle \hat{a}, b \rangle \delta_{b, c+a} \quad (a \in G, \hat{a} \in \hat{G})$$

We shall define line bundles  $L_{\eta(a)}$  ( $a \in G$ ) over  $\mathbf{C}^r / \omega(\hat{G}^*)$  as follows. Let  $\{\eta_\beta^{(a)}(z)\}_{\beta \in \hat{G}^*}$  ( $a \in G$ ) be families of functions defined by

$$\eta_\beta^{(a)}(z) = \langle \hat{a}(\beta), a \rangle \xi_\beta(z) \quad (\beta \in \hat{G}^*; a \in G).$$

Then it follows

$$\eta_\alpha^{(a)}(z) \eta_\beta^{(a)}(z + \omega(\alpha)) = \eta_{\alpha+\beta}^{(a)}(z) \quad (\alpha, \beta \in \hat{G}^*; a \in G),$$

and thus there exist line bundles  $L_{\eta(a)}$  ( $a \in G$ ) over  $\mathbf{C}^r / \omega(\hat{G}^*)$  associated with

multipliers  $\{\eta_{\beta}^{(\omega)}(z)\}_{\beta \in \hat{G}^*}$  ( $a \in G$ ), respectively. The direct sum

$$\bigoplus_{a \in G} L_{\eta(a)}$$

may be regarded as the pull back by  $\lambda$  of the vector bundle on  $T_{\omega} = \mathbf{C}^r / \omega(\mathbf{Z}^{2r})$  associated with the multifier  $\{\mu_{\alpha}(z)\}_{\alpha \in \mathbf{Z}^{2r}}$  such that

$$\mu_{\alpha}(z) = U_{\rho(\alpha)} \xi_{\alpha}(z), \quad (\alpha \in \mathbf{Z}^{2r}),$$

This means that the vector bundle  $V_{\mu}$  is the direct image  $\lambda_*(L_{\eta(a)})$  of any one of the line bundles  $L_{\eta(a)}$  with respect to the isogeny  $\lambda$ .

Then next is the geometric expression of Theorem 1.

**THEOREM 2.** *Let  $V$  be a holomorphic vector  $n$ -bundle over a complex torus  $T$  such that*

i)  $End_{\mathbf{C}}(V) = \mathbf{C}$ ,

ii) *the projective  $(n-1)$ -bundle  $P(V)$  associated with  $V$  has a family of constant transition functions. Then there exist an isogeny  $\lambda: S \rightarrow T_{\omega}$  of degree  $n$  and a line bundle  $L$  over the complex torus  $S$  such that  $V$  is isomorphic to the direct image  $\lambda_*(L)$  of  $L^{2)}$  with respect to  $\lambda$ .*

#### REFERENCES

- [ 1 ] Gunning, Rossi; Analytic Functions of Several Complex Variables, Prentice-Hall, 1965.  
 [ 2 ] T. Oda; Vector bundles on an elliptic curve. (to appear in Nagoya Math. J.).

*Mathematical Institute of Nagoya University*

<sup>1)</sup> See [1].

<sup>2)</sup> T. Oda dealt with such direct images  $\lambda_*(L)$  systematically in his paper [2]; our result is nothing else than a characterization of simple direct images  $\lambda_*(L)$ .