A NEW journal in the applications of categorical, algebraic and geometric methods in Computer Science

Mathematical Structures in Computer Science

Why a new journal?

Mathematical Structures in Computer Science (MSCS) is a new journal of theoretical computer science which focuses on the application of ideas from the structural side of mathematics and mathematical logic to computer science. The journal aims to bridge the gap between theoretical contributions and software design, publishing original papers or broad surveys with original perspectives in all areas of computing, provided that ideas or results from algebra, geometry or category theory form a basis for the work. MSCS is distinct from existing titles in that it specialises in the art of applying mathematics of genuine interest and general applicability to computer science; its objective is to promote the useful application of high level mathematics to language design and software implementation. The journal will increase the circulation of new results in this fast growing area.

Editor-in-Chief

G Longo Ecole Normale Supérieure, 45 rue d'Ulm, 75230 Paris (email: longo@earn.frulm63)

Associate Editors

P L Curien Laboratoire d'Informatique, Ecole Normale Supérieure, Paris

A M Pitts

Computer Laboratory, University of Cambridge

Forthcoming articles

From Petri Nets to Linear Logic Narcisio Marti-Oliet & Jose Meseguer Impredicative Calculi of Dependent Types

THOMAS STREICHER

Operations on Records Luca Cardelli & John C Mitchell

Predicate Transformer Semantics ERNIE MANES

Declarative Continuations: An Investigation of Duality in Programming ANDRZEJ FILINSKI

A Category-theoretic Account of Program Modules EUGENIO MOGGI

What is Functional Programming? R F C WALTERS

Subscription information

Mathematīcal Structures in Computer Science, Vol 1: March, July and November 1991: £75/\$135. Delivery by airmail £9.50 per year extra (dollar price includes airfreight to New York). ISSN 0960-1295

Send your order to Journals Publicity Department, Cambridge University Press, FREEPOST*, The Edinburgh Building, Shaftesbury Road, Cambridge CB2 1BR, England TEL: (0223) 325806, FAX (0223) 315052

* no postage stamp necessary if posted in UK

In US & Canada, write to: Cambridge University Press, 40 West 20th Street, New York,

NY 10011-4211, USA



1 Submission of typescripts

Two copies of the manuscript should be submitted to one of the four Executive Editors (addresses on outside front cover). The editor will acknowledge receipt of the manuscripts. It is important that authors inform the editor of any changes of address whilst their paper is under consideration.

2 Typescript

Papers should be typed, double-spaced, on one side only and with generous margins. The pages must be numbered.

The first page should give the title, the author's name and institution, and a short abstract intelligible to mathematicians.

The title, while brief, must be informative (e.g. A new proof of the ergodic theorem, whereas Some applications of a theorem of Birkhoff would be useless).

3 Notation

It is important that mathematical expressions are clear to a printer (who is not a mathematician). For instance, n_k (n sub k) is common usage, but avoid if possible using c sub n sub k. Fractions are generally best expressed by a solidus. Complicated exponentials like

$$\exp\left\{z^2\sin\theta/(1+y^2)\right\}$$

should be shown in this and no other way.

In the typescript, italics, small capitals and capitals are specified by single, double and triple underlining. Bold-faced type is shown by wavy underlining.

It helps if displayed, equations or statements which will be quoted later are numbered in order on the right of their line. They can then be referred to by, for example, 'from (7)'.

The author must enable the printer (if necessary by pencilled notes in the margin) to distinguish between similar symbols such as o, O, o O, 0; x, X, \times ; ϕ , Φ , \emptyset ; l, 1; ε , \in ; κ , k.

There is no need to underline Greek or script letters provided these are clearly typed. Any special symbols should be explained on a separate sheet of directions for the printer.

If an author wishes to mark the end of the proof of a theorem, the sign \Box may be used.

Footnotes should be avoided.

4 Diagrams

Figures and drawings should be on separate sheets in black ink. Photocopies are acceptable only if they are as clear as the originals. Symbols, legends and captions should be given on a transparent overlay. Each text figure must be numbered as Figure 1, Figure 2,... and its intended position clearly indicated in the typescript. The author's name in pencil must be on all separate sheets of diagrams.

A figure is expensive to reproduce and should be included only when the subject matter demands it, or when it greatly clarifies the exposition.

The publisher recognizes that some authors do not have the facilities for producing drawings of a sufficiently high standard to be reproduced directly and is therefore willing to have such diagrams re-drawn, provided that they are clear.

5 Tables

Tables should be numbered (above the table) and set out on separate sheets. Indicate the position of each in the text as for figures.

6 References

References should be collected at the end of the paper numbered in alphabetical order of the authors' names. A reference to a book should give the title, in italics, and then in roman type the publisher's name and the place and year of publication;

[4] N. Dunford & J. T. Schwartz Linear Operators Part I. Wiley: New York, 1958.

A reference to a paper should give in italics the title of the periodical, the number of the volume and year, and the beginning and end pages of the paper. Titles should be abbreviated as in *Mathematical Reviews*:

[6] J. E. Littlewood. The 'pits effect' for functions in the unit circle. J. Analyse Math. 23 (1970), 236-268.

7 Proofs

Authors receive one set of proofs for correction. If excessive alterations to the original manuscript are requested after the paper has been typeset, the author will be charged the cost of resetting. For papers with more than one author the proofs are sent to the first named author unless the editor receives other instructions. It is important that proofs are corrected and returned promptly.

8 Reprints

There are 100 reprints, free of charge, for each paper. For papers with several authors these reprints are divided between the authors. There are no page charges.

Ergodic theory and dynamical systems

VOLUME 11 PART 1 MARCH 1991

CONTENTS

Bick, T. A. and Coffey, J. A class of examples of D-sequences	1
Blanchard, F. Extensions à fibre constante	7
Boivin, D. and Derriennic, Y. The ergodic theorem for additive	
cocycles of \mathbb{Z}^d or \mathbb{R}^d	19
Craig, W. Floquet exponents for Jacobi fields	41
Hulse, P. Uniqueness and ergodic properties of attractive g-measures	65
Jungreis, I. A method for proving that monotone twist maps have no invariant circles	79
Kitchens, B., Marcus, B. and Trow, P. Eventual factor maps and compositions of closing maps	85
Llibre, J. and MacKay, R. S. Rotation vectors and entropy for homeomorphisms of the torus isotopic to the identity	115
Marcus, B. and Tuncel, S. The weight-per-symbol polytope and scaffolds of invariants associated with Markov chains	129
Norton, A. An area approach to wandering domains for smooth surface endomorphisms	181
Nusse, H. E. and Yorke, J. A. Analysis of a procedure for finding numerical trajectories close to chaotic saddle hyperbolic sets	189
Ramsay, A. Local product structure for group actions	209

© Cambridge University Press 1991

CAMBRIDGE UNIVERSITY PRESS

The Pitt Building, Trumpington Street, Cambridge CB2 1RP

40 West 20th Street, New York, NY 10011-4211, USA

10 Stamford Road, Oakleigh, Melbourne 3166, Australia

Printed in Great Britain by J. W. Arrowsmith Ltd, Bristol