

**ADDENDUM TO “ON THE BERGMAN KERNEL
OF HYPERCONVEX DOMAINS”,
NAGOYA MATH. J. 129 (1993), 43–52**

TAKEO OHSAWA

0. In [O-1] it was proved that for any bounded hyperconvex domain D in \mathbf{C}^2 the Bergman kernel function $K(z, w)$ of D satisfies

$$\lim_{z \rightarrow \partial D} K(z, z) = \infty.$$

In case $n = 1$, this is due to a behavior of sublevel sets of the Green function. The general case then follows by the extendability of L^2 holomorphic functions.

1. After the author finished typing the manuscript of [O-1], H. Tanigawa suggested to him an alternative proof of the one variable case. Her argument consisted of an observation that the logarithmic capacity $c_\beta(z)$ of any bounded hyperconvex domain in \mathbf{C} is exhaustive and an assertion that $K(z, z)$ is exhaustive whenever so is $c_\beta(z)$. Unfortunately, her proof of the latter statement was too difficult for the author to follow, and seemingly not to be published anywhere. Therefore, he decided to fix her idea by giving a straightforward proof to the following.

THEOREM. There exists a constant $A \in [\pi, 750\pi]$ such that, for any Riemann surface S and for any local coordinate z on S , $\sqrt{AK(z, z)} \geq c_\beta(z)$ holds.

2. N. Suita [S] conjectured that π can be taken as the above A . In fact he showed that

$$\sqrt{\pi K(z, z)} = c_\beta(z)$$

if $S = \{z \in \mathbf{C} \mid |z| < 1\}$, and that

$$\sqrt{\pi K(z, z)} > c_\beta(z)$$

if $S = \{z \in \mathbf{C} \mid r < |z| < 1\}$ for some $r \in (0, 1)$. The author hopes that our method may give a new insight into this subtle question.

Received November 19, 1993.

3. Let S be any Riemann surface, and let z be any local coordinate of S defined on a coordinate neighborhood, say U . If S admits the Green function g , the logarithmic capacity $c_\beta(z) (= c_\beta(z(p)))$ is defined by

$$-\log c_\beta(z(p)) = \lim_{q \rightarrow p} (g(p, q) + \log |z(p) - z(q)|)$$

for any $p \in U$. Otherwise we set $c_\beta(z) \equiv 0$. The Bergman kernel function $K(z) (= K(z, z))$ is defined by

$$\log K(z) = \sup \log |Q(z)|^2.$$

Here Q runs through the set $\{Q \mid Q(z) \text{ is holomorphic on } U \text{ and there exists a holomorphic 1-form } f \text{ on } S \text{ of } L^2 \text{ norm } 1 \text{ such that } Q(z)dz = f \mid U\}$.

4. For the proof of theorem we may assume that $c_\beta(z) \neq 0$, since the result is trivial otherwise. For any point $p \in U$, we shall prove that there exists a holomorphic 1-form B_p on S such that

$$B_p \mid p = c_\beta(z) dz \mid p$$

and

$$\|B_p\|^2 \leq 750 \pi.$$

Here $\| \cdot \|$ denotes the L^2 norm.

5. Let $\chi : R \rightarrow R$ be any C^∞ function satisfying $\chi(t) = 1$ on $(-\infty, 1]$, $\chi(t) = 0$ on $(2, \infty)$ and $|\chi'(t)| < 2 \log 2$ everywhere. For simplicity we put

$$g_p = g(p, \cdot).$$

Then we put

$$f_\varepsilon = \begin{cases} \chi\left(\frac{-g_p - \log \varepsilon}{\log 2}\right) c_\beta(z(p)) & \text{on } U \\ 0 & \text{on } S \setminus U. \end{cases}$$

Clearly, for sufficiently small ε , f_ε is a C^∞ function satisfying $f_\varepsilon(p) = c_\beta(z(p))$ and $\|f_\varepsilon dz\| \rightarrow 0$ as $\varepsilon \rightarrow 0$.

6. We assert that there exists a $c_0 > 0$ such that, for any $\varepsilon \in (0, c_0)$ one can find a square integrable $(1, 0)$ form α_ε on S satisfying

$$(1) \quad \bar{\partial} \alpha_\varepsilon = \bar{\partial} f_\varepsilon \wedge dz$$

$$(2) \quad \left| \int_U |z|^{-2} \alpha_\varepsilon \wedge \bar{\alpha}_\varepsilon \right| < \infty$$

and

$$(3) \quad \|\alpha_\varepsilon\|^2 < 750 \pi.$$

This suffices, since the required 1-form B_p will be obtained by putting $B_p = f_\varepsilon \wedge dz - \alpha_\varepsilon$ for sufficiently small ε .

7. For that, given ε we look for a positive number δ , a C^∞ function $\rho : S \rightarrow (0, \infty)$ and a conformal metric ds^2 on S satisfying the following conditions (i) through (iii).

$$(i) \quad i\delta \int_U |z|^{-2} \alpha \wedge \bar{\alpha} + i \int_S \alpha \wedge \bar{\alpha} \leq 5i \int_S \rho \alpha \wedge \bar{\alpha}$$

for any square integrable (1, 0) form α on S .

(ii) For any C^∞ (1,1) form β on $S \setminus \{p\}$ with $\text{supp } \beta \subset \{\log 2 < g_p + \log \varepsilon < 2 \log 2\}$, there exists a solution to $\bar{\partial}\alpha = \beta$ satisfying

$$i \int_S \rho \alpha \wedge \bar{\alpha} \leq \int_S e^{2g_p} |\beta|^2 \, dvol.$$

Here $|\beta|$ denotes the pointwise norm of β and $dvol$ denotes the volume form, both with respect to ds^2 .

$$(iii) \quad \int_S e^{2g_p} |\bar{\partial}f_\varepsilon \wedge dz|^2 \, dvol < 150 \pi.$$

8. Obviously, we are through if there exist δ , ρ and ds^2 as above.

9. It is easy to see that (iii) is satisfied if we put

$$ds^2 = 4e^{-2g_p} \varepsilon^2 (e^{-2g_p} + \varepsilon^2)^{-2} \partial g_p \bar{\partial} g_p.$$

On the other hand, a general nonsense of elementary functional analysis tells us that (ii) is satisfied provided that there exists a C^∞ positive function η on S such that,

$$(4) \quad -i(\partial\bar{\partial}\eta + \eta^{-2}\partial\eta \wedge \bar{\partial}\eta) \geq 4i\varepsilon^2 e^{-2g_p} (e^{-2g_p} + \varepsilon^2)^{-2} \partial g_p \wedge \bar{\partial} g_p$$

and

$$(5) \quad \rho \leq e^{2g_p} (\eta + \eta^2)^{-1}$$

(cf. [O-2] Theorem 1.7).

10. Therefore our problem was reduced to finding ρ and η satisfying (4), (5) and (i) for some $\delta > 0$.

11. For that, we put

$$\eta = -\log(e^{-2(g_p+1)} + \varepsilon^2) + \log(-\log(e^{-2(g_p+1)} + \varepsilon^2))$$

for $\varepsilon \in (0, e^{-1} - e^{-2})$. To simplify the computation, let

$$\psi = \log(e^{-2(g_p+1)} + \varepsilon^2).$$

Then

$$-\partial\bar{\partial}\eta = \partial\bar{\partial}\psi - \psi^{-1}\partial\bar{\partial}\psi + \psi^{-2}\partial\psi \wedge \bar{\partial}\psi$$

and

$$\eta^{-2}\partial\eta \wedge \bar{\partial}\eta = (\psi + \log(-\psi))^{-2}(1 - \psi^{-1})^2\partial\psi \wedge \bar{\partial}\psi.$$

Hence

$$-i(\partial\bar{\partial}\eta + \eta^{-2}\partial\eta \wedge \bar{\partial}\eta) \geq i\partial\bar{\partial}\psi.$$

But a straightforward computation shows that

$$i\partial\bar{\partial}\psi \geq 4i\varepsilon^2(e^{-2g_p} + \varepsilon^2)^{-2}e^{-2g_p}\partial g_p\bar{\partial}g_p.$$

Thus (4) is satisfied by the above η . As ρ , we have only to put

$$\rho = e^{2g_p}(\eta + \eta^2)^{-1}.$$

In fact, since

$$\begin{aligned} & \sup_{t>0} e^{-2t}(-\log(e^{-2(t+1)} + \varepsilon^2) + \log(-\log(e^{-2(t+1)} + \varepsilon^2))) \\ & \leq \sup_{t>0} e^{2-2T}(2T + \log T + \log 2) \\ & \leq \sup_{t>0} e^{2-2T}(3T - 1 + \log 2) \\ & < \frac{3}{2}e < 5, \end{aligned}$$

one has (i) for sufficiently small δ , in view of the behavior of ρ near p .

REFERENCES

- [O-1] Ohsawa, T., On the Bergman kernel of hyperconvex domains, Nagoya Math. J., **129** (1993), 43–52.
- [O-2] ———, On the extension of L^2 holomorphic functions III: negligible weights, to appear in Math. Z.
- [S] Suita, N., Capacities and kernels on Riemann surfaces, Arch. Rational Mech. Anal., **46** (1972), 212–217.

*Department of Mathematics
Nagoya University
Nagoya, 464-01 Japan*