



RESEARCH ARTICLE

Transference of multilinear Fourier and Schur multipliers acting on noncommutative L_p -spaces for non-unimodular groups

Gerrit Vos 

TU Delft, EWI/DIAM, Delft, The Netherlands
Email: G.M.Vos@tudelft.nl

Received: 14 September 2023; **Accepted:** 14 October 2024; **First published online:** 2 December 2024

Keywords: Locally compact groups; noncommutative L_p spaces; noncommutative harmonic analysis; Fourier multipliers; Schur multipliers

2020 Mathematics Subject Classification: *Primary* - 22D25; *Secondary* - 43A15, 46L51

Abstract

In Caspers et al. (Can. J. Math. 75[6] [2022], 1–18), transference results between multilinear Fourier and Schur multipliers on noncommutative L_p -spaces were shown for unimodular groups. We propose a suitable extension of the definition of multilinear Fourier multipliers for non-unimodular groups and show that the aforementioned transference results also hold in this more general setting.

1. Introduction

A central problem in classical harmonic analysis is finding conditions on symbols $\phi \in L_\infty(\mathbb{R}^n)$ such that the associated Fourier multiplier T_ϕ is bounded on $L_p(\mathbb{R}^n)$. One can replace \mathbb{R}^n here by any locally compact abelian group in a straightforward way. For non-abelian groups G , there is no Pontryagin dual; instead, the Fourier multiplier corresponding to a function ϕ on G is a map on the group von Neumann algebra. It is given by $\lambda_s \mapsto \phi(s)\lambda_s$ for $s \in G$, where λ is the left regular representation. Equivalently, it is given by $\lambda(f) \mapsto \lambda(\phi f)$ for $f \in L_1(G)$. It turns out that symbols $\phi \in L_\infty(G)$ give rise to bounded Fourier multipliers on $\mathcal{L}G$ precisely when ϕ defines a multiplier on the Fourier algebra $A(G)$, which coincides with the predual of $\mathcal{L}G$.

Another interesting question is which symbols $\phi \in L_\infty(G)$ give rise to a completely bounded multiplier on $\mathcal{L}G$. Bozejko and Fendler [1] showed that this happens exactly when ϕ defines a bounded Schur multiplier of Toeplitz type on $B(L_2(G))$, and in that case, the completely bounded norms are equal. This is called a transference result between Fourier and Schur multipliers. A different proof of this transference result was later provided by Jolissaint [13]. This relation between Fourier and Schur multipliers has been an important tool to prove several multiplier results. For instance, bounding the norm of Fourier multipliers by that of Schur multipliers played a crucial role in [21]. The converse transference was used in [20] to give examples of bounded multipliers on L_p -spaces that are not completely bounded. Similar transference techniques were used in [4] to prove Hörmander-Mikhlin criteria for the boundedness of Schur multipliers and in [17] to find examples of noncommutative L_p -spaces without the completely bounded approximation property.

There are several papers treating transference results for the noncommutative L_p -spaces $L_p(\mathcal{L}G)$. Neuwirth and Ricard studied this question for discrete groups [18]. In this case, the Fourier multipliers are relatively straightforward to define on $L_p(\mathcal{L}G)$. They proved that if ϕ defines a completely bounded Fourier multiplier on $L_p(\mathcal{L}G)$, then it defines a completely bounded Schur multiplier on the Schatten class $S_p(L_2(G))$. Moreover, they proved that the converse implication also holds provided that the group G is amenable. Later, Caspers and De la Salle [3] defined Fourier multipliers on the noncommutative

L_p -spaces of general locally compact groups and proved that the same transference results also hold here. An analogous result was proved by Gonzalez-Perez [10] for crossed products.

Juschenko, Todorov and Turowska [14] introduced multilinear Schur multipliers with respect to measure spaces. Such multipliers and the related notion of multiple operator integrals have been used to prove several interesting results such as the resolution of Koplienko's conjecture on higher order spectral shift functions in [22]. Therefore, it is also interesting to consider transference between multilinear Fourier and Schur multipliers. Todorov and Turowska [26] defined a multidimensional Fourier algebra and proved a transference result for multiplicatively bounded multilinear Fourier and Schur multipliers.

To consider multilinear results on noncommutative L_p -spaces, one needs a (p_1, \dots, p_n) -version of multiplicative boundedness. This was introduced by Caspers, Janssens, Krishnaswamy-Usha and Miaskiowski [5] along the lines of Pisier's characterisation of completely bounded norms in terms of Schatten norms [20, Lemma 1.7]. Moreover, they proved a bilinear transference result for discrete groups 'along the way' in [5, Proof of Theorem 7.2], in order to provide examples of L_p -multipliers for semidirect products of groups. Multilinear transference was studied in a more general sense by Caspers, Krishnaswamy-Usha and the author of the present manuscript in [6], but only for unimodular groups. They obtained a generalisation of the linear transference results. As a direct consequence of this transference result, a De Leeuw-type restriction theorem was proven for the multiplicatively bounded norms. In this paper, we complete the picture by proving a multilinear transference result for general locally compact groups.

The main difficulty for non-unimodular groups comes from the fact that the Plancherel weight is not tracial. This means that we have to deal with spatial derivatives, which in our case will just be the multiplication operator with the modular function. It will be denoted by Δ . In particular, this raises the question of how the multilinear Fourier multiplier should be defined for $p < \infty$. It turns out that, in order to prove transference results, one needs to use the definition that 'leaves the Δ 's in place'. More precisely, for 'suitable' f_i (this will be defined later) and $x_i = \Delta^{\frac{1}{2p_i}} \lambda(f_i) \Delta^{\frac{1}{2p_i}}$, the Fourier multiplier is defined as

$$T_\phi(x_1, \dots, x_n) = \int_{G^{\times n}} \phi(s_1, \dots, s_n) f_1(s_1) \dots f_n(s_n) \Delta^{\frac{1}{2p_1}} \lambda_{s_1} \Delta^{\frac{1}{2p_1}} \dots \Delta^{\frac{1}{2p_n}} \lambda_{s_n} \Delta^{\frac{1}{2p_n}} ds_1 \dots ds_n.$$

A major drawback of this definition is that it is not suitable for interpolation results when $n > 2$, unless the 'intermediate' p_i 's are all equal to ∞ , in which case it is open. All this will be discussed in Section 3. Our first main result gives the multilinear transference from Fourier multipliers as defined above to Schur multipliers. This is Theorem 4.1. The definitions of (p_1, \dots, p_n) -multiplicative norms are given in Section 2.2.

Theorem A. *Let G be a locally compact first countable group, and let $1 \leq p \leq \infty$, $1 < p_1, \dots, p_n \leq \infty$ be such that $p^{-1} = \sum_{i=1}^n p_i^{-1}$. Let $\phi \in C_b(G^{\times n})$ and define $\tilde{\phi} \in C_b(G^{\times n+1})$ by*

$$\tilde{\phi}(s_0, \dots, s_n) = \phi(s_0 s_1^{-1}, s_1 s_2^{-1}, \dots, s_{n-1} s_n^{-1}), \quad s_i \in G.$$

If ϕ is the symbol of a (p_1, \dots, p_n) -multiplicatively bounded Fourier multiplier T_ϕ of G , then $\tilde{\phi}$ is the symbol of a (p_1, \dots, p_n) -multiplicatively bounded Schur multiplier $M_{\tilde{\phi}}$ of G . Moreover,

$$\begin{aligned} \|M_{\tilde{\phi}} : S_{p_1}(L_2(G)) \times \dots \times S_{p_n}(L_2(G)) &\rightarrow S_p(L_2(G))\|_{(p_1, \dots, p_n) \rightarrow mb} \\ &\leq \|T_\phi : L_{p_1}(\mathcal{L}G) \times \dots \times L_{p_n}(\mathcal{L}G) \rightarrow L_p(\mathcal{L}G)\|_{(p_1, \dots, p_n) \rightarrow mb}. \end{aligned}$$

The proof is mostly an adaptation of the proof of [6, Theorem 3.1]. [6, Theorem 3.1] in turn has a large overlap with [5, Lemma 4.6]. For that reason, the proof of [6, Theorem 3.1] only sketches the changes compared to [5, Lemma 4.6]. As it seems undesirable to keep stacking sketches of changes, we have chosen to include the proof in full detail here. However, most of the work in generalising to the non-unimodular case comes from generalising the reduction lemmas [5, Lemmas 4.3 and 4.4]. We do this already in Section 3. Note that [5, Lemma 4.4] does not give any details for the proof even though it is not that trivial even in the unimodular case. In Lemma 3.8, we give an elegant induction argument that fills this gap.

Also, we need an extension of the intertwining result [5, Proposition 3.9] for non-unimodular groups, which we state in Proposition 4.3. We will sketch the proof in a separate technical section at the end. We also note that in [6, Theorem 3.1], the group was required to be second countable, but in the proof, actually, only first countability was needed.

For amenable groups, we also have the converse transference result. In fact, one no longer needs a continuous symbol nor the first countability condition on the group. This is Corollary 5.2.

Theorem B. *Let G be an amenable locally compact group and $1 \leq p, p_1, \dots, p_n \leq \infty$ be such that $\frac{1}{p} = \sum_{i=1}^n \frac{1}{p_i}$. Let $\phi \in L_\infty(G^{\times n})$ and define $\tilde{\phi}$ as in Theorem 1.1. If $\tilde{\phi}$ is the symbol of a (p_1, \dots, p_n) -bounded (resp. multiplicatively bounded) Schur multiplier, then ϕ is the symbol of a (p_1, \dots, p_n) -bounded (resp. multiplicatively bounded) Fourier multiplier. Moreover,*

$$\|T_\phi\|_{(p_1, \dots, p_n)} \leq \|M_{\tilde{\phi}}\|_{(p_1, \dots, p_n)}, \quad \|T_\phi\|_{(p_1, \dots, p_n)-mb} \leq \|M_{\tilde{\phi}}\|_{(p_1, \dots, p_n)-mb}.$$

Again, the proof is similar to [6] but with additional technical complications. We also abstain from using ultraproduct techniques since they were not actually necessary for the proof. It should be noted that if $p_i = \infty$ for some $1 \leq i \leq n$, then our methods only yield the above boundedness results of the multilinear Fourier multiplier on $C_\lambda^*(G)$ in the i 'th input (and conversely, boundedness on $C_\lambda^*(G)$ is all we need for the converse direction in Theorem A). Of course, if $p_1 = \dots = p_n = p = \infty$, then the result from [26] guarantees that the Fourier multiplier is indeed bounded on $(\mathcal{L}G)^{\times n}$.

As a result of Theorem B, we again get a multilinear De Leeuw-type restriction theorem.

Finally, we describe the structure of the paper. We start by giving the necessary preliminaries in Section 2. We also give a new definition of (linear) p -Fourier multipliers here. In Section 3, we discuss possible definitions of the multilinear Fourier multiplier and explain why the definition as stated above is the correct one for transference. We also prove some properties of the multilinear Fourier multiplier that we will need later. In Sections 4 and 5, we prove the transference from Fourier to Schur (Theorem A) and transference from Schur to Fourier (Theorem B), respectively. In Section 6, we sketch the proof of Proposition 4.3 using Haagerup reduction. This section is rather technical and not essential to understand the bigger picture.

2. Preliminaries

A more elaborate discussion of some of the statements in this section is contained in the author's PhD thesis [27].

2.1. L_p -spaces of group von Neumann algebras for noncommutative groups

Let G be a locally compact group, not necessarily unimodular. We will denote the left Haar measure of such a group by $\mu := \mu_G$. Its modular function will be denoted by Δ . Recall that $\Delta : G \rightarrow (\mathbb{R}_{>0}, \times)$ is a continuous group homomorphism satisfying

$$\int_G f(s^{-1})\Delta(s^{-1})d\mu(s) = \int_G f(s)d\mu(s) = \Delta(t) \int_G f(st)d\mu(s), \quad t \in G, f \in L_1(G).$$

In the sequel, we will write ds for the left Haar measure.

The left regular representation λ acts on $L_2(G)$ by $(\lambda_s f)(t) = f(s^{-1}t)$. It also defines a $*$ -representation of the $*$ -algebra $L_1(G)$ on $L_2(G)$ by the formula

$$\lambda(f)g = \int_G f(s)(\lambda_s g)ds = f * g.$$

The group von Neumann algebra $\mathcal{L}G$ of G is defined as

$$\mathcal{L}G = \{\lambda_s : s \in G\}'' = \{\lambda(f) : f \in L_1(G)\}''.$$

The group von Neumann algebra admits a canonical weight φ , named the Plancherel weight. It is defined for $x \in \mathcal{LG}$ by

$$\varphi(x^*x) = \begin{cases} \|f\|_2^2 & \text{if } x = \lambda(f) \text{ for some } f \in L_2(G) \\ \infty & \text{else.} \end{cases}$$

Here, we extend λ to all functions that define a bounded convolution operator. The Plancherel weight is tracial if and only if G is unimodular. Similarly, there is a right regular representation ρ defined by $(\rho_s f)(t) = \Delta^{1/2}(s)f(ts)$ and a Plancherel weight ψ on $(\mathcal{LG})'$ defined as $\psi(x^*x) = \|f\|_2^2$ if x is given by $x = \rho(f)$ for some $f \in L_2(G)$ and $\psi(x) = \infty$ otherwise. In practice, the group \mathcal{LG} is often semifinite even when G is not unimodular, but it is more natural to work with the Plancherel weight. For instance, the definition of φ gives the Plancherel identity (2.2) below.

Let $L_p(\mathcal{LG})$ denote the Connes–Hilsum L_p -space corresponding to the Plancherel weight ψ on $(\mathcal{LG})'$ ([7], [11]; see also [25]). To keep this paper more accessible, we will not be using the Tomita–Takesaki theory or the theory of spatial derivatives, except for the last section. Instead, we will use some facts about the Connes–Hilsum L_p -spaces as a black box. First, for $1 \leq p < \infty$, elements of $L_p(\mathcal{LG})$ are closed unbounded operators on $L_2(G)$, and sums and products of such operators are densely defined and preclosed. Addition and multiplication on $L_p(\mathcal{LG})$ are defined by taking the closures of the resulting sum resp. product, and we will use the usual addition and multiplication notations for this. For any $x \in L_p(\mathcal{LG})$, $1 \leq p \leq \infty$, we have

$$\|\lambda_s x \lambda_t\|_{L_p(\mathcal{LG})} = \|x\|_{L_p(\mathcal{LG})}, \quad s, t \in G. \quad (2.1)$$

The spatial derivative $\frac{d\varphi}{d\psi}$ is just multiplication with the modular function Δ ; see, for instance, [3, Section 3.5] for more details. With slight abuse of notation, we will denote this operator by Δ as well. The domain of this operator is exactly the set of functions $f \in L_2(G)$ such that $\int_G \Delta^2(s)|f(s)|^2 ds < \infty$. Usually, we will only apply Δ to continuous compactly supported functions so that no technical complications can arise.

Now define $L := \lambda(C_c(G) \star C_c(G))$, where $C_c(G) \star C_c(G)$ is the linear span of elements of the form $f * g$, $f, g \in C_c(G)$. For $\theta \in [0, 1]$ and $1 \leq p < \infty$, there is an embedding $\kappa_p^\theta : L \rightarrow L_p(\mathcal{LG})$ given by

$$\kappa_p^\theta(x) = \Delta^{(1-\theta)/p} x \Delta^{\theta/p}.$$

For $p = \infty$, we simply set κ_p^θ to be the identity. The images of the embeddings κ_p^θ are dense in $L_p(\mathcal{LG})$, which will be crucial in the rest of the paper.

We now state some other facts for later use. From the definitions of the L_p -norm and the Plancherel weight, we have the following Plancherel identity:

$$\|\lambda(f)\Delta^{1/2}\|_{L_2(\mathcal{LG})} = \|f\|_2, \quad f \in L_2(G) \cap L_1(G). \quad (2.2)$$

As a side note, this implies that the map $C_c(G) \mapsto \kappa_2^1(\lambda(C_c(G)))$, $f \mapsto \lambda(f)\Delta^{1/2}$ extends to a unitary $L_2(G) \cong L_2(\mathcal{LG})$. Next, a straightforward calculation yields the following commutation formulae:

$$\Delta^z \lambda_s = \Delta^z(s) \lambda_s \Delta^z, \quad z \in \mathbb{C}, s \in G \quad (2.3)$$

and

$$\Delta^z \lambda(f) = \lambda(\Delta^z f) \Delta^z, \quad z \in \mathbb{C}, f \in C_c(G). \quad (2.4)$$

Also, note that for $f = g * h$, $g, h \in C_c(G)$ and $z \in \mathbb{C}$, we have

$$(\Delta^z f)(s) = \Delta^z(s) \int_G g(t)h(t^{-1}s)dt = \int_G \Delta^z(t)g(t)\Delta^z(t^{-1}s)h(t^{-1}s)dt = ((\Delta^z g) * (\Delta^z h))(s) \quad (2.5)$$

and therefore $\Delta^z(C_c(G) \star C_c(G)) = C_c(G) \star C_c(G)$. Hence, (2.4) yields that

$$\kappa_p^\theta(L) = \kappa_p^0(L), \quad \forall \theta \in [0, 1]. \quad (2.6)$$

2.2. Operator spaces and multiplicatively bounded maps

Let E_1, \dots, E_n, E be operator spaces and $T: E_1 \times \dots \times E_n \rightarrow E$ a linear map. For $N \geq 1$, the *multiplicative amplification* $T^{(N)}: M_N(E_1) \times \dots \times M_N(E_n) \rightarrow M_N(E)$ of T is defined as

$$T^{(N)}(\alpha_1 \otimes x_1, \dots, \alpha_n \otimes x_n) = \alpha_1 \dots \alpha_n \otimes T(x_1, \dots, x_n), \quad \alpha_i \in M_N(\mathbb{C}), x_i \in E_i$$

and extended linearly. The map T is said to be *multiplicatively bounded* if

$$\|T\|_{mb} := \sup_{N \geq 1} \|T^{(N)}\| < \infty.$$

Let us now define a notion of p -completely bounded maps. Set $E = L_p(\mathcal{M}, \psi)$ for some von Neumann algebra \mathcal{M} and normal faithful semifinite weight ψ on \mathcal{M}' . Let $x \in M_N(E)$; then x is closable and $[x] \in L_p(M_N(\mathcal{M}), \text{tr}_N \otimes \psi)$. Here tr_N is the trace on $M_N(\mathbb{C})$. Now we set $S_p^N \otimes L_p(\mathcal{M}, \psi)$ to be the space $M_N(L_p(\mathcal{M}, \psi))$ equipped with the norm

$$\|x\|_{S_p^N \otimes L_p(\mathcal{M}, \psi)} := \|[x]\|_{L_p(M_N(\mathcal{M}), \text{tr}_N \otimes \psi)}.$$

We say that an operator $T: L_p(\mathcal{M}, \psi) \rightarrow L_p(\mathcal{M}, \psi)$ is *p -completely bounded* if

$$\|T\|_{p-cb} := \sup_{N \geq 1} \|T^{(N)}: S_p^N \otimes L_p(\mathcal{M}, \psi) \rightarrow S_p^N \otimes L_p(\mathcal{M}, \psi)\| < \infty.$$

The matrix norms satisfy

$$\|\alpha \otimes x\|_{S_p^N \otimes L_p(\mathcal{M}, \psi)} = \|\alpha\|_{S_p^N} \|x\|_{L_p(\mathcal{M}, \psi)}, \quad \alpha \in S_p^N, x \in L_p(\mathcal{M}, \psi). \quad (2.7)$$

For semifinite \mathcal{M} , this is an easy exercise; for arbitrary von Neumann algebras, this requires some analysis on the interplay between spatial derivatives and matrices.

We note that the norm on $S_p^N \otimes L_p(\mathcal{M}, \psi)$ does not give an operator space structure on $L_p(\mathcal{M}, \psi)$, as it does not satisfy the axioms. However, if \mathcal{M} is semifinite, then the notion of p -completely bounded maps coincides with the usual notion of completely bounded maps by [20].

We now define a (p_1, \dots, p_n) -multiplicatively bounded norm in a similar manner. Let $1 \leq p_1, \dots, p_n, p \leq \infty$ with $p^{-1} = \sum_{i=1}^n p_i^{-1}$. We define a map $T: L_{p_1}(\mathcal{M}) \times \dots \times L_{p_n}(\mathcal{M}) \rightarrow L_p(\mathcal{M})$ to be (p_1, \dots, p_n) -multiplicatively bounded if

$$\|T\|_{(p_1, \dots, p_n)-mb} := \sup_{m \geq 1} \|T^{(m)}: S_{p_1}^m \otimes L_{p_1}(\mathcal{M}) \times \dots \times S_{p_n}^m \otimes L_{p_n}(\mathcal{M}) \rightarrow S_p^m \otimes L_p(\mathcal{M})\| < \infty.$$

This turns out to be the correct notion to prove our transference results. We note that this time, there does not seem to be a case for which this coincides with ‘normal’ multiplicative boundedness, as [20, Lemma 1.7] does not generalise to the multilinear case. If \mathcal{M} is semifinite, then the above definition does coincide with the definition of (p_1, \dots, p_n) -multiplicative boundedness from [5] and [6]. Even in the semifinite case, it is unclear if this definition corresponds to the complete boundedness of some linear map on some appropriate tensor product; see [6, Remark 2.1].

2.3. Fourier multipliers

Again let G be a locally compact group. For a bounded function $\phi: G \rightarrow \mathbb{C}$, the associated Fourier multiplier $T_\phi: \mathcal{L}G \rightarrow \mathcal{L}G$ is given for $f \in L_1(G)$ by $\lambda(f) \mapsto \lambda(\phi f)$, whenever this map extends weak-* continuously. This definition has been extended to $L_p(\mathcal{L}G)$ for general locally compact groups in [3]. However, this was done only for symbols $\phi \in M_{cb}A(G)$, that is, those symbols for which T_ϕ defines a completely bounded multiplier on $\mathcal{L}G$. We give a broader definition here as preparation for the multilinear definition. Define $\mathcal{L}G_{(\gamma)}$ to be the space of closed densely defined γ -homogeneous operators on $L_2(G)$. See [25, Section III, IV] for a definition and some properties. We use here the facts that this space contains $L_p(\mathcal{L}G)$ and that the right-hand side of (2.8) is always in $\mathcal{L}G_{(-1/p)}$ ([25, III.(19) and Corollary III.34]).

For $\phi \in L_\infty(G)$, define $T_\phi : \kappa_p^\theta(L) \rightarrow \overline{\mathcal{LG}}_{(-1/p)}$ by

$$T_\phi(\kappa_p^\theta(\lambda(f))) = \Delta^{\frac{1-\theta}{p}} \lambda(\phi f) \Delta^{\frac{\theta}{p}}, \quad f \in C_c(G) \star C_c(G), \quad 1 \leq p < \infty, \quad \theta \in [0, 1] \quad (2.8)$$

The map T_ϕ does not depend on the choice of θ ; this follows from (2.6) and some manipulations using the commutation formula (2.4). When the image of T_ϕ lies in $L_p(\mathcal{LG})$ and T_ϕ extends continuously to a bounded map on $L_p(\mathcal{LG})$, we say that ϕ defines a p -Fourier multiplier. When the extension is moreover p -completely bounded on $L_p(\mathcal{LG})$, we say ϕ defines a p -cb Fourier multiplier. For $\phi \in M_{cb}A(G)$, the definition of the Fourier multiplier coincides with that of [3]. For $p = 1$, this can be shown by using [3, Proposition-Definition 3.5] and (2.10) below together with [3, Theorem 6.2 (ii), (iv)]; for other p , this follows by interpolation.

Let us now state some known results about when symbols define p - or p -cb Fourier multipliers. Let $A(G)$ be the Fourier algebra; it is defined as $A(G) = \{f * \tilde{g} : f, g \in L_2(G)\}$ where $\tilde{g}(s) = \overline{g(s^{-1})}$. It is isomorphic to $L_1(\mathcal{LG})$, the predual of \mathcal{LG} , through the pairing

$$\langle \psi, \lambda(f) \rangle = \int_G \psi(s) f(s) ds, \quad \psi \in A(G), \quad f \in L_1(G).$$

From classical theory (see, e.g. [16, Proposition 5.1.2]), it is known that $\phi \in M(A(G))$ if and only if ϕ defines a ∞ -Fourier multiplier. In the same way as in [3, Proof of Definition-Proposition 3.5], one proves that in this case, ϕ defines a p -Fourier multiplier for all $1 \leq p \leq \infty$. Trivially, $A(G) \subseteq M(A(G))$, and hence $A(G)$ provides us with plenty of symbols defining p -Fourier multipliers; we will use this fact later on. As mentioned above, if ϕ is a ∞ -cb Fourier multiplier (i.e. $\phi \in M_{cb}A(G)$), then ϕ is also a p -cb Fourier multiplier for $1 \leq p \leq \infty$. This is proven in [3, Definition-Proposition 3.5].

Let us now turn our attention to the multilinear case. In [26], $M_n^{cb}A(G)$ was defined to be the space of all symbols $\phi \in L_\infty(G^{\times n})$ such that the map

$$(\lambda_{s_1}, \dots, \lambda_{s_n}) \mapsto \phi(s_1, \dots, s_n) \lambda_{s_1 \dots s_n}$$

extends to a multiplicatively bounded normal map $(\mathcal{LG})^{\times n} \rightarrow \mathcal{LG}$. In [6], multilinear Fourier multipliers on the noncommutative L_p -spaces were defined for unimodular groups G as follows. Let $\phi \in L_\infty(G^{\times n})$ and $1 \leq p_1, \dots, p_n, p < \infty$ with $p^{-1} = \sum_{i=1}^n p_i^{-1}$. Consider the map $T_\phi : L^{\times n} \rightarrow \mathcal{LG}$ defined by

$$T_\phi(\lambda(f_1), \dots, \lambda(f_n)) = \int_{G^{\times n}} \phi(t_1, \dots, t_n) f_1(t_1) \dots f_n(t_n) \lambda_{t_1 \dots t_n} dt_1 \dots dt_n$$

for $f_i \in C_c(G) \star C_c(G)$. If this map takes values in $L_p(\mathcal{LG})$ and extends continuously to $L_{p_1}(\mathcal{LG}) \times \dots \times L_{p_n}(\mathcal{LG})$, then we say that ϕ defines a (p_1, \dots, p_n) -Fourier multiplier. The extension is again denoted by T_ϕ . In case $p_i = \infty$, we replace $L_{p_i}(\mathcal{LG})$ by $C_\lambda^*(G)$ in the i 'th coordinate. If the extension is (p_1, \dots, p_n) -multiplicatively bounded, then we say that ϕ defines a (p_1, \dots, p_n) -mb Fourier multiplier.

This definition works only for unimodular groups if $p < \infty$ since L is not contained in $L_p(\mathcal{LG})$ otherwise. In Section 3, we will give the definition for non-unimodular groups.

Remark 2.1. We note that a priori, the set of symbols of (∞, \dots, ∞) -mb Fourier multipliers is smaller than $M_n^{cb}A(G)$. However, these sets are actually the same. This follows, for instance, from a combination of our results and [26, Theorem 5.5]. One does not need the complicated machinery of Section 4 however. It follows already from the proof of [26, Theorem 5.5] or from the alternative proof of the Fourier to Schur direction in [6, Proposition 2.3], which it suffices to require that T_ϕ is bounded on $(C_\lambda^*(G))^{\times n}$.

2.4. Schatten classes and Schur multipliers

We denote by $S_p(H)$ the standard Schatten classes on the Hilbert space H . If (X, μ) is some measure space, then $S_2(L_2(X))$ can be isometrically identified with the space of kernels $L_2(X \times X)$. Through this identification, a kernel $A \in L_2(X \times X)$ corresponds to the operator $(A\xi)(s) = \int_X A(s, t)\xi(t)dt$. This should

be seen as a continuous version of matrix multiplication. We will make no distinction between an operator A and its kernel. For $1 \leq p \leq 2 \leq p' \leq \infty$ with $\frac{1}{p} + \frac{1}{p'} = 1$, the dual pairing between $S_p(L_2(X))$ and $S_{p'}(L_2(X))$ is given by

$$\langle A, B \rangle_{p,p'} = \int_{X \times X} A(s, t) B(t, s) dt ds, \quad A \in S_p(L_2(X)), B \in S_{p'}(L_2(X)). \quad (2.9)$$

This assignment is extended continuously for general $B \in S_{p'}(L_2(X))$. We refer to [17] Section 1.2] for more details.

For $\phi \in L_\infty(X^{\times n+1})$, the associated Schur multiplier is the multilinear map $S_2(L_2(X)) \times \dots \times S_2(L_2(X)) \rightarrow S_2(L_2(X))$ determined by

$$M_\phi(A_1, \dots, A_n)(t_0, t_n) = \int_{X^{\times n-1}} \phi(t_0, \dots, t_n) A_1(t_0, t_1) A_2(t_1, t_2) \dots A_n(t_{n-1}, t_n) dt_1 \dots dt_{n-1}.$$

It follows by Cauchy–Schwarz and a straightforward calculation that M_ϕ does indeed take values in $S_2(L_2(X))$ (see, for instance, [6]). Now let $1 \leq p, p_1, \dots, p_n \leq \infty$, with $p^{-1} = \sum_{i=1}^n p_i^{-1}$. Restrict M_ϕ in the i -th input to $S_2(L_2(X)) \cap S_{p_i}(L_2(X))$. Assume that this restriction maps to $S_p(L_2(X))$ and has a bounded extension to $S_{p_1}(L_2(X)) \times \dots \times S_{p_n}(L_2(X))$. Then we say that ϕ defines a (p_1, \dots, p_n) -Schur multiplier. Its extension is again denoted by M_ϕ . If M_ϕ is (p_1, \dots, p_n) -multiplicatively bounded, then we say that ϕ defines a (p_1, \dots, p_n) -mb Schur multiplier.

The following theorem is [6, Theorem 2.2]; see also [17, Theorem 1.19] and [3, Theorem 3.1]. It will be the starting point for the proof of Theorem 4.1.

Theorem 2.2. *Let μ be a Radon measure on a locally compact space X and $\phi : X^{n+1} \rightarrow \mathbb{C}$ a continuous function. Let $K > 0$. The following are equivalent for $1 \leq p_1, \dots, p_n, p \leq \infty$:*

- (i) ϕ defines a bounded Schur multiplier $S_{p_1}(L_2(X)) \times \dots \times S_{p_n}(L_2(X)) \rightarrow S_p(L_2(X))$ with norm less than K .
- (ii) For every σ -finite measurable subset X_0 in X , ϕ restricts to a bounded Schur multiplier $S_{p_1}(L_2(X_0)) \times \dots \times S_{p_n}(L_2(X_0)) \rightarrow S_p(L_2(X_0))$ with norm less than K .
- (iii) For any finite subset $F = \{s_1, \dots, s_N\} \subset X$ belonging to the support of μ , the symbol $\phi|_{F^{\times(n+1)}}$ defines a bounded Schur multiplier $S_{p_1}(\ell_2(F)) \times \dots \times S_{p_n}(\ell_2(F)) \rightarrow S_p(\ell_2(F))$ with norm less than K .

The same equivalence is true for the (p_1, \dots, p_n) -mb norms.

Now let G again be a locally compact group. In general, one has $L_p(\mathcal{L}G) \cap S_p(L_2(G)) = \{0\}$, so we cannot directly link Fourier and Schur multipliers as in the case $p = \infty$. In Section 5, we will use the following trick from [3] to circumvent this difficulty. Let $F \subseteq G$ be a relatively compact Borel subset of G with positive measure and $P_F : L_2(G) \rightarrow L_2(F)$, $f \mapsto 1_F f$ the orthogonal projection. Then for $x \in L_p(\mathcal{L}G)$, one can formally define the operator $P_F x P_F$, which lies in $S_p(L_2(G))$. We refer to [3, Proposition 3.3, Theorem 5.1] for details.

Let $F \subseteq G$ compact. We calculate the kernel of $P_F \Delta^a \lambda(f) \Delta^b P_F$: for $g \in C_c(G)$ and $s \in G$, we have

$$\begin{aligned} (P_F \Delta^a \lambda(f) \Delta^b P_F g)(s) &= 1_F(s) \Delta^a(s) \int_G f(t) \Delta^b(t^{-1}s) 1_F(t^{-1}s) g(t^{-1}s) dt \\ &= 1_F(s) \Delta^a(s) \int_G f(st) \Delta^b(t^{-1}) 1_F(t^{-1}) g(t^{-1}) dt \\ &= 1_F(s) \Delta^a(s) \int_G f(st^{-1}) \Delta^b(t) 1_F(t) \Delta(t^{-1}) g(t) dt. \end{aligned}$$

Hence the kernel of $P_F \Delta^a \lambda(f) \Delta^b P_F$ is given by

$$(s, t) \mapsto 1_F(s) \Delta^a(s) f(st^{-1}) \Delta^{b-1}(t) 1_F(t). \quad (2.10)$$

3. The definition of multilinear Fourier multipliers for non-unimodular groups

In this section, G is an arbitrary locally compact group. Let $1 \leq p_1, \dots, p_n, p \leq \infty$ with $p^{-1} = \sum_{i=1}^n p_i^{-1}$ and $\phi \in L_\infty(G^{\times n})$. In this section, we explore what a suitable definition for the Fourier multiplier $T_\phi: L_{p_1}(\mathcal{L}G) \times \dots \times L_{p_n}(\mathcal{L}G) \rightarrow L_p(\mathcal{L}G)$ might be. Our first requirement is that it must coincide with the linear definition for $n = 1$; that is, it must satisfy (2.8).

Second, we would like the definition to be compatible with interpolation arguments. More precisely, if T_ϕ is bounded as a map $\mathcal{L}G \times \dots \times \mathcal{L}G \rightarrow \mathcal{L}G$ and as a map $L_{p_1}(\mathcal{L}G) \times \dots \times L_{p_n}(\mathcal{L}G) \rightarrow L_p(\mathcal{L}G)$, then it should also be bounded as map $L_{\frac{p_1}{\nu}}(\mathcal{L}G) \times \dots \times L_{\frac{p_n}{\nu}}(\mathcal{L}G) \rightarrow L_{\frac{p}{\nu}}(\mathcal{L}G)$ for all $0 < \nu < 1$. This means that the definition must be ‘compatible’ with the definition on $(\mathcal{L}G)^{\times n}$, in the sense that in each input, the maps T_ϕ must coincide on the intersection space of some compatible couple (with respect to some θ). This tells us what the Fourier multiplier should look like on the dense subsets $\kappa_{p_i}^\theta(L)$:

Definition 3.1 (‘Wrong definition’). Let $\theta_1, \dots, \theta_n, \theta \in [0, 1]$ and $x_i = \kappa_{p_i}^{\theta_i}(\lambda(f_i))$ for $i = 1, \dots, n$, where $f_i \in C_c(G) * C_c(G)$. We set

$$T_{\phi, \text{int}}^{\theta_1, \dots, \theta_n, \theta}(x_1, \dots, x_n) = \kappa_p^\theta(T_\phi(\lambda(f_1), \dots, \lambda(f_n))). \quad (3.1)$$

Definition 3.1 might seem reasonable at first glance; it coincides with the linear definition for $n = 1$, and it is the only option if we want interpolation results. However, there are several problems with Definition 3.1. First, the definition depends on the choice of embeddings, which is not an issue in the linear case. Second, there are several properties of multilinear Fourier multipliers on unimodular groups, which do not carry over. This includes, for instance, [5, Lemma 4.3 and Lemma 4.4], which are crucial in the proof of the transference from Fourier to Schur multipliers. Moreover, if we want to prove an approximate intertwining property as in (5.1), Corollary 5.5 tells us that the definition of the Fourier multiplier has to ‘preserve products of linear multipliers’, in the sense that

$$T_\phi(x_1, \dots, x_n) = T_{\phi_1}(x_1) \dots T_{\phi_n}(x_n)$$

whenever $\phi(s_1, \dots, s_n) = \phi_1(s_1) \dots \phi_n(s_n)$. Definition 3.1 does not do this. This means that there is essentially no hope of proving the transference from Schur to Fourier multipliers either.

The above requirement on the preservation of products leads us to consider instead the following definition. Let $\theta_i \in [0, 1]$ and set $a_i = \frac{1-\theta_i}{p_i}$ and $b_i = \frac{\theta_i}{p_i}$ so that $\kappa_{p_i}^{\theta_i}(x) = \Delta^{a_i} x \Delta^{b_i}$. Now for $f_i \in C_c(G) * C_c(G)$, we formally define the Fourier multiplier corresponding to $\theta_1, \dots, \theta_n$ by

$$T_{\phi, (\theta_1, \dots, \theta_n)}(\kappa_{p_1}^{\theta_1}(\lambda(f_1)), \dots, \kappa_{p_n}^{\theta_n}(\lambda(f_n))) = \int_{G^{\times n}} \phi(t_1, \dots, t_n) f_1(t_1) \dots f_n(t_n) \times \\ \Delta^{a_1} \lambda_{t_1} \Delta^{b_1+a_2} \lambda_{t_2} \dots \Delta^{b_{n-1}+a_n} \lambda_{t_n} \Delta^{b_n} dt_1 \dots dt_n. \quad (3.2)$$

A priori, it is not clear how to define the integral in (3.2). After all, the integrand

$$H(t_1, \dots, t_n) := \phi(t_1, \dots, t_n) f_1(t_1) \dots f_n(t_n) \Delta^{a_1} \lambda_{t_1} \Delta^{b_1+a_2} \lambda_{t_2} \dots \Delta^{b_{n-1}+a_n} \lambda_{t_n} \Delta^{b_n}$$

is a function that has unbounded operators as values. However, on closer inspection, the ‘unbounded part’ of this operator doesn’t really depend on the integration variables. Indeed, using the commutation formula (2.3), we can write

$$H(t_1, \dots, t_n) \\ = \phi(t_1, \dots, t_n) f_1(t_1) \dots f_n(t_n) \Delta^{a_1}(t_1) \Delta^{a_1+a_2+b_1}(t_2) \dots \Delta^{\sum_{i=1}^n a_i + \sum_{i=1}^{n-1} b_i}(t_n) \lambda_{t_1 \dots t_n} \Delta^{1/p} \\ = \phi(t_1, \dots, t_n) (\Delta^{b_1} f_1)(t_1) \dots (\Delta^{b_n} f_n)(t_n) \lambda_{t_1 \dots t_n} \cdot \Delta^{1/p}, \quad \beta_j = \sum_{i=1}^j a_i + \sum_{i=1}^{j-1} b_i.$$

Note here that the functions $\Delta^{\beta_i} f_i$ are still in $C_c(G) \star C_c(G)$ by (2.5). Hence, a more rigorous way to define the Fourier multiplier is

$$T_{\phi, (\theta_1, \dots, \theta_n)}(\kappa_{p_1}^{\theta_1}(\lambda(f_1)), \dots, \kappa_{p_n}^{\theta_n}(\lambda(f_n))) = T_{\phi}(\lambda(\Delta^{\beta_1} f_1), \dots, \lambda(\Delta^{\beta_n} f_n)) \Delta^{1/p}.$$

However, we will keep the notation from (3.2). The integral is justified through the above arguments. The latter expression also makes clear that (3.2) takes values in the space of closed densely defined $(-1/p)$ -homogeneous operators on $L_2(G)$ (see [25, III.(19) and Corollary III.34]). Just as in the linear case, it is not clear that (3.2) takes values in $L_p(\mathcal{L}G)$ in general; this will be part of the assumptions.

It turns out that the operator $T_{\phi, (\theta_1, \dots, \theta_n)}$ in (3.2) does not depend on the choice of θ_i 's:

Proposition 3.2. *Let $1 \leq p_1, \dots, p_n, p < \infty$ and $\theta_1, \dots, \theta_n \in [0, 1]$. The maps $T_{\phi, (\theta_1, \dots, \theta_n)}$ and $T_{\phi, (0, \dots, 0)}$ coincide on the space $\kappa_{p_1}^0(L) \times \dots \times \kappa_{p_n}^0(L)$. Consequently, if one of the maps has an image in $L_p(\mathcal{L}G)$ and extends continuously to $L_{p_1}(\mathcal{L}G) \times \dots \times L_{p_n}(\mathcal{L}G)$, then the other does as well, and these extensions are equal.*

Proof. Recall that by (2.6), $\kappa_{p_i}^{\theta_i}(L) = \kappa_{p_i}^0(L)$ for $i = 1 \dots, n$. For any such i , take a_i, b_i as above, that is, so that $\kappa_{p_i}^{\theta_i}(x) = \Delta^{a_i} x \Delta^{b_i}$. Let $f_i \in C_c(G) \star C_c(G)$ and set $g_i = \Delta^{-b_i} f_i$. Then by (2.4), $\Delta^{a_i} \lambda(f_i) \Delta^{b_i} = \Delta^{1/p_i} \lambda(g_i) =: x_i$, that is, $\kappa_{p_i}^{\theta_i}(\lambda(f_i)) = \kappa_{p_i}^0(\lambda(g_i))$. By (2.3), we find

$$\begin{aligned} T_{\phi, (\theta_1, \dots, \theta_n)}(x_1, \dots, x_n) &= \int_{G^{\times n}} \phi(t_1, \dots, t_n) f_1(t_1) \dots f_n(t_n) \times \\ &\quad \Delta^{a_1} \lambda_{t_1} \Delta^{b_1+a_2} \lambda_{t_2} \dots \Delta^{b_{n-1}+a_n} \lambda_{t_n} \Delta^{b_n} dt_1 \dots dt_n \\ &= \int_{G^{\times n}} \phi(t_1, \dots, t_n) \Delta^{-b_1}(t_1) f_1(t_1) \Delta^{-b_2}(t_2) f_2(t_2) \dots \Delta^{-b_n}(t_n) f_n(t_n) \times \\ &\quad \Delta^{1/p_1} \lambda_{t_1} \Delta^{1/p_2} \dots \Delta^{1/p_n} \lambda_{t_n} dt_1 \dots dt_n \\ &= T_{\phi, (0, \dots, 0)}(\Delta^{1/p_1} \lambda(g_1), \dots, \Delta^{1/p_n} \lambda(g_n)) = T_{\phi, (0, \dots, 0)}(x_1, \dots, x_n). \end{aligned}$$

□

With this issue out of the way, we can now define Fourier multipliers independent of the choice of θ_i 's:

Definition 3.3 ('Correct definition'). *Let $1 \leq p_1, \dots, p_n, p \leq \infty$ with $p^{-1} = \sum_{i=1}^n p_i^{-1}$. Also let $\phi \in L_{\infty}(G^{\times n})$. For $i = 1, \dots, n$, take any $a_i, b_i \in [0, 1]$ such that $a_i + b_i = p_i^{-1}$. If the map*

$$T_{\phi} : \kappa_{p_1}^0(L) \times \dots \times \kappa_{p_n}^0(L) \rightarrow \overline{\mathcal{L}G}_{(-1/p)}$$

which is given for $x_i = \Delta^{a_i} \lambda(f_i) \Delta^{b_i}$ with $f_i \in C_c(G) \star C_c(G)$ by

$$\begin{aligned} T_{\phi}(x_1, \dots, x_n) &= \int_{G^{\times n}} \phi(t_1, \dots, t_n) f_1(t_1) \dots f_n(t_n) \times \\ &\quad \Delta^{a_1} \lambda_{t_1} \Delta^{b_1+a_2} \lambda_{t_2} \dots \Delta^{b_{n-1}+a_n} \lambda_{t_n} \Delta^{b_n} dt_1 \dots dt_n, \end{aligned} \quad (3.3)$$

takes values in $L_p(\mathcal{L}G)$ and extends boundedly to $L_{p_1}(\mathcal{L}G) \times \dots \times L_{p_n}(\mathcal{L}G)$ in the norm topology (in case $p_i = \infty$ for some i , we use the space $C_{\lambda}^(G)$ instead of $L_{\infty}(\mathcal{L}G) = \mathcal{L}G$ in the i 'th leg), then we say that ϕ defines a (p_1, \dots, p_n) -Fourier multiplier. We denote the extension by T_{ϕ} or $T_{\phi}^{p_1, \dots, p_n}$ when we wish to emphasise the domain of the operator. This is especially useful when writing an operator norm since writing out the full domain and codomain generally makes equations too long. If T_{ϕ} is (p_1, \dots, p_n) -multiplicatively bounded, then we say that ϕ defines a (p_1, \dots, p_n) -mb Fourier multiplier.*

Clearly, for $n = 1$, Definition 3.3 reduces to (2.8). It does not give the problems that Definition 3.1 does; as we saw already, it does not depend on the choice of embeddings. Moreover, the properties [5, Lemma 4.3 and 4.4] do carry over, as we show in Lemmas 3.6 and 3.8. Finally, it preserves products in

the following more general way: if ϕ is such that there exist $m < n$ and $\phi_1 : G^{\times m} \rightarrow \mathbb{C}$, $\phi_2 : G^{\times n-m} \rightarrow \mathbb{C}$ such that

$$\phi(s_1, \dots, s_n) = \phi_1(s_1, \dots, s_m)\phi_2(s_{m+1}, \dots, s_n),$$

then

$$T_\phi(x_1, \dots, x_n) = T_{\phi_1}(x_1, \dots, x_m)T_{\phi_2}(x_{m+1}, \dots, x_n). \quad (3.4)$$

However, we have to give up interpolation results in general. The only instances where interpolation might work is when the L_p -spaces ‘in the middle’ are all equal to $C_\lambda^*(G)$. Indeed, in that case, we can take $\theta_1 = 0$, $\theta_n = 1$, $\theta = \frac{p}{p_n}$ so that the Fourier multiplier T_ϕ from Definition 3.3 also satisfies (3.1). We note that for $n > 2$ and $p_i < \infty$ for some $2 \leq i \leq n-1$, (3.3) is not of the form (3.1) for any $\theta_1, \dots, \theta_n, \theta$, and hence T_ϕ cannot be a compatible morphism for any ‘usual’ compatible couple structures on $(\mathcal{L}G, L_{p_i}(\mathcal{L}G))_{\theta_i}$.

Remark 3.4. Although (3.1) is a necessary condition for the Fourier multiplier to allow interpolation, we have not been able to prove that it is a sufficient condition. The issue is that to prove that the mapping for (p_1, \dots, p_n) is compatible with the one for (∞, \dots, ∞) , we have to prove that they coincide on the entire intersection space $L_p(\mathcal{L}G) \cap \mathcal{L}G$ (within the appropriate compatible couple structure). However, we do not know whether L is dense in this space in the intersection norm. In fact, for $p > 2$, we do not even know if T_ϕ^2 is dense in the intersection norm.

Remark 3.5. We could have just taken (the extension of) the map $T_\phi^{\frac{1}{2}, \dots, \frac{1}{2}}$ as the definition of our Fourier multiplier. This would have allowed us to skip Proposition 3.2, and all the proofs further on in this paper would still work by approximating only with elements in the central embedding. However, the more general definition allows some flexibility to choose convenient embeddings for notation or to avoid some technicalities (in particular in Lemma 3.6).

Let us now prove some properties of the multilinear Fourier multiplier for later use. Lemmas 3.6, 3.7 and 3.8 are used in the proof of Theorem 4.1. Here, Lemma 3.6 generalises [5, Lemma 4.3], and Lemma 3.8 generalises [5, Lemma 4.4]. Since the proofs of these two lemmas require careful bookkeeping with modular functions, we will give the full details. The proof of [5, Lemma 4.4] was omitted, but it is not that trivial; our argument fills that gap.

Lemma 3.6. Let $1 \leq p_j, p \leq \infty$ and fix some $1 \leq i \leq n$. Suppose that $\phi : G^{\times n} \rightarrow \mathbb{C}$ is bounded and measurable and set for $r, t, r' \in G$:

$$\bar{\phi}(s_1, \dots, s_n; r, t, r') := \phi(rs_1, \dots, s_i t, t^{-1}s_{i+1}, \dots, s_n r').$$

Then ϕ defines a (p_1, \dots, p_n) -Fourier multiplier (resp. (p_1, \dots, p_n) -mb Fourier multiplier) iff $\bar{\phi}(\cdot; r, t, r')$ defines a (p_1, \dots, p_n) -Fourier multiplier (resp. (p_1, \dots, p_n) -mb Fourier multiplier). In that case, for $x_j \in L_{p_j}(\mathcal{L}G)$,

$$T_{\bar{\phi}(\cdot; r, t, r')}(x_1, \dots, x_n) = \lambda_r^* T_\phi(\lambda_r x_1, x_2, \dots, x_i \lambda_t, \lambda_t^* x_{i+1}, \dots, x_n \lambda_{r'}) \lambda_{r'}^*. \quad (3.5)$$

Further, we have

$$\|T_\phi^{p_1, \dots, p_n}\| = \|\bar{\phi}(\cdot; r, t, r')^{p_1, \dots, p_n}\|$$

and $(r, t, r') \mapsto T_{\bar{\phi}(\cdot; r, t, r')}$ is strongly continuous. In the multiplicatively bounded case, we have for any $N \geq 1$

$$\|(T_\phi^{p_1, \dots, p_n})^{(N)}\| = \|(\bar{\phi}(\cdot; r, t, r')^{p_1, \dots, p_n})^{(N)}\|$$

as maps $S_{p_1}^N[L_{p_1}(\mathcal{L}G)] \times \dots \times S_{p_n}^N[L_{p_n}(\mathcal{L}G)] \rightarrow S_p^N[L_p(\mathcal{L}G)]$, and $(r, t, r') \mapsto T_{\bar{\phi}(\cdot; r, t, r')}^{(N)}$ is strongly continuous.

Proof. It is straightforward to check that for $s \in G, f \in C_c(G)$, we have $\lambda_s \lambda(f) = \lambda(\lambda_s(f)) = \lambda(f(s^{-1} \cdot))$; moreover, we have

$$\lambda(f) \lambda_s = \int_G f(t) \lambda_{ts} dt = \Delta(s^{-1}) \int_G f(ts^{-1}) \lambda_t dt = \Delta(s^{-1}) \lambda(f(\cdot s^{-1})). \quad (3.6)$$

We will only make a choice for some of the embeddings and leave the rest open; this is notationally more convenient. Let $x_j = \Delta^{a_j} \lambda(f_j) \Delta^{b_j} \in L_{p_j}(\mathcal{L}G)$, with $f_j \in C_c(G) \star C_c(G)$ and $a_1 = b_i = a_{i+1} = b_n = 0$ (hence $b_1 = \frac{1}{p_1}, a_i = \frac{1}{p_i}$, etc). We compute

$$\begin{aligned} T_{\tilde{\phi}(\cdot; r, t, r')}(x_1, \dots, x_n) &= \int_{G^{\times n}} \tilde{\phi}(s_1, \dots, s_n; r, t, r') f_1(s_1) \dots f_n(s_n) \lambda_{s_1} \Delta^{b_1+a_2} \lambda_{s_2} \dots \Delta^{b_{n-1}+a_n} \lambda_{s_n} ds_1 \dots ds_n \\ &= \int_{G^{\times n}} \phi(s_1, \dots, s_n) f_1(r^{-1}s_1) \dots \Delta(t)^{-1} f_i(s_i t^{-1}) f_{i+1}(ts_{i+1}) \dots \Delta(r')^{-1} f_n(s_n (r')^{-1}) \times \\ &\quad \lambda_{r^{-1}s_1} \Delta^{b_1+a_2} \dots \Delta^{b_{i-1}+a_i} \lambda_{s_i s_{i+1}} \Delta^{b_{i+1}+a_{i+2}} \dots \Delta^{b_{n-1}+a_n} \lambda_{s_n (r')^{-1}} ds_1 \dots ds_n \\ &= \lambda_r^* T_{\phi}(\tilde{x}_1, x_2, \dots, \tilde{x}_i, \widetilde{x_{i+1}}, \dots, \tilde{x}_n) \lambda_{r'}^*. \end{aligned}$$

Here

$$\begin{aligned} \tilde{x}_1 &:= \lambda(f_1(r^{-1} \cdot)) \Delta^{b_1}; & \tilde{x}_i &:= \Delta^{-1}(t) \Delta^{a_i} \lambda(f_i(\cdot t^{-1})); \\ \widetilde{x_{i+1}} &:= \lambda(f_{i+1}(t \cdot)) \Delta^{b_{i+1}}; & \tilde{x}_n &:= \Delta^{-1}(r') \Delta^{a_n} \lambda(f_n(\cdot (r')^{-1})). \end{aligned}$$

By (3.6), we can write

$$\tilde{x}_n = \Delta^{a_n} \lambda(f_n) \lambda_{r'} = x_n \lambda_{r'}$$

and similarly

$$\tilde{x}_1 = \lambda_r x_1; \quad \tilde{x}_i = x_i \lambda_t; \quad \widetilde{x_{i+1}} = \lambda_t^* x_{i+1}.$$

Combining everything together, we conclude

$$T_{\tilde{\phi}(\cdot; r, t, r')}(x_1, \dots, x_n) = \lambda_r^* T_{\phi}(\lambda_r x_1, x_2, \dots, x_i \lambda_t, \lambda_t^* x_{i+1}, \dots, x_n \lambda_{r'}) \lambda_{r'}^*.$$

By (2.1), we have

$$\begin{aligned} \|T_{\tilde{\phi}(\cdot; r, t, r')}(x_1, \dots, x_n)\|_p &= \|T_{\phi}(\lambda_r x_1, x_2, \dots, x_i \lambda_t, \lambda_t^* x_{i+1}, \dots, x_n \lambda_{r'})\|_p \\ &\leq \|T_{\phi}^{p_1, \dots, p_n}\| \|x_1\|_{p_1} \dots \|x_n\|_{p_n}. \end{aligned}$$

Hence, on the dense subsets of elements x_j as above, we have $\|T_{\tilde{\phi}(\cdot; r, t, r')}\| \leq \|T_{\phi}\|$. If we set $\psi = \tilde{\phi}(\cdot; r, t, r')$, then $\psi(\cdot; r^{-1}, t^{-1}, (r')^{-1}) = \phi$. Hence, applying the above result to $\psi(\cdot; r^{-1}, t^{-1}, (r')^{-1})$ yields the reverse inequality. By density, we conclude that the first three statements of the lemma hold. By [15, Lemma 2.3], the (left or right) multiplication with $\lambda_s, s \in G$ is strongly continuous in s . This implies the strong continuity of $(r, t, r') \mapsto T_{\tilde{\phi}(\cdot; r, t, r')}$.

Now assume ϕ defines a (p_1, \dots, p_n) -mb Fourier multiplier and let $N \geq 1$. Denote by ι_N the $N \times N$ -identity matrix. Then by writing out the definitions and using (3.5), we find, for $x_i \in S_{p_i}^N \otimes L_{p_i}(\mathcal{L}G)$,

$$\begin{aligned} T_{\tilde{\phi}(\cdot; r, t, r')}^{(N)}(x_1, \dots, x_n) &= (\iota_N \otimes \lambda_r^*) T_{\phi}^{(N)}((\iota_N \otimes \lambda_r) x_1, \dots, x_i (\iota_N \otimes \lambda_t), (\iota_N \otimes \lambda_t^*) x_{i+1}, \dots, x_n (\iota_N \otimes \lambda_{r'})) (\iota_N \otimes \lambda_{r'}^*). \end{aligned}$$

Hence, by a complete/matrix amplified version of the above arguments, we deduce the last two statements. \square

Lemma 3.7. *Let $1 \leq p_j, p \leq \infty$ and fix some $1 \leq i \leq n$. Suppose that $\phi : G^{\times n} \rightarrow \mathbb{C}$ defines a (p_1, \dots, p_n) -Fourier multiplier and $\phi_i : G \rightarrow \mathbb{C}$ defines a p_i -Fourier multiplier. Set*

$$\bar{\phi}(s_1, \dots, s_n) = \phi(s_1, \dots, s_n) \phi_i(s_i).$$

Then $\bar{\phi}$ defines a (p_1, \dots, p_n) -Fourier multiplier and for $x_j \in L_{p_j}(\mathcal{L}G)$,

$$T_{\bar{\phi}}(x_1, \dots, x_n) = T_{\phi}(x_1, \dots, x_{i-1}, T_{\phi_i}(x_i), x_{i+1}, \dots, x_n). \quad (3.7)$$

In particular,

$$\|T_{\bar{\phi}}^{p_1, \dots, p_n}\| \leq \|T_{\phi}^{p_1, \dots, p_n}\| \|T_{\phi_i} : L_{p_i}(\mathcal{L}G) \rightarrow L_{p_i}(\mathcal{L}G)\|. \quad (3.8)$$

Proof. For $x_j \in \kappa_{p_j}^0(L)$ (or any other embedding), it follows directly from writing out the definitions that (3.7) holds (cf. (2.8)). By density, (3.7) holds for general $x_j \in L_{p_j}(\mathcal{L}G)$, which implies (3.8), so $T_{\bar{\phi}}^{p_1, \dots, p_n}$ is bounded. \square

Lemma 3.8. Let $1 \leq p_1, \dots, p_n \leq \infty$. Let $q_j^{-1} = \sum_{i=j}^n p_i^{-1}$ and suppose that $\phi_j : G \rightarrow \mathbb{C}$ defines a q_j -Fourier multiplier for $1 \leq j \leq n$. Set

$$\bar{\phi}(s_1, \dots, s_n) = \phi_1(s_1 \dots s_n) \phi_2(s_2 \dots s_n) \dots \phi_n(s_n).$$

Then $\bar{\phi}$ defines a (p_1, \dots, p_n) -Fourier multiplier, and for $x_i \in L_{p_i}(\mathcal{L}G)$, we have

$$T_{\bar{\phi}}(x_1, \dots, x_n) = T_{\phi_1}(x_1 T_{\phi_2}(x_2 \dots T_{\phi_n}(x_n) \dots)). \quad (3.9)$$

Proof. We first show (3.9) on the dense subset $\kappa_{p_1}^0(L) \times \dots \times \kappa_{p_n}^0(L)$. The lemma then follows from the boundedness of the T_{ϕ_i} together with Hölder's inequality.

We make the slightly stronger claim that for any ϕ_2, \dots, ϕ_n as in the assumptions and any $x_i \in \kappa_{p_i}^0(L)$, there exists a compactly supported function $g : G \rightarrow \mathbb{C}$ such that for all ϕ_1 as in the assumptions,

$$T_{\bar{\phi}}(x_1, \dots, x_n) = \Delta^{\frac{1}{q_1}} \lambda(\phi_1 g) = T_{\phi_1}(x_1 T_{\phi_2}(x_2 \dots T_{\phi_n}(x_n) \dots)). \quad (3.10)$$

We will prove (3.10) with induction on n . We will need this intermediate step in order to expand the outer Fourier multiplier in the right-hand side of (3.9).

The case $n=1$ follows directly from (2.8). Now assume that (3.10) holds for any choice of $\phi_1, \dots, \phi_{n-1}$ as above and x_1, \dots, x_{n-1} with $x_i \in \kappa_{p_i}^0(L)$. Fix functions ϕ_1, \dots, ϕ_n as in the assumptions and x_1, \dots, x_n so that $x_i = \Delta^{\frac{1}{p_i}} \lambda(f_i)$ for $f_i \in C_c(G) \star C_c(G)$. Take g compactly supported such that for any $\psi : G \rightarrow \mathbb{C}$ defining a q_2 -Fourier multiplier,

$$T_{\psi}(x_2 T_{\phi_3}(x_3 \dots T_{\phi_n}(x_n) \dots)) = \Delta^{\frac{1}{q_2}} \lambda(\psi g) = T_{\bar{\psi}}(x_2, \dots, x_n) \quad (3.11)$$

where $\bar{\psi}(s_2, \dots, s_n) = \psi(s_2 \dots s_n) \phi_3(s_3 \dots s_n) \dots \phi_n(s_n)$. We calculate

$$\begin{aligned} T_{\phi_1}(x_1 T_{\phi_2}(x_2 \dots T_{\phi_n}(x_n) \dots)) &\stackrel{(3.11)}{=} T_{\phi_1}(\Delta^{\frac{1}{p_1}} \lambda(f_1) \Delta^{\frac{1}{q_2}} \lambda(\phi_2 g)) \\ &\stackrel{(2.4)}{=} T_{\phi_1}(\Delta^{\frac{1}{q_1}} \lambda((\Delta^{-\frac{1}{q_2}} f_1) * (\phi_2 g))) \\ &\stackrel{(2.8)}{=} \Delta^{\frac{1}{q_1}} \lambda(\phi_1((\Delta^{-\frac{1}{q_2}} f_1) * (\phi_2 g))). \end{aligned}$$

This shows the second equality from (3.10). Continuing the previous equation,

$$\begin{aligned} T_{\phi_1}(x_1 T_{\phi_2}(x_2 \dots T_{\phi_n}(x_n) \dots)) &= \int_G \phi_1(t) \left(\int_G (\Delta^{-\frac{1}{q_2}} f_1)(s_1) (\phi_2 g)(s_1^{-1} t) ds_1 \right) \Delta^{\frac{1}{q_1}} \lambda_t dt \\ &= \int_G \int_G \phi_1(s_1 t) (\Delta^{-\frac{1}{q_2}} f_1)(s_1) (\phi_2 g)(t) \Delta^{\frac{1}{q_1}} \lambda_{s_1 t} dt ds_1 \\ &\stackrel{(2.3)}{=} \int_G f_1(s_1) \Delta^{\frac{1}{p_1}} \lambda_{s_1} \int_G \phi_1(s_1 t) \phi_2(t) g(t) \Delta^{\frac{1}{q_2}} \lambda_t dt ds_1 \\ &= \int_G f_1(s_1) \Delta^{\frac{1}{p_1}} \lambda_{s_1} \Delta^{\frac{1}{q_2}} \lambda(\phi_1(s_1 \cdot) \phi_2 g) ds_1. \end{aligned}$$

Applying (3.11) again but now with $\phi_1(s_1 \cdot) \phi_2$ in place of ψ , we get

$$\begin{aligned} & T_{\phi_1}(x_1 T_{\phi_2}(x_2 \dots T_{\phi_n}(x_n) \dots)) \\ &= \int_G f_1(s_1) \Delta^{\frac{1}{p_1}} \lambda_{s_1} \int_{G^{\times n-1}} \phi_1(s_1 s_2 \dots s_n) \phi_2(s_2 \dots s_n) \phi_3(s_3 \dots s_n) \dots \phi_n(s_n) \times \\ & \quad f_2(s_2) \dots f_n(s_n) \Delta^{\frac{1}{p_2}} \lambda_{s_2} \dots \Delta^{\frac{1}{p_n}} \lambda_{s_n} ds_2 \dots ds_n ds_1 \\ &= T_{\phi}(x_1, \dots, x_n). \end{aligned}$$

□

Finally, we calculate a convenient form for the kernel of a corner of the Fourier multiplier for use in Theorem 5.1.

Lemma 3.9. *Let $F \subseteq G$ compact and $x_i = \Delta^{a_i} \lambda(f_i) \Delta^{b_i} \in L_{p_i}(\mathcal{L}G)$ as above for $f_i \in C_c(G) \star C_c(G)$. Then the kernel of $P_F T_{\phi}(x_1, \dots, x_n) P_F$ is given by*

$$\begin{aligned} (t_0, t_n) \mapsto & 1_F(t_0) 1_F(t_n) \int_{G^{\times n-1}} \phi(t_0 t_1^{-1}, \dots, t_{n-1} t_n^{-1}) f_1(t_0 t_1^{-1}) \dots f_n(t_{n-1} t_n^{-1}) \times \\ & \Delta^{a_1}(t_0) \Delta^{b_1+a_2}(t_1) \dots \Delta^{b_n}(t_n) \Delta((t_1 \dots t_n)^{-1}) dt_1 \dots dt_{n-1}. \end{aligned}$$

Proof. Let $g \in C_c(G) \star C_c(G)$ and $t_0 \in G$. Then the function $P_F T_{\phi}^{\theta_1, \dots, \theta_n}(x_1, \dots, x_n) P_F g$ is given by

$$\begin{aligned} t_0 \mapsto & 1_F(t_0) \int_{G^{\times n}} \phi(t_1, \dots, t_n) f_1(t_1) \dots f_n(t_n) \times \\ & (\Delta^{a_1} \lambda_{t_1} \Delta^{b_1+a_2} \lambda_{t_2} \dots \Delta^{b_{n-1}+a_n} \lambda_{t_n} \Delta^{b_n} P_F g)(t_0) dt_1 \dots dt_n \\ = & 1_F(t_0) \int_{G^{\times n}} \phi(t_1, \dots, t_n) f_1(t_1) \dots f_n(t_n) \Delta^{a_1}(t_0) \Delta^{b_1+a_2}(t_1^{-1} t_0) \times \dots \\ & \Delta^{b_{n-1}+a_n}(t_{n-1}^{-1} \dots t_1^{-1} t_0) (\Delta^{b_n} 1_F g)(t_n^{-1} \dots t_1^{-1} t_0) dt_1 \dots dt_n \\ = & 1_F(t_0) \int_{G^{\times n}} \phi(t_0 t_1, t_2, \dots, t_n) f_1(t_0 t_1) f_2(t_2) \dots f_n(t_n) \Delta^{a_1}(t_0) \Delta^{b_1+a_2}(t_1^{-1}) \times \\ & \Delta^{b_2+a_3}(t_2^{-1} t_1^{-1}) \dots \Delta^{b_{n-1}+a_n}(t_{n-1}^{-1} \dots t_1^{-1}) (\Delta^{b_n} 1_F g)(t_n^{-1} \dots t_1^{-1}) dt_1 \dots dt_n \\ = & 1_F(t_0) \int_{G^{\times n}} \phi(t_0 t_1^{-1}, t_2, \dots, t_n) f_1(t_0 t_1^{-1}) f_2(t_2) \dots f_n(t_n) \Delta^{a_1}(t_0) \Delta^{b_1+a_2}(t_1) \times \\ & \Delta^{b_2+a_3}(t_2^{-1} t_1) \dots \Delta^{b_{n-1}+a_n}(t_{n-1}^{-1} \dots t_2^{-1} t_1) (\Delta^{b_n} 1_F g)(t_n^{-1} \dots t_2^{-1} t_1) \Delta(t_1^{-1}) dt_1 \dots dt_n \\ = & \dots \\ = & 1_F(t_0) \int_{G^{\times n}} \phi(t_0 t_1^{-1}, \dots, t_{n-1} t_n^{-1}) f_1(t_0 t_1^{-1}) \dots f_n(t_{n-1} t_n^{-1}) \Delta^{a_1}(t_0) \Delta^{b_1+a_2}(t_1) \times \dots \\ & \Delta^{b_{n-1}+a_n}(t_{n-1}) (\Delta^{b_n} 1_F g)(t_n) \Delta((t_1 \dots t_n)^{-1}) dt_1 \dots dt_n. \end{aligned}$$

It follows that the kernel has the required form. □

4. Fourier to Schur transference

Let G be a locally compact first countable group. In this section, we prove the transference from Fourier to Schur multipliers for such groups. An important ingredient will be the following ‘split’ coordinate-wise convolution: fix functions $\varphi_k \in C_c(G) \star C_c(G) \subseteq A(G)$ such that $\|\varphi_k\|_1 = 1$ and the supports of φ_k form a decreasing neighbourhood basis of $\{e\}$. In other words, (φ_k) is an approximate unit for the Banach

*-algebra $L_1(G)$. Note that we use the first countability of G here. Now, given a bounded function $\phi : G^{\times n} \rightarrow \mathbb{C}$ and some fixed $1 \leq i \leq n$, we define

$$\phi_{t_1, \dots, t_n}(s_1, \dots, s_n) = \phi(t_1^{-1}s_1t_2, \dots, t_{i-1}^{-1}s_{i-1}t_i, t_i^{-1}s_i, s_{i+1}t_{i+1}^{-1}, t_{i+1}s_{i+2}t_{i+2}^{-1}, \dots, t_{n-1}s_nt_n^{-1})$$

and

$$\begin{aligned} \phi_k(s_1, \dots, s_n) &:= \int_{G^{\times n}} \phi_{t_1, \dots, t_n}(s_1, \dots, s_n) \left(\prod_{j=1}^n \varphi_k(t_j) \right) dt_1 \dots dt_n \\ &= \int_{G^{\times n}} \phi_{t_1, \dots, t_n}(e, \dots, e) \left(\prod_{j=1}^i \varphi_k(s_j \dots s_it_j) \right) \left(\prod_{j=i+1}^n \varphi_k(t_js_{i+1} \dots s_j) \Delta(s_{i+1} \dots s_j) \right) dt_1 \dots dt_n. \end{aligned} \quad (4.1)$$

The last term of (4.1) in combination with Lemma 3.8 will allow us to reduce the problem to the linear case. The necessity of the ‘split’ between indices i and $i+1$ will be explained later.

Theorem 4.1. *Let G be a locally compact first countable group, and let $1 \leq p \leq \infty$, $1 < p_1, \dots, p_{n-1} \leq \infty$ be such that $p^{-1} = \sum_{i=1}^n p_i^{-1}$. Let $\phi \in C_b(G^{\times n})$ and define $\tilde{\phi} \in C_b(G^{\times n+1})$ by*

$$\tilde{\phi}(s_0, \dots, s_n) = \phi(s_0s_1^{-1}, s_1s_2^{-1}, \dots, s_{n-1}s_n^{-1}), \quad s_i \in G.$$

If ϕ defines a (p_1, \dots, p_n) -mb Fourier multiplier T_ϕ of G , then $\tilde{\phi}$ defines a (p_1, \dots, p_n) -mb Schur multiplier $M_{\tilde{\phi}}$ of G . Moreover,

$$\begin{aligned} \|M_{\tilde{\phi}} : S_{p_1}(L_2(G)) \times \dots \times S_{p_n}(L_2(G)) &\rightarrow S_p(L_2(G))\|_{(p_1, \dots, p_n)-mb} \\ &\leq \|T_\phi : L_{p_1}(\mathcal{L}G) \times \dots \times L_{p_n}(\mathcal{L}G) \rightarrow L_p(\mathcal{L}G)\|_{(p_1, \dots, p_n)-mb}. \end{aligned}$$

Proof. Let $F \subseteq G$ finite with $|F| = N$. By Theorem 2.2, it suffices to show that the norm of

$$M_{\tilde{\phi}} : S_{p_1}(\ell_2(F)) \times \dots \times S_{p_n}(\ell_2(F)) \rightarrow S_p(\ell_2(F))$$

and its matrix amplifications are bounded.

For $s \in F$, let $p_s \in B(\ell_2(F))$ be the orthogonal projection onto the span of δ_s . Define the unitary $U = \sum_{s \in F} p_s \otimes \lambda_s \in B(\ell_2(F)) \otimes \mathcal{L}G$. In the case $p = \infty$, the Fourier to Schur transference is proven through the transference identity

$$T_\phi^{(N)}(U(a_1 \otimes 1)U^*, \dots, U(a_n \otimes 1)U^*) = U(M_{\tilde{\phi}}(a_1, \dots, a_n) \otimes 1)U^*, \quad a_i \in B(\ell_2(F)).$$

The idea is to do something similar in the case $p < \infty$. However, the unit does not embed in $L_p(\mathcal{L}G)$, so we need to use some approximation of the unit instead. We construct this as follows: let $\mathcal{V} = (V_i)_{i \in \mathbb{N}}$ be a decreasing symmetric neighbourhood basis of the identity (this is possible because G is first countable). For $V \in \mathcal{V}$, we define the operator

$$k_V = \|1_V \Delta^{-1/4}\|_2^{-1} \lambda(1_V \Delta^{-1/4}) \Delta^{1/2} \in L_2(\mathcal{L}G)$$

which is proven to be self-adjoint in [8, Section 8.3]. Let $k_V = u_V h_V$ be its polar decomposition. Then we have $h_V^{2/p} \in L_p(\mathcal{L}G)$, and by (2.2), $\|h_V^{2/p}\|_p = 1$. Now for any $V \in \mathcal{V}$, we have, by (2.7),

$$\|M_{\tilde{\phi}}(a_1, \dots, a_n)\|_{S_p^N} = \|M_{\tilde{\phi}}(a_1, \dots, a_n) \otimes h_V^{\frac{2}{p}}\|_{S_p^N \otimes L_p(\mathcal{L}G)}, \quad a_i \in B(\ell_2(F)). \quad (4.2)$$

Next, fix an $i \in \{1, \dots, n\}$ such that $\bar{p}_1 := (\sum_{l=1}^i p_l^{-1})^{-1} > 1$ and $\bar{p}_2 := (\sum_{l=i+1}^n p_l^{-1})^{-1} > 1$. This is possible by our assumption that $p_1, \dots, p_n > 1$. We now define the functions ϕ_{t_1, \dots, t_n} and ϕ_k as in (4.1) for the chosen i .

The condition $\bar{p}_1, \bar{p}_2 > 1$ is necessary for the use of Proposition 4.3 at the end of the proof of Lemma 4.2; this also explains why we need the ‘split’ in the pointwise convolutions. If $p > 1$, then one can take $i = n$ in which case the proof of Lemma 4.2 simplifies somewhat. Note that in [5] and [6], the convolutions were defined for $i = n - 1$. In the latter paper, this creates a problem in case $p_n = \infty, p = 1$; this problem is resolved by splitting instead at some i chosen as above.

Let $a_1, \dots, a_n \in B(\ell_2(F))$. By continuity of ϕ , we have that $\phi_k \rightarrow \phi$ pointwise. Indeed, for $\varepsilon > 0$, we can take K such that for $t_1, \dots, t_n \in \text{supp} \varphi_K$, $|\phi_{t_1, \dots, t_n}(s_1, \dots, s_n) - \phi(s_1, \dots, s_n)| < \varepsilon$. Then for $k > K$, we get $|\phi_k(s_1, \dots, s_n) - \phi(s_1, \dots, s_n)| < \varepsilon$. Since we are working in finite dimensions, this implies

$$M_{\phi_k}(a_1, \dots, a_n) \rightarrow M_{\tilde{\phi}}(a_1, \dots, a_n)$$

in S_p^N . Together with (4.2), we find

$$\begin{aligned} \|M_{\tilde{\phi}}(a_1, \dots, a_n)\|_{S_p^N} &= \lim_k \limsup_{V \in \mathcal{V}} \|M_{\phi_k}(a_1, \dots, a_n) \otimes h_V^{\frac{2}{p}}\|_{S_p^N \otimes L_p(\mathcal{L}G)} \\ &= \lim_k \limsup_{V \in \mathcal{V}} \|U(M_{\phi_k}(a_1, \dots, a_n) \otimes h_V^{\frac{2}{p}})U^*\|_{S_p^N \otimes L_p(\mathcal{L}G)} \\ &\leq \lim_k \limsup_{V \in \mathcal{V}} \|T_{\phi_k}^{(N)}(U(a_1 \otimes h_V^{\frac{2}{p_1}})U^*, \dots, U(a_n \otimes h_V^{\frac{2}{p_n}})U^*)\|_{S_p^N \otimes L_p(\mathcal{L}G)} \\ &\quad + \lim_k \limsup_{V \in \mathcal{V}} \|T_{\phi_k}^{(N)}(U(a_1 \otimes h_V^{\frac{2}{p_1}})U^*, \dots, U(a_n \otimes h_V^{\frac{2}{p_n}})U^*) \\ &\quad - U(M_{\phi_k}(a_1, \dots, a_n) \otimes h_V^{\frac{2}{p}})U^*\|_{S_p^N \otimes L_p(\mathcal{L}G)} \\ &:= A + B. \end{aligned}$$

First, we have

$$\begin{aligned} A &\leq \lim_k \limsup_{V \in \mathcal{V}} \|T_{\phi_k}^{(N)}\| \|a_1 \otimes h_V^{\frac{2}{p_1}}\|_{S_{p_1}^N \otimes L_{p_1}(\mathcal{L}G)} \dots \|a_n \otimes h_V^{\frac{2}{p_n}}\|_{S_{p_n}^N \otimes L_{p_n}(\mathcal{L}G)} \\ &= \lim_k \|T_{\phi_k}^{(N)}\| \|a_1\|_{S_{p_1}^N} \dots \|a_n\|_{S_{p_n}^N}. \end{aligned}$$

By repeated use of Lemma 3.6 (in particular, we can use Fubini because of the strong continuity property) we find

$$\|T_{\phi_k}^{(N)}\| \leq \int_{G^{\times n}} \|T_{\phi}^{(N)}\| \left(\prod_{i=1}^n |\varphi_k(t_j)| \right) dt_1 \dots dt_n = \|T_{\phi}^{(N)}\| \|\varphi_k\|_1^n = \|T_{\phi}^{(N)}\| \leq \|T_{\phi}\|_{(p_1, \dots, p_n) - mb}.$$

and hence

$$A \leq \|T_{\phi}\|_{(p_1, \dots, p_n) - mb} \|a_1\|_{S_{p_1}^N} \dots \|a_n\|_{S_{p_n}^N}.$$

It remains to show that $B = 0$. By the triangle inequality, it suffices to show this for $a_i = E_{r_{i-1}, r_i}$, $r_0, \dots, r_n \in F$ (for other combinations of matrix units, the term below becomes 0). In that case we get, by applying Lemma 3.6 in the second equality:

$$\begin{aligned} &T_{\phi_k}^{(N)}(U(E_{r_0, r_1} \otimes h_V^{\frac{2}{p_1}})U^*, \dots, U(E_{r_{n-1}, r_n} \otimes h_V^{\frac{2}{p_n}})U^*) - U(M_{\phi_k}(E_{r_0, r_1}, \dots, E_{r_{n-1}, r_n}) \otimes h_V^{\frac{2}{p}})U^* \\ &= E_{r_0, r_n} \otimes \left(T_{\phi_k}(\lambda_{r_0} h_V^{\frac{2}{p_1}} \lambda_{r_1}^*, \dots, \lambda_{r_{n-1}} h_V^{\frac{2}{p_n}} \lambda_{r_n}^*) - \phi_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1}) \lambda_{r_0} h_V^{\frac{2}{p}} \lambda_{r_n}^* \right) \\ &= E_{r_0, r_n} \otimes \lambda_{r_0} \left(T_{\phi_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1})}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}}) - \phi_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1}) h_V^{\frac{2}{p}} \right) \lambda_{r_n}^*. \end{aligned}$$

Hence,

$$B = \lim_k \limsup_{V \in \mathcal{V}} \left\| T_{\phi_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1})}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}}) - \phi_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1}) h_V^{\frac{2}{p}} \right\|_{L_p(\mathcal{L}G)}. \quad (4.3)$$

The limit over k exists and is 0; we postpone the proof to Lemma 4.2 below.

For the multiplicatively bounded estimate, we prove using similar methods that for $K \geq 1$ and $a_1, \dots, a_n \in M_K(B(\ell_2(F)))$,

$$\|M_{\phi}^{(K)}(a_1, \dots, a_n)\|_{S_p^{KN}} \leq \|T_{\phi}^{(KN)}\| \|a_1\|_{S_p^{KN}} \dots \|a_n\|_{S_p^{KN}}.$$

Here, we use $1_{M_K} \otimes U$ in place of U . Moreover, by the triangle inequality, it suffices to prove the estimate for B for $a_i = E_{j_{i-1}, j_i} \otimes E_{r_{i-1}, r_i}$, with $1 \leq j_i \leq K$ and $r_i \in F$; the expression for B then reduces to (4.3) again. \square

The following Lemma is similar to [5, Lemma 4.6]. In our case, we have $x_j = 1$, which allows us to avoid the SAIN condition used in that paper; on the other hand, we work with translated functions, and our result works for non-unimodular groups. Already in [6, Theorem 3.1], it was explained how to adapt the proof of [5, Lemma 4.6] for the translated functions. However, this paper only considered unimodular groups. Here, we spell out the full proof for the convenience of the reader.

Lemma 4.2. *In the proof of Theorem 4.1, we have that*

$$\lim_k \limsup_{V \in \mathcal{V}} \left\| T_{\phi_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1})}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}}) - \phi_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1}) h_V^{\frac{2}{p}} \right\|_{L_p(\mathcal{L}G)} = 0.$$

The main idea is to reduce the problem to the linear case using (4.1) and apply the following result for linear Fourier multipliers:

Proposition 4.3. *Let \mathcal{V} be a symmetric neighbourhood basis of the identity of G . Let $2 \leq q < p \leq \infty$ or $1 \leq p < q \leq 2$. Assume $\psi \in C_b(G)$ defines a Fourier multiplier on $L_p(\mathcal{L}G)$. Then we have*

$$\lim_{V \in \mathcal{V}} \|T_\psi(h_V^{2/q}) - \psi(1)h_V^{2/q}\|_{L_q(\mathcal{L}G)} \rightarrow 0.$$

The proof of Proposition 4.3 is essentially a matter of combining results and remarks from [5, Proposition 3.9] and [8, Claim B and Section 8] and applying Haagerup reduction to [9, Lemma 3.1] to generalise that estimate to general von Neumann algebras. We give more details in Section 6.

Proof of Lemma 4.2. The idea is to use a dominated convergence argument in the last expression of (4.1). However, the functions ϕ_k need not be integrable. We work around this by multiplying with compactly supported functions that are close to 1 around e so that as $V \in \mathcal{V}$ decreases to $\{e\}$, we are just ‘multiplying by 1’ in the limit. Define a function $\zeta \in C_c(G) \cap A(G)$ with $\zeta(e) = 1$, which is positive definite and (therefore) satisfies $\|T_\zeta : L_p(\mathcal{L}G) \rightarrow L_p(\mathcal{L}G)\| \leq 1$ for all $1 \leq p \leq \infty$. Next let

$$\zeta_j(s) = \zeta(r_{j-1}^{-1} s r_j), \quad 1 \leq j \leq n, \quad s \in G.$$

We define a product function as follows:

$$(\phi(\zeta_1, \dots, \zeta_n))(s_1, \dots, s_n) = \phi(s_1, \dots, s_n) \zeta_1(s_1) \dots \zeta_n(s_n).$$

Then

$$\begin{aligned} & \|T_{\phi_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1})}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}}) - \phi_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1}) h_V^{\frac{2}{p}}\|_{L_p(\mathcal{L}G)} \\ & \leq \|(\phi(\zeta_1, \dots, \zeta_n))_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1}) h_V^{\frac{2}{p}} - \phi_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1}) h_V^{\frac{2}{p}}\|_{L_p(\mathcal{L}G)} \\ & \quad + \|T_{(\phi(\zeta_1, \dots, \zeta_n))_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1})}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}}) - (\phi(\zeta_1, \dots, \zeta_n))_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1}) h_V^{\frac{2}{p}}\|_{L_p(\mathcal{L}G)} \\ & \quad + \|T_{\phi_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1})}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}}) - T_{(\phi(\zeta_1, \dots, \zeta_n))_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1})}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}})\|_{L_p(\mathcal{L}G)} \\ & =: A_{k,V} + B_{k,V} + C_{k,V}. \end{aligned}$$

Here, $\phi((\zeta_1, \dots, \zeta_n))_k$ is defined again by (4.1) for the same i . We will estimate these terms separately. We start by showing that $\lim_k \limsup_{V \in \mathcal{V}} A_{k,V}$ and $\lim_k \limsup_{V \in \mathcal{V}} C_{k,V}$ are 0, essentially reducing the problem to the integrable functions $\phi(\zeta_1, \dots, \zeta_n)$. We then apply the idea mentioned above to show that $\lim_{V \in \mathcal{V}} B_{k,V} = 0$ for any k .

First, since $\psi_k \rightarrow \psi$ pointwise for any $\psi \in C_b(G)^{\times n}$, we have

$$\begin{aligned} \limsup_{V \in \mathcal{V}} A_{k,V} &= |(\phi(\zeta_1, \dots, \zeta_n))(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1}) - \phi_k(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1})| \\ &\rightarrow |(\phi(r_0 r_1^{-1}, \dots, r_{n-1} r_n^{-1}))(1 - \zeta_1(r_0 r_1^{-1}) \dots \zeta_n(r_{n-1} r_n^{-1}))| = 0. \end{aligned}$$

Next, we estimate the limit in k of $\limsup_{V \in \mathcal{V}} C_{k,V}$. Set $u_j = t_j^{-1} r_{j-1}$, $v_j = t_j r_j$ and

$$C_V(t_1, \dots, t_n) = \|T_\eta(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}})\|_{L_p(\mathcal{L}G)},$$

where

$$\eta = (\phi - \phi(\zeta_1, \dots, \zeta_n))(u_1 \cdot u_2^{-1}, \dots, u_{i-1} \cdot u_i^{-1}, u_i \cdot r_i^{-1}, r_i \cdot v_{i+1}^{-1}, v_{i+1} \cdot v_{i+2}^{-1}, \dots, v_{n-1} \cdot v_n^{-1}).$$

Thanks to the strong continuity statement of Lemma 3.6, we can use Fubini to deduce

$$C_{k,V} \leq \int_{G^{\times n}} C_V(t_1, \dots, t_n) \left(\prod_{i=1}^n |\varphi_k(t_j)| \right) dt_1 \dots dt_n. \quad (4.4)$$

Now set

$$\begin{aligned} y_{j,V} &= \lambda_{u_j} h_V^{\frac{2}{p_j}} \lambda_{u_{j+1}}^* \text{ for } 1 \leq j \leq i-1, & y_{i,V} &= \lambda_{u_i} h_V^{\frac{2}{p_i}} \lambda_{r_i}^*, & y_{i+1,V} &= \lambda_{r_i} h_V^{\frac{2}{p_{i+1}}} \lambda_{v_{i+1}}^*, \\ y_{j,V} &= \lambda_{v_{j-1}} h_V^{\frac{2}{p_j}} \lambda_{v_j}^* \text{ for } i+2 \leq j \leq n. \end{aligned}$$

Denote ι_q for the identity operator on $L_q(\mathcal{L}G)$. The symbol 1 is used both for the constant 1-function and the number 1. Then we get the following estimate, where we apply Lemma 3.6 and (2.1) in the first line and Lemma 3.7 and the assumption that T_ζ is a contraction in the third line:

$$\begin{aligned} C_V(t_1, \dots, t_n) &= \|T_{(\phi - \phi(\zeta_1, \dots, \zeta_n))}(y_{1,V}, \dots, y_{n,V})\|_{L_p(\mathcal{L}G)} \\ &\leq \sum_{j=1}^n \|T_{\phi(1, \dots, 1, (\zeta_j - 1), \zeta_{j+1}, \dots, \zeta_n)}(y_{1,V}, \dots, y_{n,V})\|_{L_p(\mathcal{L}G)} \\ &\leq \|T_\phi : L_{p_1} \times \dots \times L_{p_n} \rightarrow L_p\| \sum_{j=1}^n \left(\|(T_{\zeta_j} - \iota_{p_j})(y_{j,V})\|_{L_{p_j}(\mathcal{L}G)} \prod_{i \neq j} \|y_{i,V}\|_{p_i} \right). \end{aligned} \quad (4.5)$$

By (2.1), we have $\|y_{j,V}\|_{p_j} = 1$. Further, by applying again Lemma 3.6 and Proposition 4.3,

$$\|(T_{\zeta_j} - \iota_{p_j})(y_{j,V})\|_{L_{p_j}(\mathcal{L}G)} = \|T_{\zeta_j(u_j u_{j+1}^{-1}) - 1}(h_V^{\frac{2}{p_j}})\|_{L_{p_j}(\mathcal{L}G)} \rightarrow |\zeta_j(u_j u_{j+1}^{-1}) - 1|$$

for $1 \leq j \leq i-1$. Filling in the definition of ζ_j ,

$$|\zeta_j(u_j u_{j+1}^{-1}) - 1| = |\zeta(r_{j-1}^{-1} t_j^{-1} r_{j-1} r_j^{-1} t_{j+1} r_j) - 1|$$

and this equals 0 when evaluated at $t_j, t_{j+1} = e$. Similarly, we find for $i \leq j \leq n$ that $\lim_{V \in \mathcal{V}} \|(T_{\zeta_j} - \iota)(y_{j,V})\|_{L_{p_j}(\mathcal{L}G)}$ exists and equals 0 when evaluated at the identity in the corresponding t_1, \dots, t_n . Moreover, all these values are bounded by 2. Going back to (4.4), let us write $M := \|T_\phi : L_{p_1} \times \dots \times L_{p_n} \rightarrow L_p\|$. We find

$$C_{k,V} \leq \int_{G^{\times n}} M \left(\sum_{j=1}^n \|(T_{\zeta_j} - \iota)(y_{j,V})\|_{L_{p_j}(\mathcal{L}G)} \right) \left(\prod_{j=1}^n |\varphi_k(t_j)| \right) dt_1 \dots dt_n. \quad (4.6)$$

The integrand of (4.6) is bounded by the integrable function $2M \prod_{i=1}^n |\varphi_k(t_j)|$. Hence, by Lebesgue's dominated convergence theorem, the right-hand side of (4.6) converges in V . We find that

$$\limsup_{V \in \mathcal{V}} C_{k,V} \leq M \int_{G^{\times n}} \left(\sum_{j=1}^n \lim_{V \in \mathcal{V}} \|(T_{\zeta_j} - \iota)(y_{j,V})\|_{L_{p_j}(\mathcal{L}G)} \right) \left(\prod_{i=1}^n |\varphi_k(t_j)| \right) dt_1 \dots dt_n.$$

This quantity goes to 0 in k . This concludes the proof for $C_{k,V}$.

Finally, we prove that $\lim_{V \in \mathcal{V}} B_{k,V} = 0$ for any k . We fix a k for the remainder of the proof. Recall that since $\varphi_k \in A(G)$, T_{φ_k} is bounded on $L_q(\mathcal{L}G)$ for any $1 \leq q \leq \infty$. Moreover, since $\varphi_k \in C_c(G) \star C_c(G)$, we also have $\varphi_k \Delta \in C_c(G) \star C_c(G) \subseteq A(G)$ (cf. the calculation before (2.6)), hence $T_{\varphi_k \Delta}$ is also bounded on $L_q(\mathcal{L}G)$ for any q .

We may assume, by scaling φ_k if necessary, that $T_{\varphi_k} : L_q(\mathcal{L}G) \rightarrow L_q(\mathcal{L}G)$ and $T_{\varphi_k \Delta} : L_q(\mathcal{L}G) \rightarrow L_q(\mathcal{L}G)$ are contractions for any Hölder combination q of p_1, \dots, p_n . Of course, this means that $\|\varphi_k\|_1$ need no longer be 1 from now on. Set

$$\begin{aligned} \psi_k(s_1, \dots, s_n; t_1, \dots, t_n) &:= \left(\prod_{j=1}^i \varphi_k(r_{j-1} s_j \dots s_i r_i^{-1} t_j) \right) \\ &\quad \times \left(\prod_{j=i+1}^n \varphi_k(t_j r_i s_{i+1} \dots s_j r_j^{-1}) \Delta(r_i s_{i+1} \dots s_j r_j^{-1}) \right) \\ &=: \psi_k^1(s_1, \dots, s_i; t_1, \dots, t_i) \psi_k^2(s_{i+1}, \dots, s_n; t_{i+1}, \dots, t_n). \end{aligned}$$

By using the last term of (4.1) and Fubini, we get

$$\begin{aligned} B_{k,V} &\leq \int_{G^{\times n}} |(\phi(\zeta_1, \dots, \zeta_n))_{t_1, \dots, t_n}(e, \dots, e)| \\ &\quad \times \|T_{\psi_k(\cdot; t_1, \dots, t_n)}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}}) - \psi_k(1, \dots, 1; t_1, \dots, t_n) h_V^{\frac{2}{p}}\|_{L_p(\mathcal{L}G)} dt_1 \dots dt_n. \end{aligned} \quad (4.7)$$

Note that $|\varphi_k| \leq 1$ by the assumed contractivity of T_{φ_k} . Indeed, for $s \in G$, apply T_{φ_k} to λ_s to deduce that $|\varphi_k(s)| \leq 1$. Hence, $|\psi_k(1, \dots, 1; t_1, \dots, t_n)| \leq \prod_{j=i+1}^n \Delta(r_i r_j^{-1})$. Moreover, from the expression (4.9) below, we see that $\|T_{\psi_k(\cdot; t_1, \dots, t_n)}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}})\|_{L_p(\mathcal{L}G)} \leq \Delta((t_{i+1}, \dots, t_n)^{-1})$. Since $\phi(\zeta_1, \dots, \zeta_n)$ is compactly supported, the integrand of (4.7) is dominated by an integrable function. Hence by the Lebesgue dominated convergence theorem, it suffices to show that the term

$$\|T_{\psi_k(\cdot; t_1, \dots, t_n)}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}}) - \psi_k(1, \dots, 1; t_1, \dots, t_n) h_V^{\frac{2}{p}}\|_{L_p(\mathcal{L}G)} \quad (4.8)$$

goes to 0 in V for any choice of $t_1, \dots, t_n \in G$.

Fix $t_1, \dots, t_n \in G$. For $1 \leq j \leq i$, set $q_j^{-1} = \sum_{l=j}^i p_l^{-1}$ (so $q_1 = \bar{p}_1$) and $T_j = T_{\varphi_k(r_{j-1} r_i^{-1} t_j)}$. By Lemma 3.6, T_j is a contraction on $L_{q_j}(\mathcal{L}G)$. For $i+1 \leq j \leq n$, set $q_j^{-1} = \sum_{l=j}^n p_l^{-1}$ (so $q_{i+1} = \bar{p}_2$) and $T_j = T_{\varphi_k(t_j r_i r_j^{-1}) \Delta(r_i r_j^{-1})}$. We can estimate the norm of $T_j : L_{q_j}(\mathcal{L}G) \rightarrow L_{q_j}(\mathcal{L}G)$ by using again Lemma 3.6:

$$\|T_j\| = \Delta(t_j^{-1}) \|T_{\varphi_k(t_j r_i r_j^{-1}) \Delta(r_i r_j^{-1})}\| = \Delta(t_j^{-1}) \|T_{\varphi_k \Delta}\| \leq \Delta(t_j^{-1}).$$

Now, by Lemma 3.8, we have

$$\begin{aligned} T_{\psi_k^1(\cdot; t_1, \dots, t_i)}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_i}}) &= T_1(h_V^{\frac{2}{p_1}} T_2(h_V^{\frac{2}{p_2}} \dots T_i(h_V^{\frac{2}{p_i}}) \dots)), \\ T_{\psi_k^2(\cdot; t_{i+1}, \dots, t_n)}(h_V^{\frac{2}{p_{i+1}}}, \dots, h_V^{\frac{2}{p_n}}) &= T_{i+1}(h_V^{\frac{2}{p_{i+1}}} T_{i+2}(h_V^{\frac{2}{p_{i+2}}} \dots T_n(h_V^{\frac{2}{p_n}}) \dots)). \end{aligned}$$

Clearly, $T_{\psi_k^1(\cdot; t_1, \dots, t_i)}$ is contractive as a map on $L_{p_1}(\mathcal{L}G) \times \dots \times L_{p_i}(\mathcal{L}G)$. Let $x_j \in L_{p_j}(\mathcal{L}G)$ with $\|x_j\|_{L_{p_j}(\mathcal{L}G)} \leq 1$; then, from (3.4),

$$\begin{aligned} \|T_{\psi_k(\cdot; t_1, \dots, t_n)}(x_1, \dots, x_n)\|_{L_p(\mathcal{L}G)} &\leq \|T_{\psi_k^2(\cdot; t_{i+1}, \dots, t_n)}(x_{i+1}, \dots, x_n)\|_{L_{\bar{p}_2}(\mathcal{L}G)} \\ &\leq \Delta((t_{i+1} \dots t_n)^{-1}). \end{aligned} \quad (4.9)$$

This validates the use of the dominated convergence theorem above. Now we go back to estimating (4.8). Using subsequently the triangle inequality and Hölder's inequality (with again [3.4]), we find

$$\begin{aligned}
& \|T_{\psi_k(\cdot; t_1, \dots, t_n)}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}}) - \psi_k(1, \dots, 1; t_1, \dots, t_n)h_V^{\frac{2}{p}}\|_{L_p(\mathcal{L}G)} \\
& \leq \|T_{\psi_k^1(\cdot; t_1, \dots, t_i)}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_i}}) \cdot \psi_k^2(1, \dots, 1; t_{i+1}, \dots, t_n)h_V^{\frac{2}{p_2}} - \psi_k(1, \dots, 1; t_1, \dots, t_n)h_V^{\frac{2}{p}}\|_{L_p(\mathcal{L}G)} \\
& \quad + \|T_{\psi_k(\cdot; t_1, \dots, t_n)}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_n}}) - T_{\psi_k^1(\cdot; t_1, \dots, t_i)}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_i}}) \cdot \psi_k^2(1, \dots, 1; t_{i+1}, \dots, t_n)h_V^{\frac{2}{p_2}}\|_{L_p(\mathcal{L}G)} \\
& \leq \left(\prod_{j=i+1}^n \Delta(r_j r_j^{-1}) \right) \|T_{\psi_k^1(\cdot; t_1, \dots, t_i)}(h_V^{\frac{2}{p_1}}, \dots, h_V^{\frac{2}{p_i}}) - \psi_k^1(1, \dots, 1; t_1, \dots, t_i)h_V^{\frac{2}{p_1}}\|_{L_{\bar{p}_1}(\mathcal{L}G)} \\
& \quad + \|T_{\psi_k^2(\cdot; t_{i+1}, \dots, t_n)}(h_V^{\frac{2}{p_{i+1}}}, \dots, h_V^{\frac{2}{p_n}}) - \psi_k^2(1, \dots, 1; t_{i+1}, \dots, t_n)h_V^{\frac{2}{p_2}}\|_{L_{\bar{p}_2}(\mathcal{L}G)} \\
& =: B_{k,V}^1 + B_{k,V}^2.
\end{aligned}$$

We show only that $\lim_{V \in \mathcal{V}} B_{k,V}^2 = 0$; the equality $\lim_{V \in \mathcal{V}} B_{k,V}^1 = 0$ follows similarly and is in fact slightly easier since the T_j are contractions for $j \leq i$. Now set, for $i \leq j \leq n$,

$$R_{j,V} := \left(\prod_{l=j+1}^n \varphi_k(t_l r_l r_l^{-1}) \right) T_{i+1}(h_V^{\frac{2}{p_{i+1}}}) \dots T_j(h_V^{\frac{2}{p_j}}) \dots.$$

Here $R_{i,V} = \prod_{l=i+1}^{n-1} \varphi_k(t_l r_l r_l^{-1}) h_V^{\frac{2}{q_1}}$. Then

$$B_{k,V}^2 \leq \sum_{j=i+1}^n \|R_{j,V} - R_{j-1,V}\|_{L_{\bar{p}_2}(\mathcal{L}G)}.$$

Recall that $|\varphi_k| \leq 1$. Hence

$$\begin{aligned}
& \|R_{j,V} - R_{j-1,V}\|_{L_{\bar{p}_2}(\mathcal{L}G)} \\
& = \left(\prod_{l=j+1}^n |\varphi_k(t_l r_l r_l^{-1})| \right) \|T_{i+1}(h_V^{\frac{2}{p_{i+1}}}) \dots T_{j-1}(h_V^{\frac{2}{p_{j-1}}}) (T_j(h_V^{\frac{2}{p_j}}) - \varphi_k(t_j r_j r_j^{-1}) h_V^{\frac{2}{q_j}}) \dots\|_{L_{\bar{p}_2}(\mathcal{L}G)} \\
& \leq \Delta((t_{i+1} \dots t_n)^{-1}) \|T_j(h_V^{\frac{2}{q_j}}) - \varphi_k(t_j r_j r_j^{-1}) h_V^{\frac{2}{q_j}}\|_{L_{q_j}(\mathcal{L}G)}.
\end{aligned}$$

We know that $q_j > \bar{p}_2 > 1$ for any $i+1 \leq j \leq n$. Additionally, T_j is bounded on $\mathcal{L}G$ and $L_1(\mathcal{L}G)$. By Proposition 4.3, the above terms converge to 0 in V . Hence, $\lim_{V \in \mathcal{V}} B_{k,V}^2 = 0$. This finishes the proof. \square

Remark 4.4. As in the unimodular case (see [6, Remark 3.3]) we do not know if Theorem 4.1 holds if $p = p_i = 1$ for some $1 \leq i \leq n$ (and $p_j = \infty$ for all $j \neq i$). The proof above fails in that case because we cannot apply Proposition 4.3.

5. Schur to Fourier transference for amenable groups

In this section, we extend [6, Proof of Theorem 4.1], that is, the transference from Schur multipliers to Fourier multipliers for amenable groups, to the non-unimodular setting. The proof is essentially the same but with extra technicalities due to the modular function. We also fixed a small mistake in the proof, as will be mentioned at the relevant spot.

Recall [19, Theorem 4.10] that G is amenable iff it satisfies the following Følner condition: for any $\varepsilon > 0$ and any compact set $K \subseteq G$, there exists a compact set $F \subseteq G$ with nonzero measure such that $\frac{\mu((sF \setminus F) \cup (F \setminus sF))}{\mu(F)} < \varepsilon$ for all $s \in K$. This allows us to construct a net $F_{(\varepsilon, K)}$ of such Følner sets using the ordering $(\varepsilon_1, K_1) \leq (\varepsilon_2, K_2)$ if $\varepsilon_1 \geq \varepsilon_2$, $K_1 \subseteq K_2$.

Theorem 5.1. Let G be a locally compact, amenable group and let $1 \leq p, p', p_1, \dots, p_n \leq \infty$ be such that $\frac{1}{p} = \sum_{i=1}^n \frac{1}{p_i} = 1 - \frac{1}{p'}$. Let $\phi \in L_\infty(G^{\times n})$ and define $\tilde{\phi} \in L_\infty(G^{\times n+1})$ by

$$\tilde{\phi}(s_0, \dots, s_n) = \phi(s_0 s_1^{-1}, s_1 s_2^{-1}, \dots, s_{n-1} s_n^{-1}), \quad s_i \in G.$$

Assume that $\tilde{\phi}$ defines a (p_1, \dots, p_n) -Schur multiplier of G . Then there is a net I , and there are complete contractions $i_{q,\alpha} : L_q(\mathcal{L}G) \rightarrow S_q(L_2(G))$, $\alpha \in I$, such that for all $f_i, f \in C_c(G) \star C_c(G)$,

$$\left| \langle i_{p,\alpha}(T_\phi(x_1, \dots, x_n)), i_{p',\alpha}(y) \rangle - \langle M_{\tilde{\phi}}(i_{p_1,\alpha}(x_1), \dots, i_{p_n,\alpha}(x_n)), i_{p',\alpha}(y) \rangle \right| \xrightarrow{\alpha} 0, \quad (5.1)$$

where $x_i = \Delta^{a_i} \lambda(f_i) \Delta^{b_i} \in L^{p_i}(\mathcal{L}G)$, $y = \Delta^a \lambda(f) \Delta^b \in L^{p'}(\mathcal{L}G)$ (i.e. $a_i + b_i = \frac{1}{p_i}$). In a similar way, the matrix amplifications of $i_{q,\alpha}$ approximately intertwine the multiplicative amplifications of the Fourier and Schur multipliers.

Proof. Let $F_\alpha, \alpha \in I$ be a Følner net for G , where I is the index set consisting of pairs (ε, K) for $\varepsilon > 0$, $K \subseteq G$ compact and the ordering as described above.

Let $P_\alpha = P_{F_\alpha}$ be the projection of $L_2(G)$ onto $L_2(F_\alpha)$. Consider the maps $i_{p,\alpha} : L_p(\mathcal{L}G) \rightarrow S_p(L_2(G))$ defined by $i_{p,\alpha}(x) = \mu(F_\alpha)^{-1/p} P_\alpha x P_\alpha$. They are contractions by [3, Theorem 5.1]. By replacing G by $G \times SU(2)$, one proves that they are in fact complete contractions (see also the last paragraph of [3, Proof of Theorem 5.2]).

Now fix α . From (2.10), we deduce

$$\begin{aligned} M_{\tilde{\phi}}(i_{p_1,\alpha}(x_1), \dots, i_{p_n,\alpha}(x_n))(t_0, t_n) \\ = \frac{1}{\mu(F_\alpha)^{1/p}} 1_{F_\alpha}(t_0) 1_{F_\alpha}(t_n) \int_{F_\alpha^{\times n-1}} \phi(t_0 t_1^{-1}, \dots, t_{n-1} t_n^{-1}) f_1(t_0 t_1^{-1}) \dots f_n(t_{n-1} t_n^{-1}) \times \\ \Delta^{a_1}(t_0) \Delta^{b_1+a_2}(t_1) \dots \Delta^{b_n}(t_n) \Delta((t_1 \dots t_n)^{-1}) dt_1 \dots dt_{n-1}. \end{aligned}$$

From Lemma 3.9, we have a similar expression for the kernel of $i_{p,\alpha}(T_\phi(x_1, \dots, x_n))$:

$$\begin{aligned} (t_0, t_n) \mapsto \frac{1}{\mu(F_\alpha)^{1/p}} 1_{F_\alpha}(t_0) 1_{F_\alpha}(t_n) \int_{G^{\times n-1}} \phi(t_0 t_1^{-1}, \dots, t_{n-1} t_n^{-1}) f_1(t_0 t_1^{-1}) \dots f_n(t_{n-1} t_n^{-1}) \times \\ \Delta^{a_1}(t_0) \Delta^{b_1+a_2}(t_1) \dots \Delta^{b_n}(t_n) \Delta((t_1 \dots t_n)^{-1}) dt_1 \dots dt_{n-1}. \end{aligned}$$

Now we need to take the pairing of these kernels with $i_{p',\alpha}(y)$ and calculate their difference. To that end, we define the following function Φ :

$$\begin{aligned} \Phi(t_0, \dots, t_n) = \phi(t_0 t_1^{-1}, \dots, t_{n-1} t_n^{-1}) f_1(t_0 t_1^{-1}) \dots f_n(t_{n-1} t_n^{-1}) f(t_0 t_1^{-1}) \times \\ \Delta^{a_1+b}(t_0) \Delta^{b_1+a_2}(t_1) \dots \Delta^{b_n+a}(t_n) \Delta((t_0 t_1 \dots t_n)^{-1}), \end{aligned}$$

and the function Ψ_α :

$$\Psi_\alpha(t_0, \dots, t_n) = 1_{F_\alpha}(t_0) 1_{F_\alpha}(t_n) - 1_{F_\alpha^{\times n+1}}(t_0, \dots, t_n) = 1_{F_\alpha \times (F_\alpha^{\times n-1})^c \times F_\alpha}(t_0, \dots, t_n).$$

Note that in [6], the indicator function was mistakenly taken over $F_\alpha \times (F_\alpha^c)^{\times n-1} \times F_\alpha$ instead. This correction leads to an extra term n in the choice of the lower bound of α at the end. Also note that *a priori*, it is not clear that $T_\phi(x_1, \dots, x_n)$ lies in $L_p(\mathcal{L}G)$, and hence it is not clear that $i_{p,\alpha}(T_\phi(x_1, \dots, x_n))$ lies in $S_p(L_2(G))$. However, both $i_{p,\alpha}(T_\phi(x_1, \dots, x_n))$ and $i_{p',\alpha}(y)$ are given by integration against a kernel in $L_2(G \times G)$, so the pairing (2.9) is still well-defined as a pairing in $S_2(L_2(G))$ instead. Now we have

$$\begin{aligned} \left| \langle i_{p,\alpha}(T_\phi(x_1, \dots, x_n)), i_{p',\alpha}(y) \rangle - \langle M_{\tilde{\phi}}(i_{p_1,\alpha}(x_1), \dots, i_{p_n,\alpha}(x_n)), i_{p',\alpha}(y) \rangle \right| \\ = \left| \frac{1}{\mu(F_\alpha)} \int_{G^{\times n+1}} \Phi(t_0, \dots, t_n) \Psi_\alpha(t_0, \dots, t_n) dt_0 \dots dt_n \right| \quad (5.2) \end{aligned}$$

Let $K \subseteq G$ be some compact set such that $\text{supp}(f_j), \text{supp}(f) \subseteq K$ and $e \in K$. Let t_0, \dots, t_n be such that both $\Phi(t_0, \dots, t_n)$ and $\Psi_\alpha(t_0, \dots, t_n)$ are nonzero. Since $\Psi_\alpha(t_0, \dots, t_n)$ is nonzero, we must have $t_0, t_n \in F_\alpha$

and $t_i \notin F_\alpha$ for some $i \in \{1, \dots, n-1\}$. Since $\Phi(t_0, \dots, t_n)$ is nonzero, there are $k_1, \dots, k_n \in K$ such that $t_{n-1} = k_n t_n$, $t_{n-2} = k_{n-1} k_n t_n$, \dots , $t_0 = k_1 \dots k_n t_n$. Hence we find

$$\begin{aligned} t_n &\in F_\alpha \cap (k_1 \dots k_n)^{-1} F_\alpha \setminus ((k_2 \dots k_n)^{-1} F_\alpha \cap \dots \cap k_n^{-1} F_\alpha) \\ &\subseteq F_\alpha \setminus ((k_2 \dots k_n)^{-1} F_\alpha \cap \dots \cap k_n^{-1} F_\alpha) \\ &= (F_\alpha \setminus (k_2 \dots k_n)^{-1} F_\alpha) \cup \dots \cup (F_\alpha \setminus k_n^{-1} F_\alpha) \end{aligned} \quad (5.3)$$

We want to apply a change of variables in (5.2). Let us first look at a simple case: assume $g \in L_1(G \times G)$ is such that $g(s, t) \neq 0$ only when $st^{-1} \in K$. Then

$$\begin{aligned} \int_{G \times 2} g(s, t) ds dt &= \int_{G \times 2} 1_K(st^{-1}) g(s, t) ds dt = \int_{G \times 2} 1_K(s) g(st, t) \Delta(t) ds dt \\ &= \int_G \int_K g(k_1 t, t) \Delta(t) dk_1 dt \end{aligned}$$

where we renamed the variable s in the last line. Applying the above formula twice for a function $g \in L_1(G^{\times 3})$ such that $g(r, s, t) \neq 0$ only when $rs^{-1} \in K$, $st^{-1} \in K$, we get

$$\begin{aligned} \int_{G \times 3} g(r, s, t) dr ds dt &= \int_{G \times 2} \int_K g(k_1 s, s, t) \Delta(s) dk_1 ds dt \\ &= \int_G \int_{K \times 2} g(k_1 k_2 t, k_2 t, t) \Delta(k_2 t) \Delta(t) dk_1 dk_2 dt. \end{aligned}$$

Carrying on like this, we obtain

$$\begin{aligned} &\left| \frac{1}{\mu(F_\alpha)} \int_{G^{\times n+1}} \Phi(t_0, \dots, t_n) \Psi_\alpha(t_0, \dots, t_n) dt_0 \dots dt_n \right| \\ &= \left| \frac{1}{\mu(F_\alpha)} \int_{K^{\times n}} \int_G \Phi(k_1 \dots k_n t_n, \dots, k_n t_n, t_n) \Psi_\alpha(k_1 \dots k_n t_n, \dots, k_n t_n, t_n) \times \right. \\ &\quad \left. \Delta(k_2 \dots k_n t_n) \dots \Delta(k_n t_n) \Delta(t_n) dt_n dk_1 \dots dk_n \right|. \end{aligned} \quad (5.4)$$

Note that $a + b + \sum_{i=1}^n a_i + b_i = 1$, hence

$$\begin{aligned} &|\Phi(k_1 \dots k_n t_n, \dots, k_n t_n, t_n)| \Delta(k_2 \dots k_n t_n) \dots \Delta(k_n t_n) \Delta(t_n) \\ &\leq \|\phi f_1 \dots f_n f\|_\infty \Delta^{a_1+b}(k_1 \dots k_n t_n) \Delta^{b_1+a_2+1}(k_2 \dots k_n t_n) \dots \Delta^{b_{n-1}+a_n+1}(k_n t_n) \times \\ &\quad \Delta^{b_n+a+1}(t_n) \Delta(k_1^{-1} k_2^{-2} \dots k_n^{-n} t_n^{-n-1}) \\ &= \|\phi f_1 \dots f_n f\|_\infty \Delta^{a_1+b-1}(k_1) \Delta^{a_1+a_2+b_1+b-1}(k_2) \dots \Delta^{1-b_n-a-1}(k_n) \\ &\leq \|\phi f_1 \dots f_n f\|_\infty C_{K,n} =: M. \end{aligned} \quad (5.5)$$

Here the constant $C_{K,n}$ can be chosen to be dependent only on K and n (and G).

Applying (5.2), (5.4), (5.5) and (5.3) consecutively, we get

$$\begin{aligned}
 & |\langle i_{p,\alpha}(T_\phi(x_1, \dots, x_n)), i_{p',\alpha}(y) \rangle_{p,p'} - \langle M_{\tilde{\phi}}(i_{p_1,\alpha}(x_1), \dots, i_{p_n,\alpha}(x_n)), i_{p',\alpha}(y) \rangle | \\
 &= \left| \frac{1}{\mu(F_\alpha)} \int_{K^{\times n}} \int_G \Phi(k_1 \dots k_n t_n, \dots, k_n t_n, t_n) \Psi_\alpha(k_1 \dots k_n t_n, \dots, k_n t_n, t_n) \times \right. \\
 &\quad \left. \Delta(k_2 \dots k_n t_n) \dots \Delta(k_n t_n) \Delta(t_n) dt_n dk_1 \dots dk_n \right| \\
 &\leq \frac{M}{\mu(F_\alpha)} \int_{K^{\times n}} \int_G \Psi_\alpha(k_1 \dots k_n t_n, \dots, k_n t_n, t_n) dt_n dk_1 \dots dk_n \\
 &= \frac{M}{\mu(F_\alpha)} \int_{K^{\times n}} \mu(F_\alpha \cap (k_1 \dots k_n)^{-1} F_\alpha \setminus ((k_2 \dots k_n)^{-1} F_\alpha \cap \dots \cap k_n^{-1} F_\alpha)) dk_1 \dots dk_n \\
 &\leq \frac{M}{\mu(F_\alpha)} \int_{K^{\times n}} \sum_{i=2}^n \mu(F_\alpha \setminus (k_i \dots k_n)^{-1} F_\alpha) dk_1 \dots dk_n \\
 &\leq M(n-1) \mu(K)^n \sup_{k \in K^{1-n}} \frac{\mu(F_\alpha \setminus k F_\alpha)}{\mu(F_\alpha)}. \tag{5.6}
 \end{aligned}$$

Using the ordering described earlier, if the index $\alpha \geq (\varepsilon \times (Mn\mu_G(K)^n)^{-1}, K^{1-n})$, then the Følner condition implies that (5.6) is less than ε , and hence the limit (5.1) holds.

From (5.1), it follows from writing out the definitions that the matrix amplifications of $i_{p,\alpha}$ also approximately intertwine the multiplicative amplifications of the Fourier and Schur multipliers; that is, for $\beta_i \in S_{p_i}^N$, $\beta \in S_{p'}^N$, we have

$$\begin{aligned}
 & \left| \langle \text{id} \otimes i_{p,\alpha}(T_\phi^{(N)}(\beta_1 \otimes x_1, \dots, \beta_n \otimes x_n)), \text{id} \otimes i_{p',\alpha}(\beta \otimes y) \rangle_{p,p'} \right. \\
 & \quad \left. - \langle M_{\tilde{\phi}}^{(N)}(\text{id} \otimes i_{p_1,\alpha}(\beta_1 \otimes x_1), \dots, \text{id} \otimes i_{p_n,\alpha}(\beta_n \otimes x_n)), \text{id} \otimes i_{p',\alpha}(\beta \otimes y) \rangle \right| \rightarrow 0 \tag{5.7}
 \end{aligned}$$

□

Corollary 5.2. *Let G be an amenable locally compact group and $1 \leq p, p_1, \dots, p_n \leq \infty$ be such that $\frac{1}{p} = \sum_{i=1}^n \frac{1}{p_i}$. Let $\phi \in L_\infty(G^{\times n})$. If $\tilde{\phi}$ is the symbol of a (p_1, \dots, p_n) -bounded (resp. multiplicatively bounded) Schur multiplier then ϕ is the symbol of a (p_1, \dots, p_n) -bounded (resp. multiplicatively bounded) Fourier multiplier. Moreover,*

$$\|T_\phi\|_{(p_1, \dots, p_n)} \leq \|M_{\tilde{\phi}}\|_{(p_1, \dots, p_n)}, \quad \|T_\phi\|_{(p_1, \dots, p_n)-mb} \leq \|M_{\tilde{\phi}}\|_{(p_1, \dots, p_n)-mb}.$$

Proof. Let x_i be as in the hypotheses of Theorem 5.1, and let $i_{p,\alpha}$ be as in the proof of Theorem 5.1. Let p' be the Hölder conjugate of p . In [3, Theorem 5.2], it is proven that

$$\langle i_{p,\alpha}(x), i_{p',\alpha}(y) \rangle_{p,p'} \rightarrow \langle x, y \rangle_{p,p'}, \quad x \in L_p(\mathcal{L}G), \quad y \in L_{p'}(\mathcal{L}G). \tag{5.8}$$

Note that this inequality also holds and in fact is explicitly proven for $p = \infty$; by symmetry, it also holds for $p = 1$. We remark that this result also uses the Følner condition.

Let $\varepsilon > 0$. Then we can find $y = \Delta^a \lambda(f) \Delta^b$, $f \in C_c(G) \star C_c(G)$ such that $\|y\|_{L_{p'}(\mathcal{L}G)} \leq 1$ and

$$\|T_\phi(x_1, \dots, x_n)\|_{L_p(\mathcal{L}G)} \leq |\langle T_\phi(x_1, \dots, x_n), y \rangle| + \varepsilon.$$

Next, by (5.8) and Theorem 5.1, we can find $\alpha \in I$ such that the following two inequalities hold:

$$|\langle T_\phi(x_1, \dots, x_n), y \rangle - \langle i_{p,\alpha}(T_\phi(x_1, \dots, x_n)), i_{p',\alpha}(y) \rangle| < \varepsilon$$

and

$$|\langle i_{p,\alpha}(T_\phi(x_1, \dots, x_n)), i_{p',\alpha}(y) \rangle_{p,p'} - \langle M_{\tilde{\phi}}(i_{p_1,\alpha}(x_1), \dots, i_{p_n,\alpha}(x_n)), i_{p',\alpha}(y) \rangle_{p,p'} | < \varepsilon.$$

By combining these inequalities, we find

$$\begin{aligned} \|T_\phi(x_1, \dots, x_n)\|_{L_p(\mathcal{L}G)} &\leq |\langle M_{\tilde{\phi}}(i_{p_1, \alpha}(x_1), \dots, i_{p_n, \alpha}(x_n)), i_{p', \alpha}(y) \rangle_{p, p'}| + 3\varepsilon \\ &\leq \|M_{\tilde{\phi}}\|_{(p_1, \dots, p_n)} \prod_{i=1}^n \|x_i\|_{L_{p_i}(\mathcal{L}G)} + 3\varepsilon \end{aligned}$$

The elements x_i as chosen above are norm dense in $L_{p_i}(\mathcal{L}G)$ (resp. $C_\lambda^*(G)$ when $p_i = \infty$); hence, we get the required bound. The multiplicative bound follows similarly. \square

Remark 5.3. In [3], [6], the proof runs via an ultraproduct construction. The ultraproduct is not actually necessary as demonstrated above, as all limits are usual limits and not ultralimits.

We can now extend the result of [6, Corollary 4.3] to non-unimodular groups. It is also a multiplicatively bounded, non-unimodular version of [5, Theorem 4.5]. Moreover, we no longer need the SAIN condition, and the subgroup no longer needs to be discrete.

Corollary 5.4. Let G be a locally compact, first countable group, and let $1 \leq p \leq \infty$ and $1 < p_1, \dots, p_n \leq \infty$ with $p^{-1} = \sum_{i=1}^n p_i^{-1}$. Let $\phi \in C_b(G^{\times n})$, which defines a (p_1, \dots, p_n) -mb Fourier multiplier, and let $H \leq G$ be an amenable subgroup. Then

$$\|T_\phi\|_{H^{\times n} \rightarrow H} \|(p_1, \dots, p_n)\text{-mb}\| \leq \|T_\phi\|_{(p_1, \dots, p_n)\text{-mb}}$$

Proof. The associated inequality for Schur multipliers follows from Theorem 2.2. Now Corollary 5.2 (using amenability of H) and Theorem 4.1 yield the result. \square

In the next corollary, we prove a necessary condition for a ‘Fourier multiplier’ to satisfy (5.1) for the embeddings $i_{p, \alpha}$ defined above. This was used in the discussion in Section 3.

Corollary 5.5. Fix $n > 1$, $1 \leq p_1, \dots, p_n, p, p' \leq \infty$ such that $\frac{1}{p} = \sum_{i=1}^n \frac{1}{p_i} = 1 - \frac{1}{p'}$, and let $\theta_1, \dots, \theta_n, \theta, \theta' \in [0, 1]$. Let $i_{p, \alpha}$ be as in the proof of Theorem 5.1. Assume that for each $\phi \in L_\infty(G^{\times n})$, we have a map $S_\phi : \kappa_{p_1}^{\theta_1}(L) \times \dots \times \kappa_{p_n}^{\theta_n}(L) \rightarrow \kappa_p^\theta(L)$ satisfying

$$|\langle i_{p, \alpha}(S_\phi(x_1, \dots, x_n)), i_{p', \alpha}(y) \rangle_{p, p'} - \langle M_{\tilde{\phi}}(i_{p_1, \alpha}(x_1), \dots, i_{p_n, \alpha}(x_n)), i_{p', \alpha}(y) \rangle_{p, p'}| \xrightarrow{\alpha} 0. \quad (5.9)$$

for $x_i \in \kappa_{p_i}^{\theta_i}(L)$, $y \in \kappa_{p'}^{\theta'}(L)$. Now let $\phi(s_1, \dots, s_n) = \phi_1(s_1) \dots \phi_n(s_n)$ for some functions $\phi_1, \dots, \phi_n \in L^\infty(G)$. Then S_ϕ must satisfy

$$S_\phi(x_1, \dots, x_n) = T_{\phi_1}(x_1) \dots T_{\phi_n}(x_n)$$

for $x_i \in \kappa_{p_i}^{\theta_i}(L)$, $i = 1, \dots, n$.

Proof. Fix some $y \in \kappa_{p'}^{\theta'}(L)$. By density, it suffices to show that

$$\langle S_\phi(x_1, \dots, x_n), y \rangle = \langle T_{\phi_1}(x_1) \dots T_{\phi_n}(x_n), y \rangle.$$

By (5.8), it suffices to show

$$\lim_{\alpha \in I} |\langle i_{p, \alpha}(S_\phi(x_1, \dots, x_n) - T_{\phi_1}(x_1) \dots T_{\phi_n}(x_n)), i_{p', \alpha}(y) \rangle| = 0.$$

By running the proof of Theorem 5.1 with the constant 1 function in place of ϕ and $T_{\phi_i}(x_i)$ in place of x_i , we find that

$$\lim_{\alpha \in I} |\langle i_{p, \alpha}(T_{\phi_1}(x_1) \dots T_{\phi_n}(x_n) - i_{p_1, \alpha}(T_{\phi_1}(x_1)) \dots i_{p_n, \alpha}(T_{\phi_n}(x_n))), i_{p', \alpha}(y) \rangle| = 0.$$

Since multiplication with ϕ_i only maps $C_c(G) \star C_c(G)$ to $C_c(G)$, we no longer need to have that $T_{\phi_i}(x_i) \in \kappa_{p_i}^{\theta_i}(L)$, so we cannot apply Theorem 5.1 directly. But since we have $\phi = 1$, this does not give any technical complications in the proof.

Using the kernel representations, it is straightforward to show that

$$\begin{aligned} i_{p_1, \alpha}(T_{\phi_1}(x_1)) \dots i_{p_n, \alpha}(T_{\phi_n}(x_n)) &= M_{\widehat{\phi}_1}(i_{p_1, \alpha}(x_1)) \dots M_{\widehat{\phi}_n}(i_{p_n, \alpha}(x_n)) \\ &= M_{\widehat{\phi}}(i_{p_1, \alpha}(x_1), \dots, i_{p_n, \alpha}(x_n)). \end{aligned}$$

By combining the above observations with (5.9), we get the required result. \square

6. Linear intertwining result

In this section, we sketch the proof of Proposition 4.3. The main ingredient to be added to already existing results is the extension of [9, Lemma 3.1] to general von Neumann algebras via Haagerup reduction. The Haagerup reduction method is described by Theorem 6.1, proved for σ -finite von Neumann algebras in [12] and extended to the weight case in [8, Section 8]. We will assume that the reader is familiar with Tomita–Takesaki theory, conditional expectations and such. We refer to [24] for the background.

Denote by σ^φ the modular automorphism group of a normal faithful semifinite (nfs) weight φ . Recall that the centraliser \mathcal{N}_φ of a nfs weight φ on a von Neumann algebra \mathcal{M} is given by

$$\mathcal{N}_\varphi = \{x \in \mathcal{M} : \sigma_t^\varphi(x) = x \ \forall t \in \mathbb{R}\}.$$

Theorem 6.1. *Let (\mathcal{M}, φ) be any von Neumann algebra equipped with a nfs weight. There is another von Neumann algebra $(\mathcal{R}, \widehat{\varphi})$ containing \mathcal{M} and with nfs weight $\widehat{\varphi}$ extending φ and elements a_n in the center of the centraliser of $\widehat{\varphi}$ such that the following properties hold:*

1. *There is a conditional expectation $\mathcal{E} : \mathcal{R} \rightarrow \mathcal{M}$ satisfying*

$$\varphi \circ \mathcal{E} = \widehat{\varphi}, \quad \sigma_s^\varphi \circ \mathcal{E} = \mathcal{E} \circ \sigma_s^{\widehat{\varphi}}, \quad s \in \mathbb{R}.$$

2. *The centralisers \mathcal{R}_n of the weights $\varphi_n := \varphi(e^{-a_n} \cdot)$ are semifinite for $n \geq 1$.*
3. *There are conditional expectations $\mathcal{E}_n : \mathcal{R} \rightarrow \mathcal{R}_n$ satisfying*

$$\widehat{\varphi} \circ \mathcal{E}_n = \widehat{\varphi}, \quad \sigma_s^{\widehat{\varphi}} \circ \mathcal{E}_n = \mathcal{E}_n \circ \sigma_s^{\widehat{\varphi}}, \quad s \in \mathbb{R}$$

4. *$\mathcal{E}_n(x) \rightarrow x$ σ -strongly for $x \in \mathfrak{n}_{\widehat{\varphi}}$, and $\bigcup_{n \geq 1} \mathcal{R}_n$ is σ -strongly dense in \mathcal{R} .*

We denote by D_φ the spatial derivative with respect to φ (and some weight on the commutant, whose choice is unimportant). Assume that $T : \mathcal{M} \rightarrow \mathcal{M}$ is unital completely positive (ucp) and satisfies $\varphi \circ T \leq \varphi$. Then by [12, Section 5], T ‘extends’ to a map $T^{(\varphi)}$ on $L_p(\mathcal{M})$, in the sense that $T^{(\varphi)}(D_\varphi^{1/2p} x D_\varphi^{1/2p}) = D_\varphi^{1/2p} T(x) D_\varphi^{1/2p}$ for $x \in \mathfrak{m}_{\widehat{\varphi}}$. If T satisfies $\sigma_s^\varphi \circ T = T \circ \sigma_s^\varphi$, then we moreover have $T^{(\varphi)}(D_\varphi^{\theta/p} x D_\varphi^{(1-\theta)/p}) = D_\varphi^{\theta/p} T(x) D_\varphi^{(1-\theta)/p}$ for any $0 \leq \theta \leq 1$ and $x \in \mathfrak{m}_{\widehat{\varphi}}$.

In particular, the conditional expectations $\mathcal{E}, \mathcal{E}_n$ ‘extend’ to maps $\mathcal{E}^{(\varphi)}, \mathcal{E}_n^{(\varphi)}$ from $L_p(\mathcal{R}, \widehat{\varphi})$ to $L_p(\mathcal{M}, \varphi)$ resp. $L_p(\mathcal{R}_n, \widehat{\varphi})$. The following statement is [8, Lemma 8.3]:

$$\lim_{n \rightarrow \infty} \|\mathcal{E}_n^{(\varphi)}(x) - x\|_p = 0, \quad 1 \leq p < \infty, \ x \in L_p(\mathcal{R}, \widehat{\varphi}). \quad (6.1)$$

We need a few more facts; we refer to [8, Section 8.2] for the details. First, there is an isometric isomorphism $\kappa_p : L_p(\mathcal{R}_n, \widehat{\varphi}) \rightarrow L_p(\mathcal{R}_n, \varphi_n)$ given by $\kappa_p(D_\varphi^{1/2p} x D_\varphi^{1/2p}) = e^{a_n/2p} x e^{a_n/2p}$ for $x \in \mathfrak{m}_{\widehat{\varphi}}$. Next, assume that $T : \mathcal{M} \rightarrow \mathcal{M}$ is ucp and preserves φ and σ_s^φ . Then by [12, Section 4], there exists an extension $\widehat{T} : \mathcal{R} \rightarrow \mathcal{R}$, which is also ucp and preserves $\widehat{\varphi}$ and $\sigma_s^{\widehat{\varphi}}$. Hence \widehat{T} itself also ‘extends’ to the various noncommutative L_p -spaces. Moreover, the following diagram commutes:

$$\begin{array}{ccccc} L_p(\mathcal{R}, \widehat{\varphi}) & \xrightarrow{\mathcal{E}_n^{(p)}} & L_p(\mathcal{R}_n, \widehat{\varphi}) & \xrightarrow{\kappa_p} & L_p(\mathcal{R}_n, \varphi_n) \\ \widehat{T}^{(p)} \uparrow & & \widehat{T}^{(p)} \uparrow & & \widehat{T} \uparrow \\ L_p(\mathcal{R}, \widehat{\varphi}) & \xrightarrow{\mathcal{E}_n^{(p)}} & L_p(\mathcal{R}_n, \widehat{\varphi}) & \xrightarrow{\kappa_p} & L_p(\mathcal{R}_n, \varphi_n). \end{array}$$

Note that since φ_n is tracial on \mathcal{R}_n , the \widehat{T} in the rightmost upwards arrow is actually an extension of the operator \widehat{T} on \mathcal{R}_n , so we do not need to use the notation $\widehat{T}^{(p)}$ here.

Finally, for $1 \leq p, q < \infty$, we define the Mazur maps $M_{p,q} : L_p(\mathcal{M}) \rightarrow L_q(\mathcal{M})$ by $x \mapsto u|x|^{p/q}$, where $x = u|x|$ is the polar decomposition of x . The Mazur maps satisfy $\kappa_q \circ M_{p,q} = M_{p,q} \circ \kappa_p$; see, for instance, [23, end of Section 3]. We are now ready to state and prove the generalisation of [9, Lemma 3.1] for general von Neumann algebras. This result was already shown for $2 < p < \infty$ in [8, Section 8], but we will prove the result for all $1 < p < \infty$ at once since this does not take any extra effort.

Lemma 6.2. *Let (\mathcal{M}, φ) be a von Neumann algebra equipped with nfs weight. Let $T : \mathcal{M} \rightarrow \mathcal{M}$ be a unital completely positive map satisfying $\varphi \circ T = \varphi$ and $T \circ \sigma_s^\varphi = \sigma_s^\varphi \circ T$ for all $s \in \mathbb{R}$. Then there exists a universal constant $C > 0$ such that for any $x \in L_2(\mathcal{M})$ and $1 < p < \infty$:*

$$\|T^{(p)}(M_{2,p}(x)) - M_{2,p}(x)\|_p \leq C \|T^{(2)}(x) - x\|_2^\theta \|x\|_2^{1-\theta},$$

where $\theta = \frac{1}{4} \min\{\frac{p}{2}, \frac{2}{p}\}$.

Proof. The proof runs via Haagerup reduction, using the estimates for the semifinite case from [8, Claim B] for $p > 2$ and [9, Lemma 3.1] for $p < 2$. Note that the latter was stated only for finite von Neumann algebras, but the same proof works for the semifinite case as well.

Set $y = M_{2,p}(x)$. Since $T = \widehat{T}$ on \mathcal{M} and $L_p(\mathcal{M}, \varphi) \hookrightarrow L_p(\mathcal{R}, \widehat{\varphi})$ canonically and isometrically, we have $T^{(p)}(y) = \widehat{T}^{(p)}(y)$ and

$$\|T^{(p)}(y) - y\|_{L_p(\mathcal{M}, \varphi)} = \|\widehat{T}^{(p)}(y) - y\|_{L_p(\mathcal{R}, \widehat{\varphi})}.$$

Now fix $n \geq 1$. Then

$$\begin{aligned} \|\mathcal{E}_n^{(p)}(\widehat{T}^{(p)}(y)) - \mathcal{E}_n^{(p)}(y)\|_{L_p(\mathcal{R}_n, \widehat{\varphi})} &= \|\kappa_p(\mathcal{E}_n^{(p)}(\widehat{T}^{(p)}(y)) - \mathcal{E}_n^{(p)}(y))\|_{L_p(\mathcal{R}_n, \varphi_n)} \\ &= \|\widehat{T}(\kappa_p(\mathcal{E}_n^{(p)}(y))) - \kappa_p(\mathcal{E}_n^{(p)}(y))\|_{L_p(\mathcal{R}_n, \varphi_n)}. \end{aligned}$$

Now we can apply the result for the semifinite case on $\kappa_p(\mathcal{E}_n^{(p)}(y))$ to obtain

$$\begin{aligned} &\|\mathcal{E}_n^{(p)}(\widehat{T}^{(p)}(y)) - \mathcal{E}_n^{(p)}(y)\|_{L_p(\mathcal{R}_n, \widehat{\varphi})} \\ &\leq C \|\widehat{T}(M_{p,2}(\kappa_p(\mathcal{E}_n^{(p)}(y)))) - M_{p,2}(\kappa_p(\mathcal{E}_n^{(p)}(y)))\|_{L_2(\mathcal{R}_n, \varphi_n)}^\theta \cdot \|M_{p,2}(\kappa_p(\mathcal{E}_n^{(p)}(y)))\|_{L_2(\mathcal{R}_n, \varphi_n)}^{1-\theta} \\ &= C \|\kappa_2(\widehat{T}^{(2)}(M_{p,2}(\mathcal{E}_n^{(p)}(y)))) - \kappa_2(M_{p,2}(\mathcal{E}_n^{(p)}(y)))\|_{L_2(\mathcal{R}_n, \varphi_n)}^\theta \cdot \|\kappa_2(M_{p,2}(\mathcal{E}_n^{(p)}(y)))\|_{L_2(\mathcal{R}_n, \varphi_n)}^{1-\theta} \\ &= C \|\widehat{T}^{(2)}(M_{p,2}(\mathcal{E}_n^{(p)}(y))) - M_{p,2}(\mathcal{E}_n^{(p)}(y))\|_{L_2(\mathcal{R}_n, \widehat{\varphi})}^\theta \cdot \|M_{p,2}(\mathcal{E}_n^{(p)}(y))\|_{L_2(\mathcal{R}_n, \widehat{\varphi})}^{1-\theta} \\ &=: CA_n^\theta B_n^{1-\theta}. \end{aligned}$$

By the triangle inequality, the main result from [23] and (6.1), we find

$$\begin{aligned} B_n &\leq \|M_{p,2}(\mathcal{E}_n^{(p)}(y)) - M_{p,2}(y)\|_{L_2(\mathcal{R}_n, \widehat{\varphi})} + \|M_{p,2}(y)\|_{L_2(\mathcal{R}_n, \widehat{\varphi})} \\ &\leq C_{x,p} \|\mathcal{E}_n^{(p)}(y) - y\|_{L_p(\mathcal{R}_n, \widehat{\varphi})}^{\min\{\frac{p}{2}, 1\}} + \|x\|_{L_2(\mathcal{R}_n, \widehat{\varphi})} \rightarrow \|x\|_{L_2(\mathcal{M}, \varphi)} \end{aligned}$$

for some constant $C_{x,p}$ independent of n . Similarly, we find

$$A_n \leq C_{x,p} \|\widehat{T}^{(2)} - 1_{\mathcal{R}}\| \|\mathcal{E}_n^{(p)}(y) - y\|_{L_p(\mathcal{R}_n, \widehat{\varphi})}^{\min\{\frac{p}{2}, 1\}} + \|\widehat{T}^{(2)}(x) - x\|_{L_2(\mathcal{R}_n, \widehat{\varphi})} \rightarrow \|T^{(2)}(x) - x\|_{L_2(\mathcal{M}, \varphi)}.$$

Hence, taking limits and applying again (6.1), we conclude

$$\|T^{(p)}(y) - y\|_p = \lim_{n \rightarrow \infty} \|\mathcal{E}_n^{(p)}(\widehat{T}^{(p)}(y)) - \mathcal{E}_n^{(p)}(y)\|_{L_p(\mathcal{R}_n, \widehat{\varphi})} \leq C \|T^{(2)}(x) - x\|_2^\theta \|x\|_2^{1-\theta}.$$

□

Proof of Proposition 4.3. We indicate only the changes to [8, Proof of Claim B]. The statement we have to prove is precisely [8, Equation (9)], but without the u_j (this is just a different choice based on convenience). The T_ζ constructed in [8, Proof of Claim B] is a φ -preserving ucp map that commutes with the modular automorphism group; one can see this from (2.4). Hence, we can apply Lemma 6.2 on T_ζ and h_V to show [8, Equation (10)] (but without the u_j). Then, setting $z_j = h_V^{2/q}$, the rest of the proof is the same. □

Acknowledgement. The author thanks Martijn Caspers for useful discussions and a thorough proofreading of the manuscript.

Competing interests. The author declares none.

Funding statement. GV is supported by the NWO Vidi grant VI.Vidi.192.018 ‘Non-commutative harmonic analysis and rigidity of operator algebras’.

References

- [1] M. Bożejko and G. Fendler, Herz-Schur multipliers and completely bounded multipliers of the Fourier algebra of a locally compact group (English, with Italian summary), *Boll. Un. Mat. Ital. A.* **6**(3) (1984), 297–302.
- [2] M. Caspers, The L_p -fourier transform on locally compact quantum groups, *J. Operator Theory* **69**(1) (2013), 161–193.
- [3] M. Caspers and M. de la Salle, Schur and fourier multipliers of an amenable group acting on non-commutative L_p -spaces, *Trans. Amer. Math. Soc.* **367**(10) (2015), 6997–7013.
- [4] J.M. Conde-Alonso, A.M. González-Pérez, J. Parcet and E. Tablate, Schur multipliers in Schatten–von Neumann classes, *Ann. Math.* **198**(3) (2023), 1229–1260.
- [5] M. Caspers, B. Janssens, A. Krishnaswamy-Usha and L. Miaskiwsy, Local and multilinear noncommutative De Leeuw theorems, *Math. Ann.* **388**(4) (2024), 4251–4305.
- [6] M. Caspers, A. Krishnaswamy-Usha and G. Vos, Multilinear transference of Fourier and Schur multipliers acting on non-commutative L_p -spaces, *Canadian J Math* **75**(6) (2022), 1–18. DOI: [10.4153/S0008414X2200058X](https://doi.org/10.4153/S0008414X2200058X)
- [7] A. Connes, On the spatial theory of von Neumann algebras, *J. Funct. Anal.* **35**(2) (1980), 153–164.
- [8] M. Caspers, J. Parcet, M. Perrin and É. Ricard, Noncommutative de Leeuw theorems, *Forum Math. Sigma, Paper* **3**(e21) (2015), 59.
- [9] J. Conde-Alonso, J. Parcet and É. Ricard, On spectral gaps of Markov maps, *Israel J. Math.* **226**(1) (2018), 189–203.
- [10] A. Gonzalez-Perez, Crossed-products extensions, of L_p -bounds for amenable actions, *J. Functional Analysis* **274**(10) (2018), 2846–2883.
- [11] M. Hilsen, Les espaces L^p d’une algèbre de von Neumann définies par la dérivée spatiale, *J. Funct. Anal.* **40**(2) (1981), 151–169.
- [12] U. Haagerup, M. Junge and Q. Xu, A reduction method for noncommutative. L_p -spaces and applications, *Trans. Amer. Math. Soc.* **362**(4) (2010), 2125–2165.
- [13] P. Jolissaint, A characterization of completely bounded multipliers of fourier algebras, *Colloq. Math.* **63**(2) (1992), 311–313.
- [14] K. Juschenko, I. G. Todorov and L. Turowska, Multidimensional operator multipliers, *Trans. Am. Math. Soc.* **361**(9) (2009), 4683–4720.
- [15] M. Junge and D. Sherman, Noncommutative L^p modules, *J. Operator Theory* **53**(1) (2005), 3–34.
- [16] E. Kaniuth and A. Lau, *Fourier and fourier-stieltjes algebras on locally compact groups*, Mathematical surveys and Monographs (American Mathematical Society, Providence, RI, 2018), xi+306 pp.
- [17] V. Lafforgue and M. de la Salle, Noncommutative L_p -spaces without the completely bounded approximation property, *Duke Math. J.* **160**(1) (2011), 71–116.
- [18] S. Neuirth and É. Ricard, Transfer of Fourier multipliers into Schur multipliers and sumsets in a discrete group, *Canad. J. Math.* **63**(5) (2011), 1161–1187.
- [19] A.L.T. Paterson and Amenability, *Mathematical surveys and Monographs* 29 (American Mathematical Society, Providence, RI, 1988), xx+452 pp.
- [20] G. Pisier, Non-commutative vector valued. L_p -spaces and completely p -summing maps, *Astérisque* **247** (1998), vi+131

- [21] J. Parcet, É. Ricard and M. de la Salle, Fourier multipliers in $\mathrm{SLn}(\mathbb{R})$, *Duke Math. J.* **171**(6) (2022), 1235–1297.
- [22] D. Potapov, A. Skripka and F. Sukochev, Spectral shift function of higher order, *Invent. Math.* **193**(3) (2013), 501–538.
- [23] É. Ricard, Hölder estimates for the noncommutative Mazur maps, *Arch. Math. (Basel)* **104**(1) (2015), 37–45.
- [24] M. Takesaki, Theory of operator algebras. II, in *Encyclopaedia of Mathematical Sciences*, vol. 125, Operator Algebras and Noncommutative Geometry (Springer-Verlag, Berlin, 2003, xxii+518 pp)
- [25] M. Terp, L_p spaces associated with von Neumann algebras, *Notes*. Københavns Universitets Matematiske Institut, Juni 1981, 3a + 3b
- [26] I. G. Todorov and L. Turowska *multipliers of multidimensional Fourier algebras*, *Oper. Matrices.* **4**(4) (2010), 459–484.
- [27] G. Vos, Multipliers and transference on noncommutative L_p -spaces, *PhD Thesis* (TU Delft repository, 2024).