

defined. Boys were more likely to be rated screenpositive than girls. Share of boys rated in the top 5% by parents is 5.6% compared to 4.3% of girls. Teachers place boys in the top 5% even more frequently – 6.4% versus 3.4% girls. Pairwise correlation coefficients (0.53) revealed moderate correlations between scores and according to p-values (< 0.05) all correlations were statistically significant.

**Conclusions:** The study defined the cut-off scores of ASSQ for 8-10 y old Georgian children and gender difference in prevalence of risk for ASD. Using the ASSQ was an effective instrument and could be used in school settings to identify children with special needs. \*This work was supported by Shota Rustaveli National Science Foundation of Georgia (SRNSFG), grant - FR-18-304.

**Disclosure:** No significant relationships.

**Keywords:** ASSQ; Prevalence of autism in Georgia; autism

## EPV0081

### Quality of life in neurotypical siblings of children with an autism spectrum disorder: Detecting children at risk

E. Koukouriki<sup>1,2</sup>

<sup>1</sup>Special Education Lab, Department Of Primary Education, University of Ioannina, Ioannina, Greece and <sup>2</sup>Centre For Educational And Counseling Services Of Trikala, Hellenic Ministry of Education and Religious Affairs, Trikala, Greece  
doi: 10.1192/j.eurpsy.2021.1689

**Introduction:** Quality of life (QOL) instruments can provide unique information on various aspects of children's everyday functioning beyond clinical symptoms and clinical measures. The study of the QOL in the siblings of individuals with an Autism Spectrum Disorder (ASD-siblings) is a new direction in the field of ASD-research.

**Objectives:** This study aimed to investigate the QOL in ASD-siblings in comparison with an age and sex matched group of siblings of neurotypical children.

**Methods:** The sample included 233 neurotypical children (8-13 years old) of whom 118 comprised the observational group (ASD-siblings) and 115 comprised the comparison group. The Kidscreen-27 and a demographics questionnaire were administered to all participants.

**Results:** The two groups differed significantly in all subscales and in the overall score of the Kidscreen-27, with the children in the observation group having a significantly worse quality of life (Mean±sd 111,62±12,43, p<0,001). Additionally, the largest difference in the averages of the subscales occurs in the subscale General Mood and Emotions, where the observation group had significantly lower scores (Mean±sd 23,23±3,69, p<0,001) than the comparison group (Mean±sd 31,27±2,74).

**Conclusions:** QOL in children has been recognized as an important outcome indicator in detecting subgroups of children at risk within the general population, while ASD-siblings' QOL is an important concept in the implementation of appropriate services for these children. This study revealed poorer QOL in ASD-siblings and highlighted the importance of assessing QOL in those siblings as well as the use of the Kidscreen-27 as a screening tool in order to detect children at risk of maladjustment.

**Disclosure:** No significant relationships.

**Keywords:** Quality of life; siblings; autism spectrum disorders

## EPV0082

### Estimation of prevalence of mental health problems in 8-10 years old georgian children by using the strengths and difficulties questionnaire\*

G. Chvamanian<sup>1,2\*</sup>, M. Zirakashvili<sup>1,3</sup>, T. Mikiashvili<sup>1,4</sup>, N. Mebonia<sup>1,5</sup> and M. Gabunia<sup>1,6</sup>

<sup>1</sup>Mental Health, Georgian Academy of Childhood Disability, Tbilisi, Georgia; <sup>2</sup>Psychology, Ilia State University, Tbilisi, Georgia; <sup>3</sup>Medical Sciences, Ilia State University, Tbilisi, Georgia; <sup>4</sup>Psychology, Javakishvili Tbilisi State University, Tbilisi, Georgia; <sup>5</sup>Department Of Chronic Deaseases, National Center for Disease Control and Public Health, Tbilisi, Georgia and <sup>6</sup>Child And Adolescent Mental Health, Mental Health Center, Tbilisi, Georgia

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1690

**Introduction:** Mental health problems are frequent among children and seems to predict mental disorders in adulthood.

**Objectives:** The study aimed whether the gender differences affects the Strengths and Difficulties Questionnaire (SDQ) assessments performed by parents and teachers in Republic of Georgia.

**Methods:** In 2019 a cross sectional survey in four main cities of Georgia was conducted; Totally 8-10 y old 16654 children from 211 public schools were included. SDQ completed by parents and school teachers was used to determine emotional and behavioral problems among Georgian children.

**Results:** 16654 (74%) parents out of 22553 were agreed to participate in the study. 1565 (9.39%) children were rated screen positive in top five percentile by either parent or teacher or both of them. Cut-off scores for 99-95 percentiles (top 1-5%) was defined. Boys were more likely to be rated screen positive than girls, especially by teachers: parents rated screen positive 7.5% of females, teachers - 7.2%, while males 9.4% and 11.5% respectively. Pairwise correlation coefficients (0.53) revealed moderate correlations according to p-values (< 0.05) between scores and all correlations were statistically significant.

**Conclusions:** The study defined the cut-off scores of SDQ for 8-10 y old children and a gender differences in prevalence of mental health problems in Georgia. SDQ could be used in primary healthcare and school settings to identify children with special needs. This work was supported by Shota Rustaveli National Science Foundation of Georgia (SRNSFG), grant - FR-18-304.

**Disclosure:** No significant relationships.

**Keywords:** SDQ; Prevalence in Georgia; Children Mental Health

## EPV0083

### Street working children in kurdistan region of Iraq; Mental health and traumatization

N. Taib\*, H. Arinell, A. Ahmad and M. Ramklint

Neuroscience/child And Adolescent Psychiatry, Uppsala University, Uppsala, Sweden

\*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1691

**Introduction:** Street working children are often poor, deprived of love and care, and lack supervision by responsible adults. The Kurdistan region of Iraq has experienced war conflicts for decades.