Vol. LXXXVI No. 9

The Journal of Laryngology and Otology

EDITED BY
SIR GEOFFREY BATEMAN

ASSISTANT EDITOR

Contents

Otogenic meningitis: A five-year study	B. BRYDØY and E. F. ELLEKJÆR	871
✓ Grommets and glue ears: Two-year results	D. KILBY, S. H. RICHARDS and G. HART	881
✓ The surgical treatment of hemi-facial spasm	J. POTTER	889
Lecological alterations of oral microflora subsequent	Dayman W. Vormer	
to tonsillectomy and adenoidectomy	ROBERT W. VELTRI, PHILIP M. SPRINKLE,	
	SUE ANN KELLER	
	and James M. Chicklo	893-
Immunoglobulin changes in a pediatric otolaryngic patient sample subsequent to T & A	ROBERT W. VELTRI,	
patient sample subsequent to 1 te A	PHILIP M. SPRINKLE,	
	SUE ANN KELLER	
	and JAMES M. CHICKLO	905
✓ A comparative study of the actiology of vertigo in the African	CHRISTIAN NSAMBA	917-
Clinical records—		
✓ Acute epiglottitis in adults	J. L. HANDA	927
Localized amyloidosis of the larynx	Francis D'Arcy	929
Basal cell carcinoma of the nose: with report of two		
atypical cases	HUSSEIN H. TOPPOZADA	
	and Mostafa A. Shehata	933
Melkersson's syndrome: an unusual case	DONALD B. HAWKINS	943
Protection against blast injuries of the ear	DARYAO SINGH and I. J. MARHIJA	040
JGeneral notes	and I. J. MARITA	949
		2000

Headley Brothers Ltd

Ashford, Kent

The Journal of Laryngology and Otology

(FOUNDED in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY SIR GEOFFREY BATEMAN

ASSISTANT EDITOR
LIONEL TAYLOR

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this *Journal* solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgement must be made.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. Green, C., and Brown, D. (1951), Journal of Laryngology and Otology, 65, 33. The titles of all Journals should be given without abbreviation.

It is most important that authors should verify personally the accuracy of every reference before submitting a paper for publication.

- 3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.
- 4. Illustration blocks will normally be held by the Printers for three years, after which they will be destroyed. Any author is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to Headley Brothers Little, The Invicta Press, Ashford, Kent. Coloured illustrations will be charged to authors, unless a special grant is authorized by the Editor.
- Orders for reprints must be sent when returning galley proofs, and for this purpose special forms are supplied.
- 6. Editorial communications may be addressed to The EDITOR, Journal of Laryngology, c/o Headley Brothers Ltd., The Invicta Press, Ashford, Kent.
- 7. The annual subscription is £8 (eight pounds sterling) post free; U.S.A. \$23 post free, and payable in advance.
- Single copies of current or back numbers (when available) will be on sale at £1 (including postage) each.
- All subscriptions, advertising and business communications should be sent to the publishers, or subscription agents.

HEADLEY BROTHERS LTD.

The Invicta Press, Ashrond, Kent.

O Journal of Laryngology and Otology Ltd., 1972

AMPLIVOX 84

The successor to the world's most widely used clinical Audiometer.



Take a look at all the advantages you get with this superb audiometer.


- Pure tone at eleven frequencies—now including 750 Hz
- Bone conduction at eight frequencies
- Narrow band masking via insert and headphones Automatic presentation
- ISO specification
- Trimpot calibration
- Continuously variable second channel for ABLB (Fowler) recruitment test
- Tone decay test
- Illuminated patient's signal

PLUS Calibrated speech audiometry via microphone or tape.

Write now for illustrated brochure to:

AMPLIVOX HEARING CONSERVATION LTD

Beresford Avenue, Wembley, Middx. HAO 1RU. Phone: 01-902 8991





PETERS

Audiometers and Sound Proof Rooms for clinics, hospitals, schools and factories all over the world

Literature and further information from:

PETERS

Head Office: 55 Clarkehouse Road, Sheffield S10 2LE Telephone 0742 64434 Regional Sales and Service Centres

London: 45 New Cavendish Street, London W1 Telephone 01-935 2604

South West England: 7 St. Augustine's Parade, Bristol BS1 4 HX Telephone 0272 20737

Please mention The Journal of Laryngology and Otology when replying to advertisements

THE LARYNGOSCOPE

A Monthly Journal devoted to the disease of EAR, NOSE AND THROAT

Official organ for the American Laryngological Rhinological and Otological Society

Price \$25.00 per year Canada \$27.00 per year Foreign \$27.00 per year

ESTABLISHED 1896

JOSEPH H. OGURA, M.D. EDITOR

222 PINE LAKE ROAD, COLLINSVILLE, ILLINOIS 62234


For advertisement space in this Journal apply to

HEADLEY BROTHERS LTD.

Ashford, Kent

Please mention The Journal of Laryngology and Otology when replying to advertisements


THE STUART SILASTIC® PHARYNGO-LARYNGECTOMY TUBE WITH DACRON MESH BACKING



Repair after pharyngo-laryngectomy continues to present many problems. The method using a permanently sited plastic tube was described in 1966 (Journal of Laryngology and Otology 1966 - 80 - 382 -935). A new tube has now been designed consisting of a Silastic® Silicone rubber tube covered with Dacron Mesh. This material does not harden in situ and is easily modified in shape or length. The upper end of the tube is sutured into the floor of the mouth and the lower end slipped down the remaining oesophagus. Stabilization is facilitated by encircling the upper end with the hyoid bone which is left in place. In patients with a small pharyngostome the funnel end of the tube can be shaped with scissors to avoid compression oedema of the tongue. A feeding tube is passed through the prosthesis into the stomach for ten days, after which the patient swallows normally through the prosthesis and can then return to a normal diet avoiding large pieces of meat etc., which might obstruct the lumen; it is surprising how rarely, however, this occurs. Twenty-one patients have been treated in this way with satisfactory results.

D. W. Stuart, F.R.C.S., Consultant E.N.T. Surgeon, North Staffs Royal Infirmary, Stoke-on-Trent.

DOWN BROS AND MAYER & PHELPS LIMITED | CHURCH PATH MITCHAM SURREY ENGLAND



British evidence confirms value of

Serc is the first drug to offer comprehensive action against all the manifestations of Meniere's syndrome. A British trial* now confirms its clinical value. Objective tests of hearing and vestibular function in 11 patients before and after treatment with Serc showed a dramatic improvement in many cases. 'There was an overall average gain in hearing over the whole group in both ears.' Studies of nystagmus suggested an improvement in vestibular function. 'Subjectively, 9 out of 11 patients claimed an improvement in the symptom of vertigo and all 11 claimed reduction in tinnitus. *Journal of Laryngology, and Otology, 1971, 85:369.

> Serc is supplied as tablets each containing 8 mg. betahistine dihydrochloride. Further information available on request,

in Meniere's **syndrome**

duphar Duphar Laboratories Limited West end Southampton tel 04218 2281

Please mention The Journal of Laryngology and Otology when replying to advertisements