

Instruments and Methods

Estimation of permafrost thickness using ground geophysical measurements, and its usage for defining vertical temperature variations in continental ice and underlying bedrock

TAPIO RUOTOISTENMÄKI, JUKKA LEHTIMÄKI

Geological Survey of Finland, Geophysics Department, Betonimiehenkuja 4, SF-02150 Espoo, Finland

ABSTRACT. In bedrock covered by continental ice, the depth to electrically conductive saline waters enriched below permafrost can be defined by electromagnetic measurements. Thus, if the ice thickness is also measured we obtain estimates of permafrost thickness variations in the survey area. Moreover, using the relation between ice and permafrost thicknesses and approximations of the melting temperature of saline waters combined with the average annual temperature at the surface, we can estimate temperature variations within both the bedrock and the ice. These can further be used for estimating the temperature values at the ice base.

The method was developed for interpreting two geophysical field profiles measured on continental ice in western Dronning Maud Land, Antarctica. Gravity, magnetic and electromagnetic profiles were measured between the nunataks Basen and Plogen (≈ 22 km) and Basen and Fossilryggen (≈ 38 km). In the Basen–Fossilryggen profile the electromagnetic soundings revealed good conductors in the bedrock beneath the ice. The significant correlation between the ice thickness and the depth of the conductor in the bedrock strongly suggests that the conductors are not sedimentary rock layers but rather saline groundwater in which salinity has been enhanced by permafrost processes. From the regression curve we can conclude that the temperature vs depth curve in the ice can be represented by a first-degree line and that the “warm” ice base of about -2°C is expected to occur at depths of about 650 m, the base of permafrost in outcropping bedrock being at about 1 km. The mean temperature gradients in the ice and in the bedrock can be estimated as being about $20^{\circ}\text{C km}^{-1}$ and $13^{\circ}\text{C km}^{-1}$, respectively.

In the Basen–Plogen profile, conductors were detected only in the beginning of the profile, where ice thickness was below 650 m. In the main part of the profile where the ice thickness exceeds 650 m, no conductors were observed. In those areas our interpretation is that the ice base is “warm” and no saline waters enriched by permafrost are present.

INTRODUCTION

When predicting the durability of radioactive-waste repositories and the stability of the surrounding bedrock during future periods of glaciation, it is essential to model the behaviour and characteristics of groundwaters driven by permafrost related to continental ice sheets (e.g. Boulton and others, 1996). For example, Bath and others (1996) considered hydrogeological models in deducing how the groundwater system might have responded to changing climate, specifically with respect to glaciation. Certes and others (1996) considered the transport of radionuclides due to groundwaters driven by permafrost during glaciation cycles, while Delisle (1996a) developed a numerical model for permafrost growth and decay. King-Clayton and others (1996) presented a model for predicting potential changes in various climatic, stress, hydraulic and hydrochemical parameters over approximately the next 130 000 years when evaluating the future behaviour of a deep depository for spent fuel.

Like any other theoretical model, these also require testing using our current knowledge of permafrost and groundwater behaviour associated with existing and recent

glaciers. However, it is relatively difficult to obtain information on these parameters. In practice it can be done indirectly, for example, by observing bedrock and soil in terrains that have been covered by glacier ice (e.g. Delisle, 1996b; Glynn and Voss, 1996; Ivanovich and others, 1996). Direct observations can be made by drilling through the ice (e.g. Paterson, 1994, and references therein) and permafrost (e.g. Gold and others, 1973). However, the indirect methods are relatively uncertain, while drilling is expensive and slow and thus cannot provide extensive coverage.

A relatively rapid and economic way is to analyze the variations of permafrost geophysically using electromagnetic measurements to define the conductivity variations within and beneath the permafrost. Hoekstra and McNeill (1973) give a summary of resistivities of various soil and rock types and conclude that there is approximately a factor of 10 between the resistivity of frozen ($\approx -10^{\circ}\text{C}$) and unfrozen ($\approx 1^{\circ}\text{C}$) soil and rock. The resistivity values also differ between various rock and soil types, which thus can be used for determining lithological and soil variations in the permafrost. Sinha (1975, 1977), Daniels and others (1976) and Sinha and Stephens (1983) considered the theoretical background and gave examples of field tests of the electromag-

netic methods used in permafrost-covered areas. Odegård and others (1996) demonstrated the use of DC resistivity soundings for permafrost mapping in southern Norway.

In the following we describe a relatively simple model for explaining the electrical conductivity variations observed by us in the bedrock beneath the Antarctic continental ice sheet. The principle of this method is based on the hypothesis that the saline content of wet ground is reduced by freezing, and the resulting enriched saline waters are "squeezed" below the permafrost (e.g. Daniels and others, 1976; Pisarskii, 1983; Nurmi, 1985). The model is based on the fact that the observations we made are difficult to explain solely by lithological conductivity variations, so the influence of saline waters on the conductivity anomalies seems apparent.

ESTIMATION OF PERMAFROST THICKNESS IN THE BEDROCK, AND EVALUATION OF TEMPERATURE VARIATIONS IN BEDROCK AND ICE BASE

Presumptions and a theoretical example

- (1) In general, the temperature vs depth curve, $T_i(z_i)$ ($^{\circ}\text{C}$), of continental ice measured from the upper surface of the ice can be arbitrary, i.e. of any degree; but the isotherms in the ice are assumed to be parallel to the ice surface (see Fig. 1). In the example shown in Figure 1, $T_i(z_i)$ is modelled for simplicity by a first-degree line, $13z_i - 15$ ($^{\circ}\text{C}$), where z_i is the depth in the ice. The temperature gradient in the ice is thus $13^{\circ}\text{C km}^{-1}$ and the average temperature on the ice surface is assumed to be about -15°C .
- (2) The temperature gradient in the bedrock, T_r' ($^{\circ}\text{C km}^{-1}$), is assumed to be of first degree and not to vary across the survey area. Järvinmäki and Puranen (1979) analyzed temperature gradients in 18 drillholes covering relatively variable bedrock environments in Finland. In all these holes the temperature vs depth curve is very close to a first-degree line. Gold and others (1973) give

examples of holes drilled in permafrost where a straight line would also give a very good approximation of the temperature vs depth curve. In the theoretical example in Figure 1 the permafrost thickness has been calculated using $T_r' = 8^{\circ}\text{C km}^{-1}$.

Calculation of the temperature vs depth curve in the ice

We define the parameters (see Fig. 1) as follows. T_1 ($^{\circ}\text{C}$) is the temperature at the ice surface, roughly equal to the average annual temperature in the area (in this model -15°C); T_2 ($^{\circ}\text{C}$) is the temperature at the base of the ice, $T_i(z_i)$; and T_3 ($^{\circ}\text{C}$) is the temperature at the base of the permafrost, roughly equal to the freezing/melting temperature of saline waters enriched below permafrost ($\approx -2^{\circ}\text{C}$).

The ice thickness z_i (km) can be interpreted from gravity or radar measurements. In Figure 1 the ice thickness varies between 0 and 1.25 km.

The depth to the bottom of the permafrost, z_p (km) (depth to electrically conductive saline waters enriched below permafrost), is determined by electromagnetic measurements. We thus obtain the permafrost thickness in the bedrock from z_r (km) $= z_p - z_i$.

By plotting the permafrost thickness (z_r) as a function of ice thickness (z_i) we obtain an estimate of the temperature gradient (T_r') in the frozen bedrock from the equation:

$$T_r' = \frac{T_3 - T_1}{z_{r0}} \quad (1)$$

where $z_{r0} = z_r(z_i = 0)$ is the permafrost thickness beneath exposed bedrock.

The temperature at the base of the permafrost in ice-covered bedrock can be written:

$$T_3 = -2 = T_i(z_i) + T_r' z_r. \quad (2)$$

We thus obtain:

$$T_i(z_i) = -2 - T_r' z_r, \quad (3)$$

which is the temperature value at the base of the ice ($= T_2$). In Equation (3) the temperature gradient in the bedrock (T_r') is assumed to be constant in the survey area, and the




Fig. 1. Geometry of the theoretical permafrost model. For definitions of the parameters see text. Hatched lines: continental ice (the thinner lines refer to thermal isotherms). Lighter shading: permafrost and bedrock (base $\approx -2^{\circ}\text{C}$ isoline). Darker shading: unfrozen bedrock. The values of permafrost thickness have been calculated from Equation (3). In areas where $z_r \approx 0$ the temperature at the ice base is about -2°C ; i.e. the ice is "warm based". The vertical exaggeration is about 15:1.

ice thickness (z_i) and permafrost thickness (z_r) are known for each survey point. Thus, when measuring the permafrost thickness values for variable ice-thickness values in the survey area we obtain from Equation (3) the ice-base temperature values as a function of z_i , i.e. an estimate of the curve $T_i(z_i)$.

It must be emphasized that the temperature vs depth curve in the ice can be estimated using the model described above if the temperature isotherms in the ice follow the course of the ice topography. If this assumption is valid we can construct even arbitrary higher-degree temperature vs depth curves $T_i(z_i)$ for the ice using Equation (3). For example, temperature vs depth curves measured in various glaciers (e.g. Paterson, 1994) indicate that in many cases they are higher-degree curves, presumably due to a complicated combination of both conduction and advection. In such cases the interpretation of $T_i(z_i)$ depends on the stability of the velocity of the ice flow and heat conduction in the survey area. The reliability of the interpreted higher-degree $T_i(z_i)$ curve can be evaluated from the variation of measured (z_i , z_r) values along the regression curve.

It is not possible to obtain estimates of the temperature vs depth curve $T_i(z_i)$ in the ice using our method if it varies throughout the survey area. However, we still can estimate permafrost thickness and temperature values at the ice base. The ice and permafrost thicknesses can be evaluated by using the geophysical methods described above. The temperature value at the ice base can then be calculated using the estimation of the temperature gradient in bedrock and permafrost thickness. If the temperature gradient in the bedrock cannot be determined from the permafrost-thickness vs ice-thickness curve it can be evaluated directly by measuring the permafrost thickness in outcrop.

Effect of electrically conductive bedrock

The method described above works best in areas where the proportion of electrically conductive rock types (e.g. gra-

phite-bearing schists) is small, as within crystalline basement or volcanics. The effect of conductive rocks can be evaluated from the permafrost-thickness (z_r) vs ice-thickness (z_i) curve. If a good correlation exists between these two parameters, there is reason to believe that the conductors represent saline waters below permafrost, because a correlation between ice thickness and the depth of lithological conductors is unlikely. This is demonstrated in the following example.

EXAMPLE: MEASUREMENTS IN ANTARCTICA

During the austral summer 1993–94 a Finnish expedition team carried out geophysical and geological investigations in western Dronning Maud Land, Antarctica. The area studied belongs to the Vestfjella mountains in Dronning Maud Land, near the Finnish base station Aboa (73°03' S, 13°25' W; black rectangle in Fig. 2a). The main aim of the geophysical measurements was to study the bedrock geology between the nunataks.

Gravity and magnetic profiles were measured between the nunataks Basen and Plogen (≈22 km) and Basen and Fossilryggen (≈38 km) with a station spacing of 250 m (Fig. 2b). The corresponding electromagnetic profiles had station spacings varying from 250 to 1000 m. The profiles have a common starting-point on an outcrop on Basen, and the angle between the profiles is about 50°. The region is covered by ice up to about 1000 m thick; only nunataks rise above the ice surface. Radio-echo profiles have been measured parallel to both profiles by a Swedish expedition team (Holmlund, 1994). They also reported that the maximum velocity of the ice in the Basen–Plogen profile close to Plogen is about 90 m year⁻¹. No velocity values have been reported for the Basen–Fossilryggen profile. The geology in the area has been described by Grind and others (1991); the main rock types exposed on the nunataks are basalts intruded by mafic sills. On Fossilryggen, sedimentary rocks with some




Fig. 2. Location of the survey area (a) and the measured profiles (b).

graphite-bearing layers are also present. The petrophysical parameters of the rocks were defined from samples collected by geologists. The geophysical and petrophysical measurements and their use in the interpretation of bedrock geology and ice-thickness variations are described in more detail by Lehtimäki and Ruotoistenmäki (1995) and Ruotoistenmäki and Lehtimäki (1997). In this paper we concentrate only on the results concerning the evaluation of permafrost.

ELECTROMAGNETIC INTERPRETATION: EVALUATION OF THE THICKNESS OF THE PERMAFROST AND TEMPERATURE VARIATIONS IN ICE AND BEDROCK

The electromagnetic sounding work was carried out with the multifrequency (2–20 000 Hz) electromagnetic sounding system “Sampo” described by Soininen and Jokinen (1991). The transmitter used was a horizontal loop with a diameter of 50 m. The receiver measures three components of the magnetic field. The coil separation can be increased to 1500 m, in which case the depth penetration is also about 1500 m.

Along the Basen–Fossilryggen profile, in areas where the ice thickness was mainly less than 650 m, soundings with a 1000 m coil separation showed significant conductors (20–500 Ωm) situated within the bedrock (Figs 3 and 4). The soundings on thick ice (>650 m) revealed no conductors. In the Basen–Ploggen profile, conductors were detected only in the beginning of the profile where the ice thickness was less than about 650 m. No conductors were observed even with greater coil separations in the main part of the profile, where the ice thickness exceeds 650 m.

The deep conductors observed in the bedrock have two possible explanations: (1) graphite-bearing sedimentary layers in basalts; (2) saline water layers enriched below the base of the permafrost. The latter process has been described by Daniels and others (1976), Pisarskii (1983) and Nurmi (1985). These two alternatives were evaluated by calculating the correlation between the ice thickness and the depth of the conductor in the bedrock (Fig. 5).

The study revealed a statistically significant negative

correlation between the variables (number of samples = 29; correlation = 0.83; significance level at which null hypothesis of zero correlation is disproved = 0.25×10^{-7}). Such a correlation could be caused by glacial erosion in bedrock above conductors which are situated at constant depth. However, from Figure 4 it can be seen that the depth variation between the conductors is about 500 m with respect to the sea-level surface ($z = 0$). Moreover, such a correlation due to tectonic processes is improbable. Thus, we can conclude that the conductors are not sedimentary layers but predominantly represent saline groundwaters enriched beneath the permafrost layer. The regression curve also indicates that the temperature vs depth curve is close to linear in the profile area.

The high ice velocities in the neighbouring Basen–Ploggen profile suggest that in the Basen–Fossilryggen profile under consideration, the temperature variations in the ice are not defined merely by conduction, but probably also by advection. However, the results of measurements by Echelmeyer and Harrison (1994) and Engelhardt and Kamb (1994) from an ice stream in Antarctica show that even in ice streams the temperature vs depth curve can sometimes be reasonably approximated by a straight line through most of the ice column. The small variation in measured z_I , z_T values along the regression line in Figure 5 also suggests that the first-degree approximation is relatively good in the survey area.

From the regression line we can estimate that the ice base will be at melting point (no permafrost) for ice thicknesses greater than about 650 m ($z_T = 0$, $z_I = 658$ m in Fig. 5) and that the base of the permafrost in outcropping bedrock is at about 1 km ($z_I = 0$, $z_T = 992$ m). These results also explain why no conductors were found in the main part of the Basen–Ploggen profile: because of the greater thickness of the ice sheet the ice base was “warm” and no saline waters enriched by permafrost are present. Using the average annual surface temperature in the area of about -15°C (Taalas and others, 1991) and taking the freezing point of saline waters as -2°C , we can estimate the mean linear temperature gradients in the ice and in the bedrock to be




Fig. 3. An example of the electromagnetic interpretation along the Basen–Fossilryggen profile at point $x = 6000$. The interpreted depths to the base of ice and permafrost have been drawn with dashed lines.




Fig. 4. Interpretation of electromagnetic soundings along the Basen-Fossilryggen profile. The white dots indicate the conductor depths used in correlation analysis in Figure 5. The variations of the ice thicknesses have been evaluated by gravity interpretation (Ruotoistenmäki and Lehtimäki, 1996) and radar measurements (Holmlund, 1994). The resistivities are represented by three broad ranges. The vertical exaggeration is 11:1.

about $13^{\circ}\text{C}/658\text{ m} \approx 20^{\circ}\text{C km}^{-1}$, and $13^{\circ}\text{C}/992\text{ m} \approx 13^{\circ}\text{C km}^{-1}$, respectively.

In areas covered by thin ice or in outcrop no conductors were observed. This was because the penetration depth of the configuration of the electromagnetic soundings we used was too small.

It must be emphasised that we do not necessarily attribute all conductors to the effect of saline waters. For example, the conductors near Fossilryggen probably represent a mixture of graphite-bearing schists and saline waters.

CONCLUSIONS

In bedrock covered by continental ice, the depth to saline waters enriched below permafrost can be defined by electromagnetic measurements. Thus, if the ice thickness is known we can estimate permafrost-thickness variations in the sur-

vey area. Moreover, using the relation between the ice and permafrost thicknesses and approximations of the melting temperature of saline waters combined with the average annual surface temperature, we can estimate the temperature variations in the bedrock and the ice. Using these estimates of temperature gradients we can also evaluate the temperature values at the base of the ice.

The results of field measurements on glaciated terrain in Antarctica support the conclusion that the method is applicable even in areas where occasional lithological conductors are to be expected. In conjunction with depth information for the ice, the electromagnetic measurements provide information concerning the distribution and conductivity of the saline water layers enriched below the permafrost. Using the average annual temperature in the survey area and the melting/freezing temperature of saline waters, we can obtain estimates of temperature variations in both the bedrock and the ice. In the future, the method should be




Fig. 5. Basen-Fossilryggen profile: the relation between the ice thickness and depth of the conductor in the bedrock.

tested in various other profiles in the survey area. In particular, we should try to obtain values for the “dashed” parts of the regression line in Figure 5. Rock samples from the lower exposures of the steep, vertically faulted cliffs should also be taken to analyze the porosity of the bedrock. Using the porosity and the electrical conductivity values we can then obtain estimates of the salinity of the groundwaters. Moreover, the method should also be tested in non-glaciated terrain where thick layers of permafrost are known to have existed since the last ice age. Also, the results of our measurements indicate that in the planning of future deep-drilling projects in areas covered by continental ice, the possibility of drilling on *thinner* ice and continuing the drilling in the *bedrock* below the base of permafrost down to the underlying conductors should also be considered. In the planning of such work, our method could be appropriate and the Basen–Fossilryggen region could be suitable for this purpose.

ACKNOWLEDGEMENTS

We want to thank our Swedish colleagues J.-O. Näslund and P. Holmlund for providing us with radar data and P. Jansson for measuring the coordinates of the tie-points of our profiles. Moreover we wish to thank our fellow geologists R. Blomqvist, P. Vuorela, K. Nenonen and H. Hirvas from the Geological Survey of Finland (GSF) for pointing out possible explanations for our observations. Finally, we express our gratitude to the GSF and the Ministry of Trade and Industry, Finland, for supporting the Finnish Antarctic Research Programme “Finnarp 93–94”.

REFERENCES

- Bath, A., H. G. Richard and N. A. Feast. 1996. Hydrochemical impacts on past glacial cycles. Abstract. In King-Clayton, L., ed. *Glaciation and hydrogeology: workshop on the impact of climate change and glaciations on rock stresses, groundwater flow and hydrogeochemistry — past, present and future, 17–19 April 1996, Stockholm*. Stockholm, Nordic Nuclear Safety Research (NKS), A-1.
- Boulton, G. S., D. Read and D. Ross. 1996. Modelling the signal of long-term environmental change in the geochemistry of groundwater — an approach to testing models of long-term rock mass properties. Abstract. In King-Clayton, L., ed. *Glaciation and hydrogeology: workshop on the impact of climate change and glaciations on rock stresses, groundwater flow and hydrogeochemistry — past, present and future, 17–19 April 1996, Stockholm*. Stockholm, Nordic Nuclear Safety Research (NKS), A-9.
- Certes, C., P. Baudoin, F. Cuisiat and D. Gay. 1996. Glaciation modelling in safety evaluation of radioactive waste disposal: experience from the EVEREST project. Abstract. In King-Clayton, L., ed. *Glaciation and hydrogeology: workshop on the impact of climate change and glaciations on rock stresses, groundwater flow and hydrogeochemistry — past, present and future, 17–19 April 1996, Stockholm*. Stockholm, Nordic Nuclear Safety Research (NKS), A-10.
- Daniels, J., G. V. Keller and J. J. Jacobson. 1976. Computer-assisted interpretation of electromagnetic soundings over a permafrost section. *Geophysics*, **41** (4), 752–763.
- Delisle, G. 1996a. A numerical model on permafrost growth and decay. Abstract. In King-Clayton, L., ed. *Glaciation and hydrogeology: workshop on the impact of climate change and glaciations on rock stresses, groundwater flow and hydrogeochemistry — past, present and future, 17–19 April 1996, Stockholm*. Stockholm, Nordic Nuclear Safety Research (NKS), A-12.
- Delisle, G. 1996b. Cryogenic fractures in salt domes in northern Germany? Abstract. In King-Clayton, L., ed. *Glaciation and hydrogeology: workshop on the impact of climate change and glaciations on rock stresses, groundwater flow and hydrogeochemistry — past, present and future, 17–19 April 1996, Stockholm*. Stockholm, Nordic Nuclear Safety Research (NKS), A-13.
- Echelmeyer, K. and W. Harrison. 1994. Temperature measurements in the margin of Ice Stream B, 1992–1993. *Antarct. J. U.S.*, **28**(5), 66–67.
- Engelhardt, H. and B. Kamb. 1994. Vertical temperature profile of Ice Stream B. *Antarct. J. U.S.*, **28**(5), 63–66.
- Glynn, P. D. and C. Voss. 1996. Glaciation and ground water geochemistry in the Fennoscandian Shield: deep oxygenated ground water of glacial origin. Abstract. In King-Clayton, L., ed. *Glaciation and hydrogeology: workshop on the impact of climate change and glaciations on rock stresses, groundwater flow and hydrogeochemistry — past, present and future, 17–19 April 1996, Stockholm*. Stockholm, Nordic Nuclear Safety Research (NKS), A-17.
- Gold, L. W. and A. H. Lachenbruch. 1973. Thermal conditions in permafrost: a review of North American literature. In *Permafrost ... Second International Conference ... 1973, Yakutsk, U.S.S.R.* Washington, DC, National Academy of Sciences, 3–25.
- Grind, H., J. Siivola and A. Luttinen. 1991. A geological overview of the Vestfjella Mountains in western Dronning Maud Land, Antarctica. In *Finnarp-89 Symposium Publication*. Helsinki, Ministry of Trade and Industry, 11–15. (Report 1.)
- Hockstra, P. and D. McNeill. 1973. Electromagnetic probing of permafrost. In *Permafrost ... Second International Conference ... 1973, Yakutsk, U.S.S.R.* Washington, DC, National Academy of Sciences, 517–526.
- Holmlund, P. 1994. Depth soundings. In Näslund, J.-O. and A. Bodin, eds. *Studies on ice dynamics, mass balance and paleoclimate in East Antarctica 1991/92*. Stockholm, Stockholms Universitet. Naturgeografiska Institutionen, 6. (Forskningsrapport 98.)
- Ivanovich, M., E. McDermott, S. Frapce and C. J. Hawkesworth. 1996. Paleoclimatic controls on hydrological systems: evidence from U-Th dated calcite veins in the Fennoscandian and Canadian shields. Abstract. In King-Clayton, L., ed. *Glaciation and hydrogeology: workshop on the impact of climate change and glaciations on rock stresses, groundwater flow and hydrogeochemistry — past, present and future, 17–19 April 1996, Stockholm*. Stockholm, Nordic Nuclear Safety Research (NKS), A-29.
- Järvinmäki, P. and M. Puranen. 1979. Heat flow measurements in Finland. In Cermák, V. and L. Rybach, eds. *Terrestrial heat flow in Europe*. Berlin, Springer-Verlag, 173–177.
- King-Clayton, L., F. Kautsky, N. A. Chapman, N.-O. Svensson, G. de Marsily and E. Ledoux. 1996. The central climate change scenario: SKI's SITE-94 project to evaluate the future behaviour of deep repository for spent-fuel. Abstract. In King-Clayton, L., ed. *Glaciation and hydrogeology: workshop on the impact of climate change and glaciations on rock stresses, groundwater flow and hydrogeochemistry — past, present and future, 17–19 April 1996, Stockholm*. Stockholm, Nordic Nuclear Safety Research (NKS), A-35.
- Lehtimäki, J. and T. Ruotoistenmäki. 1995. Geophysical ground measurements on glaciated terrain in Queen Maud Land, Antarctica. Abstract. In *EAEG 57th Conference, Glasgow, 25 May–2 June 1995. Extended abstracts. Vol. 1*. Glasgow, EAGE Business Office, D056.
- Nurmi, P. A. 1985. *Mahdolliset ympäristöolosuhteiden pitkäaikaismuutokset Suomessa ja niiden vaikutukset syväällä oleviin kallioföhjavesiin* [Future long-term environmental changes in Finland: their influence on deep-seated groundwater]. Helsinki, Nuclear Waste Commission of Finnish Power Companies. (Report YJT-85-21.)
- Odegård, R. S., M. Hoelzle, K. V. Johansen and J. L. Sollid. 1996. Permafrost mapping and prospecting in southern Norway. *Nor. Geogr. Tidsskr.*, **50**(1), 41–53.
- Paterson, W. S. B. 1994. *The physics of glaciers. Third edition*. Oxford, etc., Elsevier.
- Pisarskii, B. E. 1983. Groundwater in permafrost regions. In Pinneker, E. V., ed. *General hydrogeology*. Cambridge, Cambridge University Press, 93–98.
- Ruotoistenmäki, T. and J. Lehtimäki. 1997. Analysis of bedrock geology and thermal gradients using geophysical ground measurements on glaciated terrain in Queen Maud Land, Antarctica. In *Seventh International Symposium on Antarctic Earth Sciences, 10–15 September 1995, Siena, Italy. Proceedings*.
- Sinha, A. K. 1975. Electromagnetic sounding over layered permafrost terrain. *Geol. Surv. Can. Pap.*, **75-1A**, 149–151.
- Sinha, A. K. 1977. A theoretical study on electromagnetic probing of permafrost terrains. *Can. J. Earth Sci.*, **14**(10), 2388–2401.
- Sinha, A. K. and L. E. Stephens. 1983. Permafrost mapping over a drained lake by electromagnetic induction methods. *Geol. Surv. Can. Pap.* 83-1A, 213–220.
- Soinenen, H. and T. Jokinen. 1991. Sampo, a new wide-band electromagnetic system. In *53rd Meeting of the European Association of Exploration Geophysics and Technical Exhibition, Florence, Italy. Technical programme and abstracts of papers*.
- Taalas, P., S. Huovila, K. Karlsson and J. Launiainen. 1991. Climatology of the Vestfjella region. In *Finnarp-89 Symposium Publication*. Helsinki, Ministry of Trade and Industry, 133–136. (Report 1.)