

ON A HOMOLOGY THEORY ASSOCIATED TO FOLIATIONS

YOSHIHIRO SHIKATA

Dedicated to Professor Katuzi Ono on his 60th birthday

Introduction.

In this note, we associate a foliated singular homology theory to a foliation on a compact manifold X and construct a spectral sequence which relates the foliated homology to the ordinary homology of X . Since the foliated homology is so constructed as to be related closely to a certain topological behavior of the foliation, we may expect that further study of the spectral sequence reveals some information on topology of foliation. The study and applications will be given in a sequel.

1. A double complex associated to a plane field.

Let X be a compact smooth manifold having a plane field Π . A smooth map α of $(i + j)$ -cube $I^i \times I^j$ into X is called a singular (i, j) -cube on $(X; \Pi)$, if the map α satisfies that

(i) for any $Q \in I^j$, the restriction of α to the i -plane $I^i \times Q$ of $I^i \times I^j$ is tangent to the plane field, that is,

$$d\alpha|_{I^i \times Q}(T_p(I^i)) \subset \Pi_{\alpha(P \times Q)}, \quad (\text{for all } P \in I^i),$$

(ii) for any $P \in I^i$, the restriction of α to j -plane $P \times I^j$ is transversal to the plane field, that is,

$$d\alpha|_{P \times I^j}(T_Q(I^j)) \cap \Pi_{\alpha(P \times Q)} = 0 \quad (\text{for all } Q \in I^j).$$

Denote by $C'_{i,j}(X, \Pi)$ the free abelian group generated over all the (i, j) -cubes and define face operators $F_{k; \varepsilon}^i, F_{; k}^\varepsilon$ ($\varepsilon = 0, 1$) as to be the linear maps satisfying that

Received March 31, 1969

$$\begin{aligned} (F_{;k}^\varepsilon \alpha)(s_1 \cdots s_i, t_1 \cdots t_j) &= \alpha(s_1 \cdots s_{k-1}, \varepsilon, s_k \cdots s_i, t_1 \cdots t_j) \\ (F_{;k}^\varepsilon \alpha)(s_1 \cdots s_i, t_1 \cdots t_j) &= \alpha(s_1 \cdots s_i, t_1 \cdots t_{k-1}, \varepsilon, t_k \cdots t_j). \end{aligned}$$

Define operators $\xi = \xi_{i,j}, \eta = \eta_{i,j}$ on $C'_{i,j}$ by

$$\xi = \sum_{k=1}^i (-1)^k (F_{;k}^{-1} - F_{;k}^0), \quad \eta = (-1)^i \sum_{k=1}^j (-1)^k (F_{;k}^{-1} - F_{;k}^0),$$

then ξ, η satisfy that

$$\xi \circ \xi = \eta \circ \eta = 0, \quad \xi \circ \eta + \eta \circ \xi = 0.$$

Let D_k , (resp. $D_{;k}$) be linear maps of $C'_{i-1,j}$ into $C'_{i,j}$ (resp. $C'_{i,j-1} \rightarrow C'_{i,j}$)

such that

$$\begin{aligned} (D_k \alpha)(s_1 \cdots s_i, t_1 \cdots t_j) &= \alpha(s_1 \cdots s_{k-1}, s_k \cdots s_i, t_1 \cdots t_j) \\ (D_{;k} \alpha)(s_1 \cdots s_i, t_1 \cdots t_j) &= \alpha(s_1 \cdots s_i, t_1 \cdots t_{k-1}, t_k \cdots t_j) \end{aligned}$$

then it is easy to see that

$$\begin{aligned} D_k \circ D_m &= D_{m+1} \circ D_k, \quad D_{;k} \circ D_{;m} = D_{;m+1} \circ D_{;k} \quad (k \leq m), \\ F_{;k}^\varepsilon \circ D_m &= \begin{cases} D_{m-1} \circ F_{;k}^\varepsilon & (k < m), \\ 1 & (k = m), \\ D_m \circ F_{;k-1}^\varepsilon & (k > m), \end{cases} \quad F_{;k}^\varepsilon \circ D_{;m} = \begin{cases} D_{;m-1} \circ F_{;k}^\varepsilon & (k < m) \\ 1 & (k = m) \\ E_{;m} \circ F_{;k-1}^\varepsilon & (k > m). \end{cases} \end{aligned}$$

Therefore the quotient $C_{i,j}$ of $C'_{i,j}$ given by

$$C_{i,j} = C'_{i,j} / \sum_{k=1}^{i-1} D_k C'_{i-1,j} + \sum_{k=1}^{j-1} D_{;k} C'_{i,j-1}$$

turns out to be a double complex having boundary operators ξ, η . Denote by $(S_n(X), \partial)$ the (normalized) cubical singular chain complex of X , then the single complex $C_n = \sum_{i+j=n} C_{i,j}$ is naturally imbedded into $S_n(X)$ as a subcomplex because the operators ξ, η and ∂ satisfy that

$$\xi + \eta = \partial.$$

LEMMA 1. Let N be an integer and let $e_{p,q}$ be a map of $I^i \times I^j$ into $I^{i-1} \times [p/N, p+1/N] \times I^{i-1} \times [q/N, q+1/N]$ given by

$$e_{p,q}(s_1 \cdots s_i, t_1 \cdots t_j) = (s_1 \cdots s_{i-1}, p + s_i/N, t_1 \cdots t_{j-1}, q + t_j/N).$$

then for any $\alpha \in C'_{i,j}$ there exists $f(\alpha) \in C'_{i+1,j} + C'_{i,j+1}$ such that

$$(\xi + \eta)f(\alpha) + f((\xi + \eta)\alpha) = \alpha - \sum_{p,q} \alpha \circ e_{p,q} \text{ in } C_{i,j}.$$

For a chain $\gamma = \sum c_k \alpha_k (c_i \neq 0)$ of $C'_{i,j}$, define $\text{Car}(\gamma)$ to be $\bigcup_k \alpha_k(I^i \times I^j)$ and for an open set U in X let

$$\begin{aligned} C'_{i,j}{}^U &= \{\gamma \in C'_{i,j} \mid \text{Car}(\gamma) \subset U\}, \\ C_{i,j}{}^U &= C'_{i,j}{}^U / \sum_{k=1}^{i-1} D_k C'_{i-1,j}{}^U + \sum_{k=1}^{j-1} D_k C'_{i,j-1}{}^U, \\ C_n{}^U &= \sum_{i+j=n} C_{i,j}{}^U. \end{aligned}$$

PROPOSITION 1. *Let $\{U\}$ be an open covering of X and let $C_n^{(U)}$ be the union of $C_n{}^U$, $U \in \{U\}$. Then the inclusion $C_n^{(U)}$ into C_n is a chain equivalence.*

A plane field Π on X is said to be completely integrable if there is a foliation on X such that for any $x \in X$, Π_x agrees with the tangent plane at x of the leaf through x .

LEMMA 2. *Suppose that the plane field Π on X is completely integrable then for any open set U of X and for any point $x \in U$, there is a neighbourhood V of x such that $V \subset U$ and $C_n{}^V$ is acyclic.*

Proof. Take a flat coordinate $(x_1 \cdots x_p, y_1 \cdots y_q)$ ($-\varepsilon < x_i, y_i < \varepsilon$) in a neighbourhood V of x so that p -planes defined by $\{y_i = \text{const.}\}$ form (locally) the leaves of the foliation around x . Then the homotopy h_t^1 in V defined by

$$h_t^1(x_1 \cdots x_p, y_1 \cdots y_q) = (tx_1 \cdots tx_p, y_1 \cdots y_q)$$

gives a chain homotopy in $C_n{}^V$ between $\alpha \in C'_{i,j}{}^V$ and a chain $\pi^1(\alpha) \in C'_{0,j}{}^V$, also the homotopy h_t^2 defined by

$$h_t^2(0 \cdots 0, y_1 \cdots y_q) = (0 \cdots 0, ty_1 \cdots ty_q)$$

gives a chain homotopy in $C_n{}^V$ between $\pi^1 \alpha \in C'_{0,j}{}^V$ and $\pi^2(\pi^1 \alpha) \in C'_{0,0}{}^V$.

Thus, following a standard method of homology theory of (co-)sheaves ([1] [4]), we have

PROPOSITION 2. *If the plane field Π on X is completely integrable then the Čech homology group $H_*(X; G)$ is isomorphic to the projective limit of the homology groups $H_*(C_n^{(U)}; G)$ for any module G .*

Therefore combining the propositions 1 and 2, we have

THEOREM 1. *If the plane field Π on a compact manifold X is completely integrable, then for any module G . It holds that*

$$H_*(X; G) = H_*(C_n \otimes G, \xi + \eta).$$

2. A spectral sequence associated to $C_{i,j}$.

Introduce two regular filtrations F^I and F^II into $C_{i,j}$ as follows:

$$F_p^I C = \sum_{r \leq p} \sum_q C_{r,q}, \quad F_q^{II} C = \sum_{s \leq q} \sum_p C_{p,s}.$$

Then there result filtrations of $H_*(C, \xi + \eta)$ and two spectral sequences $I_{i,j}^r$ and $II_{i,j}^r$ ([2]), we investigate several terms of the spectral sequences in case that Π is completely integrable.

2.1 On terms $II_{i,j}^1$.

It is known ([2]) that

$$I_{i,j}^1 = H(C_{i,j}, \eta), \quad II_{i,j}^1 = H(C_{i,j}, \xi).$$

On the other hand, we easily have

LEMMA 3. *If the plane field Π is completely integrable, then the chain complex $(C_{i,0}, \xi)$ is identified with the chain complex $(\sum_L S_i(L), \partial)$, where the sum \sum_L is taken over all the leaves of the foliation.*

The lemma 3 above obviously yields

PROPOSITION 3. *If the plane field Π is completely integrable, then it holds that*

$$II_{i,0}^1 = \sum_L H_i(L; Z),$$

where the sum \sum_L is taken over all the leaves of the foliation.

LEMMA 4. *If the plane field Π is completely integrable and if an (i, j) -cube α is non zero in $C'_{i,j} / \sum_k D_{i,k} C'_{i,j-1}$, then the first i -faces $F_{k_i}^\varepsilon \alpha (\varepsilon = 0, 1)$ are non zero in $C'_{i-1,j} / \sum_k D_{i,k} C'_{i-1,j-1}$.*

Proof. Suppose on the contrary that $F_{k_i}^\varepsilon \alpha$ is independent of the coordinate t_1 of I^j and let $Q(x)$ denote the point $(x, t_2 \cdots t_n)$ of I^j , then for any $x \in I$, $\alpha(I^i \times Q(x))$ lies in the same leaf L , because they have a point $\alpha(s_1 \cdots s_i, x, t_2 \cdots t_j)$ in common, therefore the vector $\partial/\partial x = \partial/\partial t_1$ is mapped by $d\alpha$ into the tangent plane of L , thus we have that $\partial\alpha/\partial t_1 = 0$ at every point of $I^i \times I^j$ and conclude that α itself is independent of t_1 .

LEMMA 5. *If the plane field Π is completely integrable, then for each chain $\gamma \in C_{i,j}$, there corresponds a chain $\gamma' \in C'_{i,j}$, such that $\gamma' = \gamma$ in $C_{i,j}$ and satisfying*

- (i) $\xi\gamma = 0$ in $C_{i,j}$ implies $\xi\gamma' = 0$ in $C'_{i,j} / \sum_k D_k; C'_{i-1,j}$,
- (ii) $\xi\gamma = \gamma_2$ in $C_{i,j}$ implies $\xi\gamma'_1 = \gamma'_2$ in $C'_{i,j} / \sum_k D_k; C'_{i-1,j}$.

Proof. Let $V_{i,j}$ be a set of all the (i, j) -cubes which are independent of t_k for some $1 \leq k \leq j$ and let $W_{i,j}$ be the set of all the (i, j) -cubes which are not in $V_{i,j}$. Then $C'_{i,j}$ decomposes into a direct sum of free groups $A_{i,j}$ and $B_{i,j}$ generated over $V_{i,j}$ and $W_{i,j}$ respectively. Denote by x' and x'' the $A_{i,j}$ and $B_{i,j}$ component of $x \in C'_{i,j}$ respectively, then Lemma 4 yields that

$$(\xi x)'' = \xi(x'') \text{ for } x \in C'_{i,j}.$$

Now choose a representative $\tilde{\gamma} \in C'_{i,j}$ for $\gamma \in C_{i,j}$ and define γ' by $\gamma' = (\tilde{\gamma})'$. Suppose that

$$\xi\gamma = 0 \text{ in } C_{i,j},$$

then

$$\xi\tilde{\gamma} = x \text{ for some } x \in \sum_k D_k; C'_{i-2,j} + \sum_k D_k; C'_{i-1,j-1}.$$

Thus we have that

$$\xi(\gamma') = \xi(\tilde{\gamma} - (\tilde{\gamma})'') = \xi\tilde{\gamma} - (\xi\tilde{\gamma})'' = x - x'',$$

therefore we see that

$$\xi(\gamma') = 0 \text{ in } C'_{i-1,j} / \sum_k D_k; C'_{i-2,j},$$

and we also see (ii) as in the same way above.

Define a linear map $F_{i,Q}(Q \in I_j)$ of $C'_{i,j}$ into $C'_{i,0}$ by

$$(F_{i,Q}\alpha)(s_1 \cdots s_i) = \alpha(s_1 \cdots s_i, t_1(Q) \cdots t_j(Q)),$$

then we easily see that $F_{i,Q}$ commutes with F_k and D_k for any $1 \leq k \leq i$.

Thus Lemma 5 yields

PROPOSITION 4. *If the plane field Π is completely integrable, then for each chain $\gamma \in C_{i,j}$ there exists a representative $\gamma' \in C'_{i,j}$ satisfying*

- (i) $\xi\gamma = 0$ in $C_{i,j}$ implies $\xi(F_{i,Q}\gamma') = 0$ in $C_{i,0}$,
- (ii) $\xi\gamma_1 = \gamma_2$ in $C_{i,j}$ implies $\xi(F_{i,Q}\gamma'_1) = F_{i,Q}\gamma'_2$ in $C_{i,0}$.

In the other words, $\gamma \in C_{i,j}$ is a ξ -cycle only if there is a j -dimensional

family of ∂ -cycles of $\sum_L S_i(L)$ and a ξ -cycle γ_1 is a ξ -boundary only if every ∂ -cycle of the family is a ∂ -boundary of $\sum_L S_{i+1}(L)$.

2.2 On terms $\Pi_{i,j}^{\xi}$

Since η commutes with ξ , η defines canonically a homomorphism of $H(C_{i,j}, \xi)$ into $H(C_{i,j-1}, \xi)$ which is denoted also by η . Then it is known ([2]) that

$$\Pi_{i,j}^{\xi} = H(H(C_{i,j}, \xi), \eta).$$

Therefore we deduce from Proposition 3 that

PROPOSITION 5. *If the plane field Π is completely integrable, then it holds that*

$$\Pi_{i,0}^{\xi} = \sum_L H_i(L; Z) \eta(H(C_{i,1}, \xi)),$$

where the sum \sum_L is taken over all the leaves of the foliation.

DEFINITION. *In case that the plane fields Π is derived from a foliation, we call the terms $\Pi_{i,0}^1$ and $\Pi_{i,0}^2$ the i -th homology groups of the foliation of the first and second kind, respectively, and we write*

$${}^1 H_i(\Pi) = \Pi_{i,0}^1, \quad {}^2 H_i(\Pi) = \Pi_{i,0}^2.$$

Roughly speaking, for $p = \dim \Pi$, the dimension of free part of $\Pi_{p,0}^1$ represents the number of oriented closed leaves of the foliation and that of $\Pi_{p,0}^2$ corresponds to the number of oriented closed leaves which can not be connected each other by any 1-dimensioned family of oriented closed leaves.

2.3 On terms $\Pi_{i,j}^{\infty}$

Since it is known ([2]) that

$$\sum_{i+j=n} \Pi_{i,j}^{\infty} = \mathcal{S}H(C_n, \xi + \eta),$$

we see that Theorem 1 implies that

PROPOSITION 6. *If the plane field Π is completely integrable, then it holds that*

$$\sum_{i+j=n} \Pi_{i,j}^{\infty} = \mathcal{S}H_n(X; Z).$$

3. Remarks for dually foliated structures.

A pair of plane fields (Π_1, Π_2) on a compact manifold X is called an orthogonal splitting of the tangent bundle $T(X)$ if it holds that

$$\Pi_1 \perp \Pi_2, \quad \Pi_1 + \Pi_2 = T(X).$$

Let $(X; \Pi_1, \Pi_2)$ be a compact manifold with an orthogonal splitting (Π_1, Π_2) . A smooth map α of $(i + j)$ -cube $I^i \times I^j$ into X is said to be a singular (i, j) -cube of $(X; \Pi_1, \Pi_2)$ if the map α satisfies that

- (i) for any $Q \in I^j$, the restriction of α to the i -plane $I^i \times Q$ of $I^i \times I^j$ is tangent to the plane field Π_1 .
- (ii) for any $P \in I^i$, the restriction α to the j -plane $P \times I^j$ of $I^i \times I^j$ is tangent to the plane field Π_2 .

The free abelian group $\Gamma'_{i,j}$ generated over all the (i, j) -cubes of $(X; \Pi_1, \Pi_2)$ has face operators $F_{i,k}, F_{k,i}$ and degeneracy operators $D_{i,k}, D_{k,i}$ defined in the same way as in §1. Therefore $\Gamma'_{i,j}$ and the quotient

$$\Gamma_{i,j} = \Gamma'_{i,j} / \sum_k D_{k,i} \Gamma'_{i-1,j} + \sum_k D_{i,k} \Gamma'_{i,j-1}$$

turn out to be double complexes with boundary operators ξ, η given by

$$\xi = \sum (-1)^k (F_{k,i}^1 - F_{k,i}^0), \quad \eta = \sum (-1)^{k+i} (F_{i,k}^1 - F_{i,k}^0).$$

An orthogonal splitting (Π_1, Π_2) of $T(X)$ is said to be completely integrable if the plane fields $\Pi_i (i = 1, 2)$ are completely integrable by foliations $(F_i) (i = 1, 2)$ on X , and if each point $x \in X$ has a coordinate neighbourhood $U(x_1 \cdots x_p, y_1 \cdots y_q)$ such that the p -plane given by $y_i = \text{const.}$ is a leaf of (F_1) in U and the q -plane given by $x_i = \text{const.}$ is a leaf of (F_2) in U . The following theorem is proved easily as in the same way in §1;

THEOREM 2. *If an orthogonal splitting (Π_1, Π_2) on a compact manifold X is completely integrable, then for the single complex $\Gamma_n = \sum_{i+j=n} \Gamma_{i,j}$, it holds that*

$$H_*(X; Z) = H_*(\Gamma_n, \xi + \eta).$$

Therefore as in §2 the spectral sequence $(E_{i,j}^r)$ associated to the filtration

$$F_q = \sum_{s \leq q} \sum_p \Gamma_{p,s}$$

relates $E_{i,j}^2 = H(H(\Gamma_{i,j}, \xi), \eta)$ to $H_*(X; Z)$.

Since $C_{i,0} = \Gamma_{i,0}$ we obviously have

LEMMA 6. *If an orthogonal splitting (Π_1, Π_2) is completely integrable, then it holds that*

$${}^1H_i(\Pi_1) = E_{i,0}^1.$$

Define linear maps $F_{;Q}$ and $F_{;P}$, ($Q \in I_j$, $P \in I_i$) of $\Gamma'_{i,j}$ into $\Gamma'_{i,0}$ and $\Gamma'_{0,j}$, respectively, by

$$\begin{aligned} (F_{;Q}\alpha)(s_1 \cdots s_i) &= \alpha(s_1 \cdots s_i, t_1(Q) \cdots t_j(Q)) \\ (F_{;P}\alpha)(t_1 \cdots t_j) &= \alpha(s_1(P) \cdots s_i(P), t_1 \cdots t_j). \end{aligned}$$

Then we have the following proposition of which proof is quite similar to that of Proposition 4.

PROPOSITION 7. *If an orthogonal splitting (Π_1, Π_2) is completely integrable, then for each chain $\gamma \in \Gamma_{i,j}$ there exists a representative $\gamma' \in \Gamma'_{i,j}$ such that*

- (i) -1 $\xi\gamma = 0$ in $\Gamma_{i,j}$ implies $\xi(F_{;Q}\gamma') = 0$ in $\Gamma_{i,0}$
- (i) -2 $\eta\gamma = 0$ in $\Gamma_{i,j}$ implies $\eta(F_{;P}\gamma') = 0$ in $\Gamma_{0,j}$
- (ii)-1 $\xi\gamma_1 = \gamma_2$ in $\Gamma_{i,j}$ implies $\xi(F_{;Q}\gamma'_1) = F_{;Q}\gamma'_2$ in $\Gamma_{i,0}$
- (ii)-2 $\eta\gamma_1 = \gamma_2$ in $\Gamma_{i,j}$ implies $\eta(F_{;P}\gamma'_1) = F_{;P}\gamma'_2$ in $\Gamma_{0,j}$.

COROLLARY 1. *Let (Π_1, Π_2) be a completely integrable splitting of $T(X)$, if there exists non trivial ξ -cycle $\gamma \in \Gamma_{i,j}$ or η -cycle $\gamma \in \Gamma_{i,j}$, then there exists a j -dimensional family of ξ -cycles $\gamma_Q \in \Gamma'_{i,0}(Q \in I^j)$ or an i -dimensional family of η -cycles $\gamma_P \in \Gamma'_{0,j}(P \in I^i)$, respectively.*

Thus we see that for a completely integrable splitting, if there is a non trivial η -cycle $x \in \Gamma_{0,j}$ of the form

$$x = \xi y + \eta z, \quad y \in \Gamma_{1,j}, \quad z \in \Gamma_{0,j+1},$$

then there is an 1-dimensional family of η -cycles $x_P \in \Gamma'_{0,j}$ ($P \in I$) or a $(j - 1)$ -dimensional family of ξ -cycles $x_Q \in \Gamma'_{1,0}$ ($Q \in I^{j-1}$).

Now assume that X is a 3-dimensional compact manifold for which

$$H_2(X; Z) = 0$$

and that $T(X)$ splits into mutually orthogonal 1-plane field Π_1 and 2-plane field Π_2 such that the pair (Π_1, Π_2) is completely integrable. Then if there is a non trivial η -cycle $x \in \Gamma_{0,2}$, an easy computation of the spectral sequence $(E_{i,j}^r)$ shows that there exists at least one non trivial ξ -cycle $y \in \Gamma_{1,0}$ or a 1-dimensional family of η -cycles $z_p \in \Gamma'_{0,2}$. Thus we have the following theorem as an elementary application of the spectral sequence.

THEOREM 3. *Let X be a compact 3-manifold having a 2-dimensional foliation (F_2) and let (F_1) be the 1-dimensional foliation defined by the 1-plane field orthogonal to (F_2) . Then, if (F_2) has only isolated closed leaves and if 2-dimensional integral homology of X vanishes, there should be at least one closed orbit in (F_1) .*

BIBLIOGRAPHY

- [1] G. Bredon, *Cosheaves and homology*, Pacific J. of Math. Vol. 25, (1968), 1–32.
- [2] H. Cartan and S. Eilenberg, *Homological algebra*, Princeton Univ. Press, (1956).
- [3] G. Reeb, *Sur les espaces fibrés et les varietés feuilletées*, Actualités scientifiques et industrielles, Hermann Paris, n° 1183.
- [4] E. Spanier, *Algebraic topology*, McGraw-Hill, (1966).

*Mathematical Institute,
Nagoya University*