

FORM RINGS AND REGULAR SEQUENCES⁽¹⁾

PAOLO VALABREGA AND GIUSEPPE VALLA

Introduction

Hironaka, in his paper [H₁] on desingularization of algebraic varieties over a field of characteristic 0, to deal with singular points develops the algebraic apparatus of the associated graded ring, introducing standard bases of ideals, numerical characters ν^* and τ^* etc. Such a point of view involves a deep investigation of the ideal \mathfrak{b}^* generated by the initial forms of the elements of an ideal \mathfrak{b} of a local ring, with respect to a certain ideal \mathfrak{a} .

The present paper has its origin in the effort of extending to a general situation the following result (due to Hironaka: [H₂]):

Let (A, \mathfrak{m}) be a local ring and $z \in \mathfrak{m} - \mathfrak{m}^2$; then the initial form z^* of z in the associated graded ring $G(\mathfrak{m})$ is a regular element if and only if z is regular in A and $(z) \cap \mathfrak{m}^{n+1} = (z) \cdot \mathfrak{m}^n$, for every integer n .

Really our paper investigates in a general way the relations between an ideal $\mathfrak{b} = (f_1, \dots, f_r)$ and the associated graded ideal \mathfrak{b}^* generated by the initial forms of the elements of \mathfrak{b} in $G_A(\mathfrak{a}) =$ graded ring with respect to the ideal \mathfrak{a} of the ring A .

We prove the following main results:

- 1—a necessary and sufficient condition for \mathfrak{b}^* to be generated by the initial forms of the f_i 's, valid for an arbitrary noetherian ring A ;
- 2—a necessary and sufficient condition for \mathfrak{b}^* to be generated by a regular sequence, valid for an arbitrary noetherian ring A ;
- 3—a condition like in 2, valid for a local ring A , and generalizing in a natural way Hironaka's result.

The paper contains also some other properties of the associated graded ideal (concerning height and minimal generating sets) and of $G_A(\mathfrak{a})$ (conditions to be Cohen-Macaulay: see also [S]).

Received September 16, 1977.

⁽¹⁾ The present paper has been written while both authors were members of CNR, section 3 of GNSAGA.

1. Form rings and ideals

If A is a noetherian ring with a unit element 1 and \mathfrak{a} any ideal in A , we denote by $G_A(\mathfrak{a})$ the graded A/\mathfrak{a} -algebra $\bigoplus_{n=0}^{\infty} \mathfrak{a}^n/\mathfrak{a}^{n+1}$ and call it the form ring of A relative to \mathfrak{a} (other term: associated graded ring).

Sometimes we will have to deal also with negative powers of the ideal \mathfrak{a} ; once for all $\mathfrak{a}^n = A$ if $n \leq 0$.

Given an element $a \in A$, we denote by $v(a)$ the largest integer n such that $a \in \mathfrak{a}^n$; if $a \in \bigcap_{n=1}^{\infty} \mathfrak{a}^n$ we set: $v(a) = \infty$. When $v(a) \neq \infty$ the residue class of a in $\mathfrak{a}^{v(a)}/\mathfrak{a}^{v(a)+1}$ is called the initial form of a and denoted by a^* . If $v(a) = \infty$, then we set: $a^* = 0$.

The definition of multiplication in $G_A(\mathfrak{a})$ shows that the two relations: $a^*b^* = (ab)^*$ and $a^*b^* \neq 0$ are equivalent, provided that a^* and $b^* \neq 0$.

Let now \mathfrak{b} be any ideal of A ; we shall denote by \mathfrak{b}^* the homogeneous ideal of $G_A(\mathfrak{a})$ generated by all the initial forms of the elements in \mathfrak{b} ; \mathfrak{b}^* is called the form ideal of \mathfrak{b} relative to \mathfrak{a} (or the associated graded ideal). For every integer n we have: \mathfrak{b}_n^* = set of homogeneous elements in \mathfrak{b}^* of degree $n = (\mathfrak{b} \cap \mathfrak{a}^n + \mathfrak{a}^{n+1})/\mathfrak{a}^{n+1}$; furthermore \mathfrak{b}^* is the kernel of a natural epimorphism of graded rings

$$\varphi: G_A(\mathfrak{a}) \rightarrow G_{A/\mathfrak{b}}(\mathfrak{b} + \mathfrak{a}/\mathfrak{b}),$$

which is homogeneous of degree 0.

The ideal \mathfrak{b}^* can be defined with respect to any other ideal \mathfrak{a} ; but, if \mathfrak{b} and \mathfrak{a} are comaximal (i.e. $\mathfrak{a} + \mathfrak{b} = A$) then $\mathfrak{b}^* = G_A(\mathfrak{a})$ and conversely (as one may easily check). Therefore we assume, once for all, that \mathfrak{a} and \mathfrak{b} are not comaximal. Of course both \mathfrak{a} and \mathfrak{b} are assumed to be proper.

Since A is noetherian, then $G_A(\mathfrak{a})$ is noetherian. In particular \mathfrak{b}^* is generated by the initial forms of a finite number of elements of \mathfrak{b} . However it is not generally true that, if $\mathfrak{b} = (f_1, \dots, f_r)$, then $\mathfrak{b}^* = (f_1^*, \dots, f_r^*)$.

The following theorem is just a necessary and sufficient condition for the equality:

THEOREM 1.1. *If \mathfrak{a} and $\mathfrak{b} = (f_1, \dots, f_r)$ are ideals of A , then $\mathfrak{b}^* = (f_1^*, \dots, f_r^*)$ in $G_A(\mathfrak{a})$ if and only if for all $n \geq 1$ the following equality holds:*

$$\mathfrak{a}^n \cap \mathfrak{b} = \sum_{i=1}^r \mathfrak{a}^{n-p_i} f_i$$

where $p_i = v(f_i)$, $i = 1, \dots, r$.

Proof. It is clear that $(f_1^*, \dots, f_r^*)_n = \left(\sum_{i=1}^r \mathfrak{a}^{n-p_i} f_i + \mathfrak{a}^{n+1} \right) / \mathfrak{a}^{n+1}$.

Hence, if $\mathfrak{b} \cap \mathfrak{a}^n = \sum_{i=1}^r \mathfrak{a}^{n-p_i} f_i$ for all $n \geq 1$, we have: $\mathfrak{b}^* = (f_1^*, \dots, f_r^*)$.

Conversely, if \mathfrak{b}^* is generated by the f_i^* 's, then we have: $\mathfrak{a}^n \cap \mathfrak{b} \subseteq \sum_{i=1}^r \mathfrak{a}^{n-p_i} f_i + \mathfrak{a}^{n+1}$ for all $n \geq 1$; it follows that $\mathfrak{a}^n \cap \mathfrak{b} = \bigcap_{t=1}^{\infty} \left(\sum_{i=1}^r \mathfrak{a}^{n-p_i} f_i + \mathfrak{a}^{n+t} \cap \mathfrak{b} \right)$. By the Artin-Rees lemma there exists an integer $q \geq 0$ such that $\mathfrak{a}^{n+t} \cap \mathfrak{b} = \mathfrak{a}^{n+t-q}(\mathfrak{a}^q \cap \mathfrak{b})$ for all $n + t \geq q$. Hence, if d is an integer such that $d \geq n - p_i$ for $i = 1, \dots, r$, we get the following equality:

$$\mathfrak{a}^n \cap \mathfrak{b} = \bigcap_{t > q - n + d} \left(\sum_{i=1}^r \mathfrak{a}^{n-p_i} f_i + \mathfrak{a}^{n+t-q}(\mathfrak{a}^q \cap \mathfrak{b}) \right) = \sum_{i=1}^r \mathfrak{a}^{n-p_i} f_i,$$

since, if $t \geq q - n + d$, then $\mathfrak{a}^{n+t-q}(\mathfrak{a}^q \cap \mathfrak{b}) \subseteq \mathfrak{a}^d \mathfrak{b} \subseteq \sum_{i=1}^r \mathfrak{a}^{n-p_i} f_i$.

Remark 1.2. Using [R, Rem. (3.7)] one can easily see (cf. [V]) that, if \mathfrak{a} and \mathfrak{b} are ideals of A such that $\mathfrak{a} + \mathfrak{b} \neq A$, then $h(\mathfrak{b}^*) \geq h(\mathfrak{b})$ at least when A is local, and certainly equality holds whenever $\mathfrak{a} \subseteq \mathfrak{b}$; moreover there are examples with strict inequality. Therefore the following result may have some interest:

PROPOSITION 1.3. *If \mathfrak{a} and $\mathfrak{b} = (f_1, \dots, f_r)$ are ideals of A such that $\mathfrak{a} + \mathfrak{b} \neq A$, then $h(\mathfrak{b}^*) \leq r$.*

Proof. Let \mathfrak{m} be a maximal ideal containing both \mathfrak{a} and \mathfrak{b} ; then we have: $h(\mathfrak{m}) - r \leq h(\mathfrak{m}/\mathfrak{b}) = h((\mathfrak{m}/\mathfrak{b})^*) = h(\mathfrak{m}^*/\mathfrak{b}^*) \leq h(\mathfrak{m}^*) - h(\mathfrak{b}^*) = h(\mathfrak{m}) - h(\mathfrak{b}^*)$ (in the preceding chain of inequalities we use the fact that, in the isomorphism $G_A(\mathfrak{a})/\mathfrak{b}^* \cong G_{A/\mathfrak{b}}(\mathfrak{a} + \mathfrak{b}/\mathfrak{b})$, $\mathfrak{m}^*/\mathfrak{b}^*$ and $(\mathfrak{m}/\mathfrak{b})^*$ are corresponding ideals).

Remark 1.4. If $h(f_1^*, \dots, f_r^*) = r$, then also $h(\mathfrak{b}^*) = r$. However the following example shows that the condition $h(f_1^*, \dots, f_r^*) = r$ does not imply the equality $\mathfrak{b}^* = (f_1^*, \dots, f_r^*)$.

EXAMPLE 1.5. Let A be the ring $k[[t^4, t^5, t^{11}]] = k[[X, Y, Z]]/(XZ - Y^3, YZ - X^4, Z^2 - X^3Y^2) = k[[x, y, z]]$, $\mathfrak{a} = (x, y, z)$, $\mathfrak{b} = (x)$; we have: $G_A(\mathfrak{a}) = k[T_1, T_2, T_3]/(T_1T_3, T_2T_3, T_3^2, T_2^4) = k[t_1, t_2, t_3]$; hence $h(\mathfrak{b}^*) = h(t_1) = 1$, but $y^3 \in \mathfrak{a}^3 \cap \mathfrak{b}$ and $y^3 \notin \mathfrak{a}^2x$ and so, by Theorem 1.1, $\mathfrak{b}^* \neq (x^*)$.

2. Regular sequences in $G_A(\mathfrak{a})$

In this section we consider a noetherian ring A and an ideal \mathfrak{a} in A ; if $f_1, \dots, f_r \in A$ we shall state some necessary and sufficient conditions for f_1^*, \dots, f_r^* to be a regular sequence in $G_A(\mathfrak{a})$ (always provided that \mathfrak{a} and (f_1, \dots, f_r) be not comaximal).

In the following we shall write $p_i = v(f_i)$ and $\mathfrak{b}_i = (f_1, \dots, f_i)$, for $i = 1, \dots, r$ ($\mathfrak{b}_0 = (0)$).

PROPOSITION 2.1. *If \mathfrak{a} and $\mathfrak{b} = (f_1, \dots, f_r)$ are ideals of A such that f_1^*, \dots, f_r^* is a $G_A(\mathfrak{a})$ -sequence, then $\mathfrak{b}^* = (f_1^*, \dots, f_r^*)$.*

Proof. We use induction on r . The case $r = 1$ is easy: if f^* is a non zero-divisor, then, for every g , $(fg)^* = f^*g^*$, which shows our claim. Assume now the theorem true for $r - 1$ and prove it for r .

If $a \in \mathfrak{a}^n \cap \mathfrak{b}$ let t be the greatest integer (if it exists, ∞ if it does not exist) such that $a \in \mathfrak{b}_{r-1} + f_r \mathfrak{a}^t$. So we can write $a = x + f_r y$, with $v(y) = t$, $x \in \mathfrak{b}_{r-1}$; if $t + p_r < n$ we have: $f_r y \in (\mathfrak{a}^n + \mathfrak{b}_{r-1}) \cap \mathfrak{a}^{t+p_r} \subseteq \mathfrak{a}^{t+p_r+1} + \mathfrak{b}_{r-1} \cap \mathfrak{a}^{t+p_r}$.

It follows that $f_r^* y^* \in \mathfrak{b}_{r-1}^*$; by our inductive hypothesis \mathfrak{b}_{r-1}^* is generated by the initial forms of the f_i 's, $i = 1, \dots, r - 1$, hence $y^* \in \mathfrak{b}_{r-1}^*$. Then $y \in \mathfrak{a}^{t+1} + \mathfrak{b}_{r-1} \cap \mathfrak{a}^t$, so that $a \in \mathfrak{b}_{r-1} + f_r \mathfrak{a}^{t+1}$, which is absurd. Therefore $t + p_r \geq n$, hence $a \in \mathfrak{a}^n \cap \mathfrak{b}_{r-1} + f_r \mathfrak{a}^{n-p_r}$. The conclusion follows immediately from the inductive assumption together with our Theorem 1.1.

Remark 2.2. The converse of Proposition 2.1 is false even if A is local. In fact, let $A = k[[X, Y]]/(XY) = k[[x, y]]$, $\mathfrak{b} = (x)$, $\mathfrak{a} = (x, y)$; then we have: $\mathfrak{a}^n \cap \mathfrak{b} = (x^n, y^n) \cap (x) = (x^n) + (y^n) \cap (x) = (x^n) = \mathfrak{a}^{n-1} \mathfrak{b}$. Hence by Theorem 1.1 $\mathfrak{b}^* = (x^*)$, but $G_A(\mathfrak{a}) = k[T_1, T_2]/(T_1 T_2) = k[t_1, t_2]$ and $x^* = t_1$ is a zero divisor in $G_A(\mathfrak{a})$.

In the following we shall denote by \bar{I} the topological closure of an ideal I with respect to the \mathfrak{a} -adic topology.

THEOREM 2.3. *Let \mathbf{a} and $\mathbf{b} = (f_1, \dots, f_r)$ be ideals of the noetherian ring A . Then the following facts are equivalent:*

- (i) (f_1^*, \dots, f_r^*) is a $G_A(\mathbf{a})$ -sequence;
- (ii) for each $i = 1, \dots, r, \mathbf{b}_{i-1}: f_i \subseteq \bar{\mathbf{b}}_{i-1}$ and $\mathbf{b}_i \cap \mathbf{a}^n = \sum_{j=1}^i \mathbf{a}^{n-p_j} f_j$, for all $n \geq 1$.

Proof. (i) \Rightarrow (ii). For Proposition 2.1 it is enough to show that $\mathbf{b}_{i-1}: f_i \subseteq \bar{\mathbf{b}}_{i-1}$. Let $a \in \mathbf{b}_{i-1}: f_i$ with $v(a) = n$; then $a^* f_i^* \in \mathbf{b}_{i-1}^* = (f_1^*, \dots, f_{i-1}^*)$, hence $a^* \in \mathbf{b}_{i-1}^*$. It follows that $a \in \mathbf{b}_{i-1} + \mathbf{a}^{n+1} \cap (\mathbf{b}_{i-1}: f_i)$; repeating the argument we see that $a \in \bar{\mathbf{b}}_{i-1}$.

(ii) \Rightarrow (i). Conversely, since $\mathbf{a} + \mathbf{b} \neq A$ and $\mathbf{b}_{i-1}: f_i \subseteq \bar{\mathbf{b}}_{i-1}$ for each $i = 1, \dots, r$, we have $p_i < \infty$ for each $i = 1, \dots, r$. In fact, assume that $p_i = \infty$ for some i ; then there exists $a \in \mathbf{a}$ such that $(1 - a)f_i = 0$. Thus $1 - a \in \bar{\mathbf{b}}_{i-1}$ and then $(1 - a)(1 - a') \in \mathbf{b}_{i-1}$, with $a' \in \mathbf{a}$; but since $\mathbf{a} + \mathbf{b} \neq A$, this is a contradiction.

Now let $a^* f_i^* \in (f_1^*, \dots, f_{i-1}^*)$ with $v(a) = n$; then $a f_i \in \mathbf{b}_{i-1} + \mathbf{a}^{n+p_i+1}$, hence $b = a f_i + \sum_{j=1}^{i-1} a_j f_j \in \mathbf{b}_i \cap \mathbf{a}^{n+p_i+1}$.

We can write $b = \sum_{j=1}^i b_j f_j$ with $b_j \in \mathbf{a}^{n+p_i+1-p_j}$, whence we deduce that $a - b_i \in \mathbf{b}_{i-1}: f_i$ and from this it follows that $a \in \mathbf{a}^{n+1} + \bar{\mathbf{b}}_{i-1}$.

Finally $a \in \mathbf{a}^{n+1} + \mathbf{b}_{i-1} \cap \mathbf{a}^n$ and this proves that $a^* \in \mathbf{b}_{i-1}^* = (f_1^*, \dots, f_{i-1}^*)$.

COROLLARY 2.4. *Let A be a local ring and I, \mathbf{a} ideals of A , such that I^* is generated by a $G_A(\mathbf{a})$ -sequence. Then I is generated by an A -sequence.*

Proof. Let I^* be generated by g_1, \dots, g_r , where the g_i 's form a regular sequence. Since all the minimal generating sets of I^* have the same number of elements, we can write $I^* = (f_1^*, \dots, f_r^*)$ with $f_i \in I$. Now $gr(I^*) = r$, hence using the homology of Koszul complex we get $H_1(f_1^*, \dots, f_r^*; G_A(\mathbf{a})) = 0$. From this it follows that f_1^*, \dots, f_r^* is a $G_A(\mathbf{a})$ -sequence (see [A-B], Prop. 2.8); hence, by Theorem 2.3, f_1, \dots, f_r is an A -sequence. Furthermore, since $I^* = (f_1^*, \dots, f_r^*)$, we get: $(f_1, \dots, f_r) \subseteq I \subseteq \overline{(f_1, \dots, f_r)} = (f_1, \dots, f_r)$.

Remark 2.5. If f_1^*, \dots, f_r^* form a regular sequence, it is not necessarily true that f_1, \dots, f_r form also an A -sequence, unless $I = \bar{I}$ for

every ideal I contained in \mathfrak{b} . In fact let $A = k[x, y, z] = k[X, Y, Z]/(XZ, X - XY)$, $\mathfrak{a} = (y)$, $f = yz$; then, since y is not a 0-divisor in A , we have: $G_A(\mathfrak{a}) = (A/\mathfrak{a})[T] = k[Z, T]$ which is a domain. Therefore $\bar{f} \in \mathfrak{a}/\mathfrak{a}^2$ is not a 0-divisor in $G_A(\mathfrak{a})$, but $xf = 0$.

PROPOSITION 2.6. *Let \mathfrak{a} and $\mathfrak{b} = (f_1, \dots, f_r)$ be two ideals of A such that f_1, \dots, f_r is an A -sequence and $\mathfrak{a}^n \cap \mathfrak{b} = \sum_{i=1}^r \mathfrak{a}^{n-p_i} f_i$ for all $n \geq 1$. Suppose either $\mathfrak{b} \subseteq \mathfrak{a}$ or A is local.*

Then $\mathfrak{a}^n \cap \mathfrak{b}_i = \sum_{j=1}^i \mathfrak{a}^{n-p_j} f_j$, for each $i = 1, \dots, r$ and for all $n \geq 1$; thus f_1^, \dots, f_r^* is a $G_A(\mathfrak{a})$ -sequence.*

Proof. It is enough to show that $\mathfrak{a}^n \cap \mathfrak{b}_{r-1} = \sum_{j=1}^{r-1} \mathfrak{a}^{n-p_j} f_j$, for all $n \geq 1$. Let $a = \sum_{i=1}^{r-1} a_i f_i$ be an element of \mathfrak{a}^n ; then $a = \sum_{i=1}^r b_i f_i$, where $b_i \in \mathfrak{a}^{n-p_i}$ and we get: $b_r \in \mathfrak{a}^{n-p_r} \cap \mathfrak{b}_{r-1}$. Thus we have: $\mathfrak{a}^n \cap \mathfrak{b}_{r-1} \subseteq \sum_{j=1}^{r-1} \mathfrak{a}^{n-p_j} f_j + f_r(\mathfrak{a}^{n-p_r} \cap \mathfrak{b}_{r-1})$. Let m be an integer such that $m \leq p_i$ for each $i = 1, \dots, r-1$. If $n \leq m + p_r$, we have that $\mathfrak{b}_{r-1} \subseteq \mathfrak{a}^{n-p_r}$ and $f_r \in \mathfrak{a}^{n-p_i}$ for each $i = 1, \dots, r-1$; hence $\mathfrak{a}^n \cap \mathfrak{b}_{r-1}$ is contained in $\sum_{j=1}^{r-1} \mathfrak{a}^{n-p_j} f_j + f_r \mathfrak{b}_{r-1} \subseteq \sum_{j=1}^{r-1} \mathfrak{a}^{n-p_j} f_j$. Therefore we may assume that n is greater than $m + p_r$ and also that $\mathfrak{a}^t \cap \mathfrak{b}_{r-1} = \sum_{j=1}^{r-1} \mathfrak{a}^{t-p_j} f_j$, for all $t \leq n-1$. If $\mathfrak{b} \subseteq \mathfrak{a}$, then $p_r > 0$; if A is local and $p_r = 0$, then we have $\mathfrak{a}^n \cap \mathfrak{b}_{r-1} \subseteq \sum_{j=1}^{r-1} \mathfrak{a}^{n-p_j} f_j + f_r(\mathfrak{a}^n \cap \mathfrak{b}_{r-1})$, hence $\mathfrak{a}^n \cap \mathfrak{b}_{r-1} = \sum_{j=1}^{r-1} \mathfrak{a}^{n-p_j} f_j$. If $p_r \geq 1$, since $n - p_r \leq n - 1$, we have $\mathfrak{a}^n \cap \mathfrak{b}_{r-1} \subseteq \sum_{j=1}^{r-1} \mathfrak{a}^{n-p_j} f_j + f_r \left(\sum_{j=1}^{r-1} \mathfrak{a}^{n-p_r-p_j} f_j \right) = \sum_{j=1}^{r-1} \mathfrak{a}^{n-p_j} f_j$, and this completes the proof.

COROLLARY 2.7. *If \mathfrak{a} and $\mathfrak{b} = (f_1, \dots, f_r)$ are ideals of a local ring A , then f_1^*, \dots, f_r^* form a $G_A(\mathfrak{a})$ -sequence if and only if f_1, \dots, f_r form an A -sequence and moreover $\mathfrak{a}^n \cap \mathfrak{b} = \sum_{i=1}^r \mathfrak{a}^{n-p_i} f_i$ for all $n \geq 1$.*

Remark 2.8. The following example justifies the hypotheses of the above proposition. Let A be the ring $k[X, Y, Z, T]/(XT - Y^2, T - Y - TZ) = k[x, y, z, t]$, $\mathfrak{a} = (x, y)$, $f_1 = x$ and $f_2 = z$; then it is easy to see that

f_1, f_2 form a regular sequence and $p_1 = 1, p_2 = 0$. We have: $xt \in \mathfrak{a}^2 \cap (x)$, but $xt \notin \mathfrak{a}x$; on the other hand $\mathfrak{a} \cap (x, z) = (x) + \mathfrak{a} \cap (z) = (x) + \mathfrak{a}z$ and, if $n \geq 2$, then $\mathfrak{a}^n \cap (x, z) = \mathfrak{a}^n = \mathfrak{a}^{n-1}x + (y^n) = \mathfrak{a}^{n-1}x + \mathfrak{a}^nz$ since $y^n = xy^{n-1} + y^nz$.

Remark 2.9. The results of the present section extend to a quite general situation the theorem proved by Hironaka ([H₂], Prop. 6) for a local ring (A, \mathfrak{m}) , when $\mathfrak{a} = \mathfrak{m}$ and $\mathfrak{b} = (f)$ = principal ideal, generated by an element in $\mathfrak{m} - \mathfrak{m}^2$.

3. Applications

In this section we discuss some applications of the preceding results.

PROPOSITION 3.1. *Let \mathfrak{a} and $\mathfrak{b} = (f_1, \dots, f_r)$ be ideals of A such that $\mathfrak{b} \subseteq \mathfrak{a}$, f_1, \dots, f_r is an A -sequence and $\mathfrak{a}\mathfrak{b} = \mathfrak{a}^2$. Then the initial forms of the f_i 's form a $G_A(\mathfrak{a})$ -sequence.*

Proof. If $f_i \in \mathfrak{a}^2 = \mathfrak{a}\mathfrak{b}$ then we would get a relation of the form

$$a_1f_1 + \dots + (1 + a_i)f_i + \dots + a_rf_r = 0, \quad a_j \in \mathfrak{a}(\forall j).$$

But since f_1, \dots, f_r is an A -sequence, in any relation $\sum x_jf_j = 0$ all the coefficients x_j must lie in the ideal (f_1, \dots, f_r) . This is well known and easy to prove by induction on r . Thus $f_i \notin \mathfrak{a}^2$ and we have $p_1 = \dots = p_r = 1$. Therefore it is enough to prove, by Proposition 2.6, that $\mathfrak{a}^n \cap \mathfrak{b} = \mathfrak{a}^{n-1}\mathfrak{b}$. This is true for $n = 1$; if $n \geq 2$ we have $\mathfrak{a}^n = \mathfrak{a}^{n-1}\mathfrak{b}$, hence $\mathfrak{a}^n \cap \mathfrak{b} = \mathfrak{a}^{n-1}\mathfrak{b} \cap \mathfrak{b} = \mathfrak{a}^{n-1}\mathfrak{b}$.

Remark 3.2. An interesting situation in which we can apply the above proposition is the following: let (A, \mathfrak{m}) be a local ring of dimension r which is Cohen-Macaulay, with embedding dimension m and multiplicity e ; then one can show ([S], Theorem 1) that $m \leq e + r - 1$ and the equality holds if and only if there is an A -sequence f_1, \dots, f_r in \mathfrak{m} such that $\mathfrak{m}^2 = \mathfrak{m}(f_1, \dots, f_r)$. The latter equality is exactly our condition on the ideals $\mathfrak{a} = \mathfrak{m}$ and $\mathfrak{b} = (f_1, \dots, f_r)$.

Remark 3.3. The results of the present paper can be used to give a new and simplified proof of ([V], Theorem 3.2), i.e. of the following claim:

Let A be a Cohen-Macaulay ring and let a_1, \dots, a_s be a regular

sequence, $I = (a_1, \dots, a_s)$, t an integer ≥ 1 . Then $G_A(\mathbf{a})$ is Cohen-Macaulay if $\mathbf{a} = I^t$.

Proof. As in [V] we may assume that A is a r -dimensional local ring with maximal ideal \mathfrak{m} . Let $a_1, \dots, a_s, f_{s+1}, \dots, f_r$ be a maximal A -sequence in \mathfrak{m} and let $J = (f_{s+1}, \dots, f_r)$, $f_i = a_i^t$, for each $i = 1, \dots, s$ and $\mathbf{b} = (f_1, \dots, f_r)$. Since f_1, \dots, f_s is a regular sequence modulo J we have, by [V], Lemma 2.1, that $\mathfrak{a}^n \cap \mathbf{b} \subseteq \mathfrak{a}^{n-1}(f_1, \dots, f_s) + J$ for all $n \geq 1$; furthermore, since f_{s+1}, \dots, f_r is a regular sequence modulo I , hence modulo \mathfrak{a}^n for all $n \geq 1$, by [R-V], Lemma 1.1, we get: $\mathfrak{a}^n \cap J = \mathfrak{a}^n J$. From this it follows that $\mathfrak{a}^n \cap \mathbf{b} \subseteq \mathfrak{a}^n \cap (\mathfrak{a}^{n-1}(f_1, \dots, f_s) + J) = \mathfrak{a}^{n-1}(f_1, \dots, f_s) + \mathfrak{a}^n \cap J = \mathfrak{a}^{n-1}(f_1, \dots, f_s) + \mathfrak{a}^n J$; thus the f_i^* 's form a $G_A(\mathbf{a})$ -sequence by Proposition 2.6. Since $\dim G_A(\mathbf{a}) = r$, by [M-R] this is enough to prove that $G_A(\mathbf{a})$ is Cohen-Macaulay.

We conclude the paper trying to compare the initial forms of a set of elements with respect to two different ideals \mathbf{a} and I .

First of all, it is easy to see that the initial form of the same element with respect to two different ideals may or may not be a 0-divisor; for instance, if $A = k[[x, y, z]] = k[[X, Y, Z]]/(XY - Z^2)$, $\mathbf{a} = (x, y, z)$, $I = (x, z)$, then it is clear that the initial form of x relative to \mathbf{a} is a non 0-divisor, while the initial form with respect to I is a 0-divisor. Therefore $(x)^* = (x^*)$ in $G_A(\mathbf{a})$ (Proposition 2.1); on the contrary $(x)^*$ in $G_A(I)$ is not generated by the initial form of x , because of Theorem 1.1.

In the following proposition we denote by f^* the initial form with respect to \mathbf{a} and by f^0 the initial form with respect to I .

PROPOSITION 3.4. *Let $I \subseteq \mathbf{a}$ be ideals of A and let f_1, \dots, f_r be elements of I such that $v_I(f_i) = v_{\mathbf{a}}(f_i)$ for each i . Assume that f_1^*, \dots, f_r^* form a $G_A(\mathbf{a})$ -sequence. Then f_1^0, \dots, f_r^0 form a minimal base of the ideal (f_1^0, \dots, f_r^0) of $G_A(I)$.*

Proof. By [A-B], Corollary 2.9, f_1^*, \dots, f_r^* is a $G_A(\mathbf{a})$ -sequence in any order. Now if $f_r^0 = \sum_{i=1}^{r-1} a_i^0 f_i^0$, let $a = \sum_{i=1}^{r-1} a_i f_i$ and $p = v_{\mathbf{a}}(f_r) = v_I(f_r)$; then $f_r = a + b$, where $b \in I^{p+1}$. Hence $a \in \mathfrak{a}^p$ and $a \notin \mathfrak{a}^{p+1}$; it follows that $f_r^* = a^* \in (f_1, \dots, f_{r-1})^* = (f_1^*, \dots, f_{r-1}^*)$, which is absurd.

REFERENCES

- [A-B] M. Auslander and D. Buchsbaum, Codimension and multiplicity, *Ann. of Math.*, **68** (1958), 625–657.
- [H₁] H. Hironaka, Resolution of singularities of an algebraic variety over a field of characteristic, 0. *Ann. of Math.*, **79** (1964), 109–326.
- [H₂] H. Hironaka, Certain numerical characters of singularities, *J. Math. Kyoto Univ.*, **10** (1970), 151–187.
- [M-R] J. Matijevich and P. Roberts, A conjecture of Nagata on graded Cohen-Macaulay rings, *J. Math. Kyoto Univ.*, **14** (1974), 125–128.
- [R] L. J. Ratiff, Jr., On quasi unmixed local domains, the altitude formula, and the chain condition for prime ideals (II), *Amer. J. Math.*, **92** (1970), 99–144.
- [R-V] L. Robbiano and G. Valla, On normal flatness and normal torsion freeness, *J. of Algebra*, **43** (1976), 552–560.
- [S] J. Sally, On the associated graded ring of a local Cohen-Macaulay ring, *J. Math. Kyoto Univ.*, **17** (1977), 19–21.
- [V] G. Valla, Certain graded algebras are always Cohen-Macaulay, *J. of Algebra*, **42** (1976), 537–548.

Istituto matematico-Politecnico di Torino
Istituto matematico-Università di Genova