¹Psychiatry And Legal Medicine, Autonomous University of Barcelona, Barcelona, Spain; ²Inad-parc De Salut Mar, Hospital del Mar, Barcelona, Spain and ³Institut De Neuropsiquiatria I Addiccions (inad), Hospital del Mar, Barcelona, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1943

Introduction: Meta-analytic evidence suggests that migrants have higher risk for psychotic disorders. Likewise, growing evidence relate developmental trauma (emotional, sexual, physical abuse and neglect in childhood or adolescence) as a causal factor for psychotic symptoms. However, few studies examine developmental trauma in migrant populations.

Objectives: The aim of this study is to describe and compare developmental trauma exposure prevalence between immigrant and non-immigrant psychotic patients in Barcelona.

Methods: Patients who have presented, according DSM-V criteria, one or more non-affective psychotic episodes, were recruited in Acute and Chronic inpatients units at Hospital del Mar (Barcelona), leading to a total sample of 77 patients. Demographic characteristics of patients, clinical data and main pharmacological treatment were recorded through a questionnaire. Developmental trauma exposure was assessed by Childhood Trauma Questionnaire (CTQ). Comparative analysis was performed with IBM SPSS using Chi-Square Test and t-Student test.

Results: From a total of 77 patients, 43 were immigrants and 34 were non-immigrants. Exposure to traumatic events showed significant differences between immigrants and non-immigrant in Child emotional abuse (64,4% immigrants, 35,3% non-immigrant), Child physical abuse (51,2% immigrants, 14,7% non-immigrant), Child Sexual Abuse (41,9% immigrants, 11,8% non-immigrant) and physical neglect (62,8% immigrants, 26,5% non-immigrant). Emotional neglect exposure was no significant between both groups. Total mean CTQ score was 37,53 in immigrants group and 52,60 in non-immigrant group.

Conclusions: According to our results, there are important and significant differences in developmental trauma exposure between immigrant and non-immigrant psychotic patients. These results should be considered by clinicians in order to design assessment program for this population.

Disclosure: No significant relationships.

Keywords: transcultural psychiatry; trauma; psychosis; migration psychiatry

EPV0384

Migration: A risk factor for psychosis?

E. Spaho¹, V. Alikaj^{1,2}, E. Dashi^{1*} and V. Skendi¹

¹Department Of Neuroscience, University Hospital Center "Mother Theresa", Tirane, Albania and ²Faculty Of Medicine, Tirana Medical University Albania, Tirane, Albania

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1944

Introduction: Emigration is a complex process of social changing through which an individual moves from a certain cultural environment/context to another, aiming to achieve persistent or long-term residency, causing distress. There is sustainable evidence that incidence of all forms of psychosis is higher in migrants.

Objectives: This study aims to gather data of other research conducted in the field according to emigration as a risk factor for development of different psychosis.

Methods: Scientific articles searched in MEDLINE, regarding the incidence of mental disorders in different emigrant populations, for the period 1995 - 2015.

Results: The average relative risk of schizophrenia and of other psychosis occurrence among first generation emigrants was 2.7 (95% confidence interval [CI]=2.3-3.2). Statistical analysis performed among studies of first and second generation of emigrants, and among studies which don't make difference between generations, results in a relative risk of 2.9 (95% CI=2.5-3.4) of mental illness.

Conclusions: The data presented in this study emphasize the impact of migration on central symptoms of schizophrenia. Emigration process, cultural and social adaptation, play an important role on the individual mental health.

Disclosure: No significant relationships.

Keywords: Risk factors; psychosis; schizophrenia; Migration

EPV0388

Roles of trait resilience, flexibility, and volitional self-control in social adaptation: An fMRI study

S. Tei^{1,2,3*} and J. Fujino^{1,4}

¹Department Of Psychiatry, Kyoto University, Kyoto, Japan; ²Institute Of Applied Brain Sciences, Waseda University, Saitama, Japan; ³School Of Human And Social Sciences, Tokyo International University, Saitama, Japan and ⁴Department Of Psychiatry And Behavioral Sciences, Tokyo Medical and Dental University, Tokyo, Japan

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1945

Introduction: Resilience and cognitive flexibility are considered as pre-adaptive traits that help individuals to deal with environmental and social distress. Although they may alleviate emotional impulses and/or support volitional self-control (VSC), these cognitive mechanisms remain insufficiently explored.

Objectives: To better understand resilience and flexibility, we built a mechanistic framework to explain variations in socially adaptive responses under distressing situations (cooperation dilemma) using economic and social decision-making paradigms.

Methods: Twenty-four university students (7 females) were enrolled. We used ego-resiliency (ER) and Machiavellian (Mach) questionnaires to measure resilience and flexibility, and applied third-party punishment (TPP) and ultimatum game (UG), as well as moral dilemma (MD) tasks to derive VSC-associated brain activity using 3T-functional magnetic resonance imaging. Mediation analysis was used to investigate whether these pre-adaptive trait levels explain cooperative decision-making (invested sum in TPP and acceptance rate of unfair offers in UG), together with VSC-associated brain activity during MD. The regions of interest included the orbitofrontal cortex (OFC), dorsolateral prefrontal cortex (DLPFC), and temporoparietal junction (TPJ).

Results: Pre-adaptive traits were a statistically significant mediator for two different models. There was an indirect effect of 1) ER on the relationship between OFC activity strength and TPP scores, and 2) Mach on the relationship between DLPFC/TPJ and UG ($p < 0.05$).