

2021 BIDMC Gram-Positive Antibigram

How to Use:

Click on the desired organism to highlight the antibiotic sensitivities below.

To select for multiple organisms, hold ctrl

Gram-Positive Organisms	Occurrence
BETA STREPTOCOCCUS GROUP B	30
ENTEROCOCCUS SPP (including E. faecalis & E. faecium)	821
ENTEROCOCCUS FAECALIS	312
ENTEROCOCCUS FAECIUM	202
ENTEROCOCCUS SPP	307
STAPHYLOCOCCUS AUREUS (all)	716
STAPHYLOCOCCUS AUREUS-MRSA	242
STAPHYLOCOCCUS AUREUS-MSSA	475
STAPHYLOCOCCUS, COAGULASE NEGATIVE	383
STREPTOCOCCUS ANGINOSUS (MILLER) GROUP	85

Methodology: Only the first isolate per patient per year is included for the antibiogram year in accordance with CLSI m39-A4E recommendations. Only cultures for diagnostic purposes are included. The BIDMC Antibigram includes isolates from all inpatient units & emergency department.

Antibiotic name Gram-Positive Organisms	AMPCILLIN n %	CLINDAMYCIN n %	DAPTOMYCIN ^A n %	GENTAMICIN n %	LEVOFLOXACIN n %	LINEZOLID n %	NITROFURANTOIN [*] n %	OXACILLIN n %	PENICILLIN G n %	TETRACYCLINE n %	TRIM/SULFA n %	VANCOMYCIN n %
BETA STREPTOCOCCUS GROUP B		30 40%							30 100%			30 100%
ENTEROCOCCUS SPP (including E. faecalis & E. faecium)	819 67%		173 95%			297 99%	504 75%		316 61%	515 20%		821 67%
ENTEROCOCCUS FAECALIS	312 100%		80 96%			54 100%	170 99%		143 100%	175 18%		312 88%
ENTEROCOCCUS FAECALIS-VRE	36 100%					36 100%						36 0%
ENTEROCOCCUS FAECIUM-VSE	276 100%		68 97%				151 99%		126 100%	154 19%		276 100%
ENTEROCOCCUS FAECIUM	202 8%		92 93%			159 99%	84 24%		118 9%	90 19%		202 24%
ENTEROCOCCUS FAECIUM-VRE	153 1%		73 95%			152 99%	65 25%		88 0%	70 17%		154 0%
ENTEROCOCCUS FAECIUM-VSE	50 30%								31 35%			49 100%
ENTEROCOCCUS SPP	305 73%					84 98%	250 76%		55 71%	250 23%		307 73%
STAPHYLOCOCCUS AUREUS (all)		666 68%	73 100%	716 99%	704 75%	46 100%	50 98%	716 66%	569 91%	576 97%	716 97%	252 100%
STAPHYLOCOCCUS AUREUS-MRSA		223 56%	61 100%	241 98%	232 38%	38 100%		242 0%	186 85%	241 94%	241 100%	
STAPHYLOCOCCUS AUREUS-MSSA		443 75%		475 99%	472 94%		31 97%	475 100%	383 94%	475 99%		
STAPHYLOCOCCUS, COAGULASE NEGATIVE		305 57%	48 100%	383 81%	375 60%	32 100%	72 99%	383 40%		217 85%		383 100%
STREPTOCOCCUS ANGINOSUS (MILLER) GROUP		85 69%							84 86%			85 100%

Nitrofurantoin* for urine isolates only

^A % susceptible for E. faecium includes susceptible dose dependent for daptomycin

Streptococcus pneumoniae isolates over the past 2 years									
Antibiotic	Ceftriaxone			Penicillin			Penicillin (oral)		
Type	n	MIC	%	n	MIC	%	n	MIC	%
meningitis	48	<=0.5	90.0%	48	<=0.06	60.0%			
non-meningitis	48	<=0.5	90.0%	48	<=2	96.0%	48	<=0.06	60.0%

Presentation Type:

Poster Presentation - Poster Presentation

Subject Category: Antibiotic Stewardship

Creating an electronic antibiogram using visualization software: Easily updatable and removes the need for yearly manual review

Ashley Dauphin; Christopher McCoy; Robert Bowden; Matthew Lee; Howard Gold and Ryan Chapin

Background: Previously, our hospital manually built a static antibiogram from a surveillance system (VigiLanz) culture report. In 2019, a collaboration between the antimicrobial stewardship team (AST) and the infection control (IC) team set out to leverage data automation to create a dynamic antibiogram. The goal for the antibiogram was the ability to easily distribute and update for hospital staff, with the added ability to perform advanced tracking and surveillance of organism and drug susceptibilities for AST and IC. By having a readily available, accurate, and Clinical and Laboratory Standards Institute (CLSI)-compliant antibiogram, clinicians have the best available data on which to base their empiric antibiotic decisions. **Methods:** First, assessment of required access to hospital databases and selection of a visualization software (MS Power BI) was performed. Connecting SQL database feeds to Power BI enabled creation of a data model using DAX and M code to comply with the CLSI, generating the first isolate per patient per year. Once a visual antibiogram was created, it was validated against compiled antibiograms using data from the microbiology laboratory middleware (bioMerieux, Observa Integrated Data Management Software). This validation process uncovered some discrepancies between the 2 reference reports due to cascade reporting of susceptibilities. The Observa-derived data were used as the source of truth. The antibiogram prototype was presented to AST/IC members, microbiology laboratory leadership, and other stakeholders to assess functionality. **Results:** Following feedback and revisions by stakeholders, the new antibiogram was published on a hospital-wide digital platform (Fig. 1). Clinicians may view the antibiogram at any time on desktops from a firewall (or password)-protected intranet. The antibiogram view defaults to the current calendar year and users may interact with the antibiogram rows and columns without disrupting the integrity of the background databases or codes. Each year, simple refreshing of the Power BI antibiogram and

changing of the calendar year allows us to easily and accurately update the antibiogram on the hospital-wide digital platform. **Conclusions:** This interdisciplinary collaboration resulted in a new dynamic, CLSI-compliant antibiogram with improved usability, increased visibility, and straightforward updating. In the future, a mobile version of the antibiogram may further enhance accessibility, bring more useful information to providers, and optimize AST/IC guidelines and education.

Disclosures: None

Antimicrobial Stewardship & Healthcare Epidemiology 2023;3(Suppl. S2):s34

doi:10.1017/ash.2023.262

Presentation Type:


Poster Presentation - Poster Presentation

Subject Category: Antibiotic Stewardship

Identifying the relationship between hospital rurality and antibiotic overuse

Hannah Hardin; Valerie Vaughn; Andrea White; Jennifer Horowitz; Elizabeth McLaughlin; Julia Szymczak; Lindsay Petty; Anurag Malani; Scott Flanders and Tejal Gandhi

Background: Antibiotic overuse and the resulting patient outcomes span all hospitals. However, although antibiotic stewardship can improve antibiotic use, effective stewardship programs require expertise and an infrastructure that are not present in all hospitals. Rural hospitals have less access to resources, infectious disease expertise, and participation in academic research. Thus, we compared antibiotic overuse at discharge between rural and nonrural hospitals for patients diagnosed with community-associated pneumonia (CAP) or urinary tract infection (UTI)—the 2 most common hospital infections. **Methods:** To determine whether antibiotic overuse at discharge was higher among rural versus nonrural hospitals, we analyzed data from a 41-hospital prospective cohort of patients treated for CAP or UTI between July 1, 2017, and July 30, 2019, in Michigan. Antibiotic overuse was defined as treatment that was unnecessary (ie, patient did not have an infection), excessive (ie, duration >4 days for CAP), or included suboptimal fluoroquinolone use (ie, safer alternative available). Overuse was determined based on patient risk



factors, symptoms, allergies, diagnostic results, and time to stability. Hospital rurality was defined using the Rural–Urban Continuum Codes (RUCC) score. We defined rural as a score ≥ 4 and very rural as a score of 7–9. We used *t* tests to compare the mean percentage of patients with antibiotic overuse at discharge between nonrural and rural (and very rural) hospitals. **Results:** Across 41 hospitals, we included 23,449 patients with CAP or UTI. There were 5 rural (and 3 very rural) hospitals with 2,039 (and 1,082) patients. Antibiotic overuse at discharge was present in 43.1% of patient cases in nonrural hospitals, 52.5% in rural hospitals ($P = .04$ vs nonrural) and 58.1% in very rural hospitals ($P = .007$ vs nonrural). Compared to nonrural hospitals, the mean percentage of cases with antibiotic overuse at discharge in rural hospitals was 9.4% higher (15.1% higher in very rural hospitals). Results were similar in a subgroup analysis of only patients with UTI (47.0% in rural vs 37.5% in nonrural, mean difference, 9.5%; $P = .03$) but were not statistically significant in patients with CAP (53.8% vs 48.0%, respectively; mean difference, 5.8%; $P = 0.23$). **Conclusions:** In this retrospective study, rural hospitals—especially very rural hospitals, had higher rates of antibiotic overuse at discharge than nonrural hospitals. Our findings suggest that antibiotic stewardship interventions tailored toward the unique differences in infrastructure, resources, and needs of rural hospitals are essential to community health.

Disclosures: None

Antimicrobial Stewardship & Healthcare Epidemiology 2023;3(Suppl. S2):s34–s35

doi:10.1017/ash.2023.263

Presentation Type:

Poster Presentation - Poster Presentation

Subject Category: Antibiotic Stewardship

Validation of antibiotic stewardship metrics for genitourinary infection management in Veterans Affairs outpatient settings

Jordan Braunfeld; Matthew Samore; Jacob Crook; McKenna Nevers; Kelly Echevarria; Ben Brintz; Matthew Goetz and Karl Madaras-Kelly

Background: Diagnosis and management of suspected urinary tract infection (UTI) in outpatient settings has been shown to be suboptimal. We previously developed a set of stewardship metrics for UTIs based on electronic health record (EHR) data (*Antimicrobial Stewardship & Healthcare*

Tier	Antibiotic Prescribed	Cases reviewed (N)	Age (SD)	Female (%)	Reviewer diagnosed GU infection, N (%)	Reviewer recommended antibiotics, N (%)
1/2	Yes	33	63.1 (15)	8 (24.2)	23 (69.7)	20 (60.6)
	No	45	64.7 (18.3)	6 (13.3)	3 (6.7)	5 (11.1)
3	Yes	29	71.4 (13.1)	0	2 (6.9)	4 (13.8)
	No	41	71.5 (11.6)	4 (9.8)	1 (2.4)	0

Epidemiology 2022;2 suppl 1:S5–S6. doi:10.1017/ash.2022). A tier-based approach was used to more fully capture antibiotic use associated with genitourinary (GU) symptoms and diagnoses. Herein we report a preliminary analysis of validity and reliability of these metrics based on chart abstraction. **Methods:** The study cohort consisted of patients who visited Veterans Affairs emergency departments or primary care clinics between 2015 and 2022 and who had a GU diagnosis based on *International Classification of Disease, Tenth Revision* (ICD-10) codes, divided into 3 categories: tier 1 (antibiotics always indicated), tier 2 (antibiotics sometimes indicated), and tier 3 (antibiotics not indicated). Visits related to urological procedures, nontarget settings, or concomitant non-GU infections were excluded. Cases were randomly sampled for manual review from within 8 strata based on tier, use of antibiotics, and visit type. An infectious disease physician and pharmacist abstracted charts using a standardized data-collection instrument. Clinical judgments regarding diagnosis and treatment were recorded on a Likert scale without knowledge of how the patient was managed. The intraclass correlation coefficient (ICC) was used to estimate interrater reliability. **Results:** To date, 148 cases have been reviewed (50 by both reviewers). Mean (SD) age was 67.5 (15.3) years and 12.2% were female. In a majority of tier 1 and 2 visits in which antibiotics were given, the reviewers found evidence for GU infection (69.7%) and favored prescribing of antibiotics (60.6%) (Table). In contrast, most patients in the tier 3 category who received antibiotics were judged to have noninfectious conditions (eg, benign prostatic hypertrophy) and to not require antibiotics. In the subset of records examined by both reviewers, the interrater reliability of judgments of whether antibiotics were warranted was good (ICC = .704). **Conclusions:** This preliminary validation provides support for a tier-based approach for stewardship metrics for GU conditions that relies upon electronic data to identify patients for whom antibiotics are generally not indicated.

Disclosures: None

Antimicrobial Stewardship & Healthcare Epidemiology 2023;3(Suppl. S2):s35

doi:10.1017/ash.2023.264

Presentation Type:

Poster Presentation - Poster Presentation

Subject Category: Antibiotic Stewardship

Understanding clinician perspectives on antibiotic associated adverse events to inform feedback

Jerald Cherian; George Jones; Taylor Helsel; Zunaira Virk; Alejandra Salinas; Suzanne Grieb; Sara Keller; Pranita Tamma and Sara Cosgrove

Background: Feedback regarding antibiotic-associated adverse events (ABX-AEs) may assist clinicians with antibiotic decision making. We

Table. Categorization of Antibiotic-Associated Adverse Events by Degree of Clinical Concern

Prespecified Categorization	Votes (n)*		
	Very Concerning	Moderately Concerning	Mildly Concerning
Very Concerning			
Nephrotoxicity – Requiring dialysis	12	-	-
<i>Clostridioides difficile</i> infection – Severe	12	-	-
Neuropathy	12	-	-
Stevens-Johnson syndrome	12	-	-
Anaphylaxis	12	-	-
DRESS Syndrome	11	1	-
Moderately Concerning			
Nephrotoxicity – Not requiring dialysis	-	11	1
<i>Clostridioides difficile</i> infection – Non-severe	-	10	2
Hepatotoxicity	1	4	4
Encephalopathy	2	9	-
Seizures	7	5	-
Hemolytic anemia	-	10	-
Neutropenia	1	9	-
Thrombocytopenia	1	8	1
Prolonged QTc	1	10	-
Mildly Concerning			
Diarrhea, nausea, or emesis	-	1	10
Non-hives rash	-	-	10
Myositis	-	-	10

*Not all participants voted for each adverse event