

EV0627

Mental health of college students: Five-year experience of the university psychiatric outpatient clinic of São João hospital centre

R. Guedes*, M. Mota-Oliveira, E. Pereira, M.J. Peixoto, I. Ferraz, M. Oliveira, C. Silveira

Centro Hospitalar de São João, Clínica de Psiquiatria e Saúde Mental, Porto, Portugal

* Corresponding author.

Introduction The clinic of psychiatry and mental health of São João hospital centre (Oporto, Portugal) has implemented a psychiatric consultation to support college students since 2007. This consultation is open to all of the universities in the metropolitan area of Oporto.

Aims/objectives This specific consultation aims to detect and intervene early in the treatment of psychiatric illness and to promote mental health in this specific population.

Methods Retrospective study conducted in the São João hospital centre, Oporto, Portugal. Patients attending university student's specific consultation between January 1st 2011 and March 31st 2016 were included. Data collection included sociodemographic variables, clinical diagnosis (ICD-10, WHO, 1992) and psychological scales (WAIS-III, BSI, HADS, SF-36 and NEO-PI-R). SPSS® software (v. 20.0, 2011) was used for statistical analysis.

Results In this study, 139 patients were included (66.2% female, medium age 23.1 years old). The majority of patients were medical, engineering or nursing students (respectively 20.9%, 18.0% and 17.9%). The most frequent diagnosis was adjustment disorders, anxiety disorders, mood disorders and personality disorders. 54.6% completed all the psychological scales.

Conclusions The number of students with mental health issues is increasing. Early detection and treatment of these pathologies may allow improvements on the educational, economic and social levels, as well as in the quality of life.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.957>

EV0628

Burnout of caregivers in geriatric Institution: "Coping" strategies

H. Ben ammar, G. Hamdi*, Z. El hechmi

Razi Hospital, F, Mannouba, Tunisia

* Corresponding author.

Introduction The health sector has long been recognized to be a very stressful work environment for teams that can lead to "burnout". Geriatric institutions are no exception to this observation; this state has deleterious effects on health care as on the quality of care.

Objectives Measure the burnout rate among caregivers in geriatric institutions and identify associated factors and coping strategies specific to this population in order to provide the necessary preventive measures.

Subjects and method A cross-sectional study, conducted among caregivers exercising at the shelter for aged subjects of Manouba, Tunisia. We used a pre-survey exploring the socio-demographic data associated with two validated scales: the Maslach Burnout Inventory assessing the level of burnout and the Brief COPE assessing coping strategies.

Results Thirty-one subjects were recruited. The prevalence of burnout was 45.16%. Respectively 32.26%, 25.80% and 45.16% of the respondents had high scores in dimensions emotional exhaustion, depersonalization and personal accomplishment at work. The analytical study revealed that seniority in work increased the risk of

burnout. The coping strategies of caregivers in geriatric institution facing burnout were mainly centered on emotion. The number of years of experience has been associated with burnout in our study.
Conclusion This work reaffirms that the population of caregivers in geriatric institution is at risk of burnout and allows to identify predictors. The establishment of individual and collective measures is essential for appropriate treatment references (In the body text, a publication should be referred to by a consecutive number between bracket)

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.958>

EV0629

The relationship between happiness, general health and life expectancy of cancer patients

S. Heydari

Islamic Azad University, Clinical Psychology, Qazvin, Iran

Introduction Cancer is a disease that all human beings are afraid of it and anyone, of any age may be affected.

Objective The objectives of this research are relationship between well-being and life expectancy in patients with cancer in Noshahr Shahid Beheshti hospital is the general purpose of this research.

Aim Studying the relationship between happiness, general health and life expectancy of cancer patients.

Method The method used in this research is descriptive and correlation. Statistical population contains all the patients who were diagnosed with cancer in Shahid Beheshti hospital Noshahr. The panel sampling used as sampling method and the sample size was limited to 50 people.

In order to collect the data, we used Oxford's happiness questionnaire containing 29 multiple-choice phrases, Life expectancy questionnaire by Schneider (1991) and general health questionnaire (GHQ-28). In descriptive review of the collected data the mean, standard deviation and frequency tables were used and in inferentially section in order to determine and compare the obtained scores in these 3 tests, according to the findings, parametric methods were used.

Result The results show that the correlation between mental health, happiness and life expectancy was significant.

Conclusion As life expectancy increases accountability; Happiness and mental health in patients with cancer increases by the same amount.

Keywords Happiness; General health; Life expectancy; Cancer
Disclosure of interest The author has not supplied his/her declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.959>

EV0630

Does the use of psychotropic medication go up during focused efforts to bring coercion and restraint down?

M. Højlund^{1,*}, L. Høgh², P. Munk-Jørgensen³, E. Stenager¹

¹ Mental Health Services, Region of Southern Denmark, Research Unit Department of Psychiatry, Aabenraa, Denmark

² Mental Health Services, Region of Southern Denmark, Department of Psychiatry, Aabenraa, Denmark

³ Mental Health Services, Region of Southern Denmark, Department of Psychiatry, Odense, Denmark

* Corresponding author.

Introduction Antipsychotic or anxiolytic medicine is widely used in agitated patients in risk of coercion. However this medication is