



Debate

D001

Con

Clinical/therapeutic: debate: sexual addiction: does it exist?

A. Weinstein

University of Ariel, Behavioral Science, Ariel, Israel

It has been argued that compulsive sexual behavior (CSB) similar to pathological gambling (PG), meets the criteria for addiction. There is evidence showing that compulsive sexual behavior has the characteristics of addiction such as salience, mood modification, tolerance, withdrawal and adverse consequences. There are studies that have shown that exposure to visual sexual stimuli in individuals with compulsive sexual behavior is associated with activation of reward mechanisms similar to drug addiction. Cross-sectional studies report high rates of co-morbidity between compulsive sexual behavior and other psychiatric disorders such as depression, anxiety; Attention Deficit Hyperactivity Disorder (ADHD), obsessive-compulsive disorder (OCD) and personality disorders. However, despite many similarities between the features of hypersexual behavior and substance-related disorders there are gaps in our knowledge on compulsive sexual behavior and its treatment which precludes a definite conclusion that this is a behavioral addiction rather than an impulse control disorder. Therefore, more research is needed before definitively characterizing HD as an addiction at this time. This talk will review the empirical evidence and it will summarize the arguments against considering sexual addiction as a behavioral addiction (the cons side).

Disclosure of interest.– The authors have not supplied a conflict of interest statement.

D002

Pro

Mental health policy: debate: do we need compulsory treatments in psychiatric practice?

T. Kallert

Psychiatric Health Care Facilities of Upper Franconia GeBO, Bezirkskrankenhaus Bayreuth, Bayreuth, Germany; Dresden University of Technology, Faculty of Medicine, Dresden, Germany; Department of Psychiatry, Psychotherapy, and Psychosomatic Medicine, Bayreuth, Germany

<https://doi.org/10.1016/j.europsy.2019.01.004>

0924-9338/© 2019 Published by Elsevier Masson SAS.

Mostly based on the results of the EUNOMIA study, still the largest prospective study on the use and outcomes of coercive measures (involuntary hospitalization, mechanical restraint, forced medication, seclusion) in general hospital psychiatry ever conducted, the presentation will outline that

1. Coercive interventions are a medico-legal and clinical reality in Europe, but show significant variation across countries; further, patients' views on involuntary hospitalization also differ across sites
2. There might be a link between the extent to which national mental health legislation protects patients' rights and the extent to which patients retrospectively evaluate that their involuntary admission was appropriate
3. Patients who feel coerced to admission may have a poorer prognosis than legally involuntary patients
4. Effective treatment of positive symptoms and improving patients' global functioning may lead to a reduction in perceived coercion
5. Caregivers' appraisals of involuntary inpatient treatment correlate with patients' symptom improvement

Conclusion.– If compulsory treatments in psychiatric practice are needed is an open question. Many aspects of the use of such interventions deserve deeper attention in research and clinical practice. The complexity of this field is such that simple pro-con answers are not possible. In general, we have to work on a standard of clinical practice guided by respecting autonomy and rights of our patients to the utmost.

Disclosure of interest.– The authors have not supplied a conflict of interest statement.

D003

Con

Mental health policy: debate: do we need compulsory treatments in psychiatric practice?

G. Szmukler

King's College London, Institute of Psychiatry- Psychology and Neuroscience, London, United Kingdom

I shall argue that involuntary treatment can be unnecessary in the practice of psychiatry. This is the position taken by a number of UN treaty bodies, including the UN Committee for the Convention on the Rights of Persons with Disabilities (CRPD), the UN Working Group on Arbitrary Detention and the UN Commissioner on

Human Rights. Other UN bodies' positions are less explicit about an absolute prohibition on involuntary interventions, but are framed in terms that support a central role for 'will and preferences', a key concept in the UN CRPD. They call for an urgent need to develop alternatives to coercive interventions. An important Resolution on Mental Health and Human Rights from the UN Human Rights Council calls upon States to "abandon all practices that fail to respect the rights, will and preferences of all persons, on an equal basis" and to "provide mental health services for persons with mental health conditions or psychosocial disabilities on the same basis as to those without disabilities, including on the basis of free and informed consent".

I shall note the huge variation, twenty- to thirty-fold, between European countries in the use of involuntary treatment, implying unacceptable arbitrariness in its use. Attention will be drawn to the negligible research effort devoted to developing treatment approaches for the avoidance of coercive interventions. I shall then show how a focus on supportive measures aimed at enhancing patients' involvement in their care, together with a focus on respecting the person's 'will and preferences' would result in involuntary treatment becoming unnecessary.

Disclosure of interest.– The authors have not supplied a conflict of interest statement.

D004

Con

Mental health policy: debate: should the UHR paradigm for transition to mental disorder be abandoned?

F. Schultze-Lutter

Heinrich-Heine University- Medical Faculty, Department of Psychiatry and Psychotherapy, Düsseldorf, Germany

Current clinical high-risk (CHR) of psychosis criteria – particularly criteria relying on attenuated or transient positive symptoms and cognitive basic symptoms – are associated with conversion rates many times higher than the general incidence of psychosis. Yet, non-conversions still outnumber conversions, and CHR-relevant phenomena are not uncommon in the community, fueling an ongoing debate about their justification. This debate, however, widely disregards main general findings: persons meeting CHR criteria already suffer from multiple mental and functional disturbances for those they seek help; they exhibit various psychological and cognitive deficits along with morphological and functional cerebral changes, whereby, the majority of them fulfils general criteria for mental disorders; and beyond their association with subsequent psychotic disorders, CHR criteria do not specifically associate with any other mental disorder. Furthermore, while CHR symptoms might not be uncommon in the general population, CHR criteria almost as rare as psychotic disorders and, already at mere symptom level, are considerably associated with proxy measures of clinical relevance on community level, including low psychosocial functioning. Hence, the clinical picture defined by current CHR criteria might not be perceived only in terms of a psychosis-risk syndrome alone but rather as a psychosis-spectrum disorder in its own right with conversion to psychosis just being one and likely the worst of several outcomes and still the best available starting-point for an early detection of psychosis. Thus, the UHR paradigm clearly should not be abandoned but might rather act as a model for the early detection of other mental disorders.

Disclosure of interest.– The authors have not supplied a conflict of interest statement.