

CORRIGENDUM

Selenium intakes in the Irish adult population – CORRIGENDUM

Maria Buffini¹, Anne P. Nugent^{1,2}, Janette Walton^{3,4}, Albert Flynn³ and Breige A. McNulty¹

¹UCD Institute of Food and Health, School of Agriculture and Food Science, University College Dublin, Belfield, Dublin 4, Republic of Ireland

²Institute for Global Food Security, Queen's University Belfast, Belfast, Northern Ireland, UK

³School of Food and Nutritional Sciences, University College Cork, Cork, Republic of Ireland

⁴Department of Biological Sciences, Munster Technological University, Cork, Republic of Ireland

Journal of Nutritional Science (2023), vol. 12, e46, page 1 of 1

doi:10.1017/jns.2023.35

In Table 2 the arrow before the letters ‘TUL’ is pointing downwards. The arrow should be pointing upwards.

The correct table is shown below.

Table 2. Percentage of population groups with mean daily Se intakes ($\mu\text{g}/\text{day}$ from all sources) below the adequate intake and lower reference nutrient intake and above the tolerable upper limit.

Total Population* (n = 1051)	Men (n = 523)	Women (n = 528)	Men				Women				
			18–35y (n = 207)	36–50y (n = 143)	51–64y (n = 98)	$\geq 65\text{y}$ (n = 75)	18–35y (n = 170)	36–50y (n = 165)	51–64y (n = 106)	$\geq 65\text{y}$ (n = 87)	
% ↓ AI	46.81	28.68	64.77	20.77	34.27	30.61	37.33	57.06	67.27	66.98	72.41
% ↓ LRNI	4.09	0.96	7.2	0	2.1	0	2.67	7.65	5.45	6.6	10.34
% ↑ TUL	0	0	0	0	0	0	0	0	0	0	0

AI (adequate intake), LRNI (lower reference nutrient intake) and TUL (tolerable upper limit) are 70 $\mu\text{g}/\text{day}$, 40 $\mu\text{g}/\text{day}$ and 300 $\mu\text{g}/\text{day}$ respectively^(31,32,33). *Under reporters excluded.