



Generic transfer from $\mathrm{GSp}(4)$ to $\mathrm{GL}(4)$

Mahdi Asgari and Freydoon Shahidi

ABSTRACT

We establish the Langlands functoriality conjecture for the transfer from the generic spectrum of $\mathrm{GSp}(4)$ to $\mathrm{GL}(4)$ and give a criterion for the cuspidality of its image. We apply this to prove results toward the generalized Ramanujan conjecture for generic representations of $\mathrm{GSp}(4)$.

1. Introduction

Let k be a number field and let G denote the group $\mathrm{GSp}(4, \mathbb{A}_k)$. The (connected component of the) L -group of G is $\mathrm{GSp}(4, \mathbb{C})$, which has a natural embedding into $\mathrm{GL}(4, \mathbb{C})$. Langlands functoriality predicts that associated to this embedding there should be a transfer of automorphic representations of G to those of $\mathrm{GL}(4, \mathbb{A}_k)$ (see [Art04]). Langlands' theory of Eisenstein series reduces the proof of this to unitary cuspidal automorphic representations. We establish functoriality for the generic spectrum of $\mathrm{GSp}(4, \mathbb{A}_k)$. More precisely (cf. Theorem 2.4), we prove the following.

Let π be a unitary cuspidal representation of $\mathrm{GSp}(4, \mathbb{A}_k)$, which we assume to be globally generic. Then π has a unique transfer to an automorphic representation Π of $\mathrm{GL}(4, \mathbb{A}_k)$. The transfer is generic (globally and locally) and satisfies $\omega_\Pi = \omega_\pi^2$ and $\Pi \simeq \tilde{\Pi} \otimes \omega_\pi$. Here, ω_π and ω_Π denote the central characters of π and Π , respectively.

Moreover, we give a *cuspidality criterion* for Π and prove that, when Π is not cuspidal, it is an isobaric sum of two unitary cuspidal representations of $\mathrm{GL}(2, \mathbb{A}_k)$ (cf. Proposition 2.2).

We give a number of applications of this result. The first is Theorem 3.3 which gives estimates toward the generalized Ramanujan conjecture for generic representations of the group $\mathrm{GSp}(4)$ (cf. § 3.1). We also prove in Theorem 3.7 that any generic unitary cuspidal representation of $\mathrm{GSp}(4, \mathbb{A}_k)$ is *weakly Ramanujan* (cf. § 3.2 for the definition). In § 3.3 we use our main result to give an immediate proof that the spin L -function of a generic unitary cuspidal representation of $\mathrm{GSp}(4, \mathbb{A})$ is entire. This is due to the fact that the spin L -function of $\mathrm{GSp}(4)$ now becomes a standard L -function (or product of two such L -functions) of general linear groups. This fact has also been proved recently for a certain class of globally generic representations of $\mathrm{GSp}(4, \mathbb{A}_k)$ by Takloo-Bighash using different methods from ours.

We should note here that the transfer from $\mathrm{GSp}(4)$ to $\mathrm{GL}(4)$ has been expected by experts in the field for a long time. It is our understanding that Jacquet, Piatetski-Shapiro and Shalika knew how to prove this result, at least in principle, but, as far as we know, their result was never published. We should also point out that their proof is based on methods that are fairly disjoint from ours.

Our method of proof is to start with our earlier, more general but weaker, result on generic transfer from GSpin groups to GL (cf. [AS06]). This gives us the existence of Π . We then use the

Received 5 January 2005, accepted in final form 15 August 2005.

2000 *Mathematics Subject Classification* 11F70, 11R42, 22E55.

Keywords: Langlands functoriality, automorphic representation, L -function.

First author partially supported by the NSF grant DMS-0111298 at IAS; second author partially supported by the NSF grant DMS-0200325.

This journal is © Foundation Compositio Mathematica 2006.

results of Piatetski-Shapiro and Soudry on analytic properties of L -functions of $\mathrm{GSp}(4)$ twisted by $\mathrm{GL}(1)$ and $\mathrm{GL}(2)$ to get more information about the representation Π . It is exactly the lack of such results in the general case of GSpin groups that prevents us from carrying out our analysis for the more general case for now. However, as we pointed out in [AS06], there are currently two ways to overcome this problem. One is to prove an analogue of descent theory for these groups as was done for classical groups by Ginzburg, Rallis and Soudry [GRS01, Sou05]. The other is to use techniques along the lines of [Kim01, Kim02].

The case of the transfer for all automorphic representations of $\mathrm{GSp}(4)$ (whether generic or not) requires Arthur’s trace formula. The process and the expected issues in this case are outlined in [Art04].

2. Main result

Let k be a number field and let $\mathbb{A} = \mathbb{A}_k$ denote its ring of adeles. We define the similitude symplectic group of degree four via

$$\mathrm{GSp}(4) = \{g \in \mathrm{GL}(4) : {}^t g J g = \mu(g) J\},$$

where

$$J = \begin{pmatrix} & & & 1 \\ & & 1 & \\ & -1 & & \\ -1 & & & \end{pmatrix}$$

and $\mu(g) \in \mathrm{GL}(1)$ is the similitude character. We fix the following parametrization of the elements of the maximal torus \mathbf{T} in $\mathrm{GSp}(4)$:

$$\mathbf{T} = \left\{ t = t(a_0, a_1, a_2) = \begin{pmatrix} a_0 a_1 a_2 & & & \\ & a_0 a_1 & & \\ & & a_0 a_2 & \\ & & & a_0 \end{pmatrix} \right\}.$$

The above agrees with our previous more general notation for the group $\mathrm{GSpin}(2n + 1)$ in [AS06]. Recall that the group $\mathrm{GSp}(4)$ is identified with $\mathrm{GSpin}(5)$.

Let $\pi = \otimes_v \pi_v$ be a globally ψ -generic unitary cuspidal automorphic representation of $\mathrm{GSp}(4, \mathbb{A})$. Here, $\psi = \otimes_v \psi_v$ is a non-trivial additive character of $k \backslash \mathbb{A}$ defining a character of the unipotent radical of the standard upper-triangular Borel subgroup in the usual way. We fix ψ now and do not repeat it in the rest of this paper. Let S be any non-empty finite set of non-archimedean places v , which includes those v with π_v or ψ_v ramified. We proved in [AS06] that there exists an automorphic representation $\Pi = \otimes_v \Pi_v$ of $\mathrm{GL}(4, \mathbb{A})$ such that Π_v is a local transfer of π_v for v outside of S .

To be more explicit, assume that $v \notin S$. If v is archimedean, then π_v is given by a parameter $\phi_v : W_v \rightarrow \mathrm{GSp}(4, \mathbb{C})$, where W_v is the Weil group of k_v (cf. [Lan89]). Let $\Phi_v : W_v \rightarrow \mathrm{GL}(4, \mathbb{A})$ be given by $\Phi_v = \iota \circ \phi_v$, where $\iota : \mathrm{GSp}(4, \mathbb{C}) \rightarrow \mathrm{GL}(4, \mathbb{C})$ is the natural embedding. Then Φ_v is the parameter of Π_v .

If $v \notin S$ is non-archimedean, then π_v is the unique unramified subquotient of the representation induced from an unramified character χ of $\mathbf{T}(k_v)$ to $\mathrm{GSp}(4, k_v)$. Writing

$$\chi(t(a_0, a_1, a_2)) = \chi_0(a_0)\chi_1(a_1)\chi_2(a_2), \tag{1}$$

where χ_i are unramified characters of k_v^\times and $a_i \in k_v^\times$, the representation Π_v is then the unique irreducible unramified subquotient of the representation of $\mathrm{GL}(4, k_v)$ parabolically induced from the

character

$$\chi_1 \otimes \chi_2 \otimes \chi_2^{-1} \chi_0 \otimes \chi_1^{-1} \chi_0 \tag{2}$$

of $\mathbf{T}(k_v)$.

Moreover, we proved that $\omega_\Pi = \omega^2$, where $\omega = \omega_\pi$ and ω_Π denote the central characters of π and Π , respectively, and for $v \notin S$ we have $\Pi_v \simeq \tilde{\Pi}_v \otimes \omega_{\pi_v}$, i.e. Π is nearly equivalent to $\tilde{\Pi} \otimes \omega$.

The representation Π is equivalent to a subquotient of some representation

$$\mathrm{Ind}(|\det|^{r_1} \sigma_1 \otimes \cdots \otimes |\det|^{r_t} \sigma_t), \tag{3}$$

where induction is from $\mathrm{GL}(n_1) \times \cdots \times \mathrm{GL}(n_t)$ with $n_1 + \cdots + n_t = 4$ to $\mathrm{GL}(4)$ and σ_i are the unitary cuspidal automorphic representation of $\mathrm{GL}(n_i, \mathbb{A})$ and $r_i \in \mathbb{R}$.

Without loss of generality we may assume that $r_1 \geq r_2 \geq \cdots \geq r_t$. Moreover, as Π is unitary we have $n_1 r_1 + \cdots + n_t r_t = 0$, which implies that $r_t \leq 0$. Let $T = S \cup \{v : v | \infty\}$ and consider

$$L^T(s, \pi \times \tilde{\sigma}_t) = L^T(s, \Pi \times \tilde{\sigma}_t) = \prod_{i=1}^t L^T(s + r_i, \sigma_i \times \tilde{\sigma}_t). \tag{4}$$

Here, L^T denotes the product over $v \notin T$ of the local L -factors.

If $n_t = 1$, then the left-hand side is entire by a result of Piatetski-Shapiro (cf. [Pia97, p. 274]). Now consider the right-hand side at $s_0 = 1 - r_t \geq 1$. The last term in the product has a pole at s_0 , whereas all of the others are non-zero there as $\Re(s_0 + r_i) = 1 + r_i - r_t \geq 1$. This is a contradiction.

Now assume that $n_t = 3$, i.e. $t = 2$ with $n_1 = 1$ and $n_2 = 3$. Replacing π and Π by their contragredients will change r_i to $-r_i$ and takes us back to the above situation, which gives a contradiction again.

Therefore, $n_t = 2$. In this case, the left-hand side of (4) may have a pole at $s = 1$ (cf. [PS84, Theorem 1.3] and the beginning of its proof), and if so, arguing as above, we conclude that $r_t = 0$. This means that we either have $t = 2$ with $n_1 = n_2 = 2$ or $t = 3$ with $n_1 = n_2 = 1$ and $n_3 = 2$. However, we can rule out the latter as follows.

Assume that $t = 3$ with $n_1 = n_2 = 1$ and $n_3 = 2$. Then, it follows from the fact that $r_3 = 0$ and the conditions $r_1 \geq r_2 \geq r_3$ and $r_1 + r_2 + 2r_3 = 0$ that all of the r_i would be zero in this case. This implies that if we consider the L -function of π twisted by $\tilde{\sigma}_1$, we have

$$L^T(s, \pi \times \tilde{\sigma}_1) = L^T(s, \sigma_1 \times \tilde{\sigma}_1) L^T(s, \sigma_2 \times \tilde{\sigma}_1) L^T(s, \sigma_3 \times \tilde{\sigma}_1). \tag{5}$$

Now the left-hand side is again entire by Piatetski-Shapiro's result mentioned above and the right-hand side has a pole at $s = 1$, which is a contradiction.

Therefore, the only possibilities are $t = 1$ (i.e. Π unitary cuspidal) or $t = 2$ and $n_1 = n_2 = 2$ with $r_2 = 0$. In the latter case, we also get $r_1 = 0$, as $r_1 + r_2 = 0$ by unitarity of the central character. Moreover, in this case we have $\sigma_1 \not\cong \sigma_2$ as, otherwise,

$$L^T(s, \pi \times \tilde{\sigma}_1) = L^T(s, \sigma_1 \times \tilde{\sigma}_1) L^T(s, \sigma_2 \times \tilde{\sigma}_1) \tag{6}$$

must have a double pole at $s = 1$ while any possible pole of the left-hand side at $s = 1$ is simple (cf. the proof of [PS84, Theorem 1.3]).

Therefore, we have proved the following.

PROPOSITION 2.1. *Let π be a globally generic unitary cuspidal automorphic representation of $\mathrm{GSp}(4, \mathbb{A})$ and let Π be any transfer of π to $\mathrm{GL}(4, \mathbb{A})$. Then Π is a subquotient of an automorphic representation as in (3) with either $t = 1$, $n_1 = 4$ and $r_1 = 0$ (i.e. Π is unitary cuspidal) or $t = 2$, $n_1 = n_2 = 2$ and $r_1 = r_2 = 0$. In the latter case, we have $\sigma_1 \not\cong \sigma_2$.*

In fact, we can get more precise information.

PROPOSITION 2.2. *Let π be a globally generic unitary cuspidal automorphic representation of $\mathrm{GSp}(4, \mathbb{A})$ with $\omega = \omega_\pi$ its central character and let Π be any transfer as above. Then, $\Pi \simeq \tilde{\Pi} \otimes \omega$ (not just nearly equivalent). Moreover:*

- (a) *the representation Π is cuspidal if and only if π is not obtained as a Weil lifting from $\mathrm{GSO}(4, \mathbb{A})$;*
- (b) *if Π is not cuspidal, then it is the isobaric sum of two representations $\Pi = \Pi_1 \boxplus \Pi_2$, where each Π_i is a unitary cuspidal automorphic representation of $\mathrm{GL}(2, \mathbb{A})$ satisfying $\Pi_i \simeq \tilde{\Pi}_i \otimes \omega$ and $\Pi_1 \not\simeq \Pi_2$.*

Proof. By Proposition 2.1, Π is not cuspidal if and only if it is a subquotient of

$$\Sigma = \mathrm{Ind}_{\mathrm{GL}(2, \mathbb{A}) \times \mathrm{GL}(2, \mathbb{A})}^{\mathrm{GL}(4, \mathbb{A})}(\sigma_1 \otimes \sigma_2), \tag{7}$$

where σ_i are unitary cuspidal automorphic representation of $\mathrm{GL}(2, \mathbb{A})$.

On the other hand, by [PS84, Theorem 1.3] mentioned above, the representation π is obtained as a Weil lifting from $\mathrm{GSO}(4, \mathbb{A})$ if and only if there exists an automorphic representation τ of $\mathrm{GL}(2, \mathbb{A})$ such that $L^T(s, \pi \times \tau)$ has a pole and in that case τ can be normalized so that the pole occurs at $s = 1$.

Now assume that Π is cuspidal. Then for any τ as above we have

$$L^T(s, \pi \times \tau) = L^T(s, \Pi \times \tau),$$

which is entire. Therefore, π is not obtained as a Weil lifting from $\mathrm{GSO}(4, \mathbb{A})$. Moreover, as Π is cuspidal, so is $\tilde{\Pi} \otimes \omega$ and they are nearly equivalent, therefore, by the strong multiplicity one theorem [JS81a, JS81b, Pia79] we have $\Pi \simeq \tilde{\Pi} \otimes \omega$.

Next, assume that Π is not cuspidal and, hence, is given as a subquotient of Σ above. We claim that each $\Sigma_v = \mathrm{Ind}(\sigma_{1,v} \otimes \sigma_{2,v})$ is irreducible. To see this note that each $\sigma_{i,v}$ is generic unitary and is either a tempered representation of $\mathrm{GL}(2, k_v)$ or a complementary series $I(\chi |^\alpha, \chi |^{-\alpha})$ with $\alpha \in (0, \frac{1}{2})$ and χ a unitary character. If both of the $\sigma_{i,v}$ are tempered, then irreducibility of Σ_v is clear. If both are complementary series of the form $I(\chi_1 |^\alpha, \chi_1 |^{-\alpha})$ and $I(\chi_2 |^\beta, \chi_2 |^{-\beta})$ with $\alpha, \beta \in (0, \frac{1}{2})$ and χ_i unitary characters, then for Σ_v to be reducible we should have $\alpha \pm \beta = \pm 1$, which is not possible. Finally, if one of $\sigma_{i,v}$ is tempered and the other is complementary series, then we have $\Sigma_v = \mathrm{Ind}(\chi |^{-\alpha}, \chi_1, \chi_2, \chi |^\alpha)$, $\Sigma_v = \mathrm{Ind}(\chi |^{-\alpha}, Q(\chi_1 |^{-1/2}, \chi_1 |^{1/2}), \chi |^\alpha)$ or $\Sigma_v = \mathrm{Ind}(\chi |^{-\alpha}, \eta, \chi |^\alpha)$. Here, χ_1, χ_2 and χ are unitary characters, $\alpha \in (0, \frac{1}{2})$, η is a unitary supercuspidal representation of $\mathrm{GL}(2, k_v)$, and $Q(\chi_1 |^{-1/2}, \chi_1 |^{1/2})$ denotes the Steinberg representation twisted by the unitary character χ_1 . Again, in all these cases the representation Σ_v is irreducible as $\alpha \in (0, \frac{1}{2})$.

Therefore, at every place v the representation Π_v is the same as the irreducible Σ_v and, hence, it is the Langlands quotient. This means that Π is an isobaric representation, i.e. the isobaric sum of σ_1 and σ_2 . Again by the strong multiplicity one theorem, which remains valid for isobaric representations [JPS83], we conclude that $\Pi \simeq \tilde{\Pi} \otimes \omega$. Now, just take Π_i to be σ_i .

Finally, by [AS06, Proposition 7.4], which was based on classification theorems of Jacquet and Shalika, we know that we either have $\Pi_i \simeq \tilde{\Pi}_i \otimes \omega$ for $i = 1, 2$ or we have $\Pi_1 \simeq \tilde{\Pi}_2 \otimes \omega$ (or equivalently, $\Pi_2 \simeq \tilde{\Pi}_1 \otimes \omega$). However, the latter case will not occur when π is cuspidal and generic as, otherwise, π will be nearly equivalent to an Eisenstein series representation, i.e. π will be a CAP representation of $\mathrm{GSp}(4, \mathbb{A})$. This is impossible by [PS87, Theorem 1.1]. This completes the proof. \square

Remark 2.3. Note that any other transfer Π' of π is also a subquotient of Σ in (7), which is irreducible. Therefore, π has a *unique* transfer to $\mathrm{GL}(4, \mathbb{A})$, which we continue to denote by Π . In particular, this implies that we have not lost any information at the places where we did not have a natural candidate for the local transfer.

Moreover, as Π is either a unitary cuspidal representation of a general linear group or an isobaric sum of two such, every local representation Π_v is full induced and generic.

Furthermore, if Π is not cuspidal, then $\Pi_1 \not\cong \Pi_2$ implies that $L(s, \Pi_1 \times \tilde{\Pi}_2)$ has no pole at $s = 1$. This implies that the Fourier coefficient of Π along the unipotent radical of our fixed Borel subgroup is non-vanishing, i.e. Π is globally generic [Sha81].

We collect the above results in the following theorem, which is our main result.

THEOREM 2.4. *Let π be a globally generic unitary cuspidal automorphic representation of $\mathrm{GSp}(4, \mathbb{A})$ with central character ω . Then π has a unique transfer Π to $\mathrm{GL}(4, \mathbb{A})$, which satisfies $\Pi \simeq \tilde{\Pi} \otimes \omega$ and its central character is ω^2 . Moreover, Π is either unitary cuspidal or an isobaric sum $\Pi_1 \boxplus \Pi_2$ of two inequivalent unitary cuspidal automorphic representations of $\mathrm{GL}(2, \mathbb{A})$ satisfying $\Pi_i \simeq \tilde{\Pi}_i \otimes \omega$. The latter is the case if and only if π is obtained as a Weil lifting from $\mathrm{GSO}(4, \mathbb{A})$. Furthermore, Π is globally generic, i.e. it has a non-vanishing Fourier coefficient along the unipotent radical of our fixed Borel subgroup.*

In fact, we can get more precise information about the local representations at places $v \in S$.

PROPOSITION 2.5. *Fix $v \in S$ and let*

$$\pi_v \simeq \mathrm{Ind}(\pi_{1,v} |\det|^{b_{1,v}} \otimes \cdots \otimes \pi_{t,v} |\det|^{b_{t,v}} \otimes \pi_{0,v}) \tag{8}$$

be an irreducible generic representation of $\mathrm{GSp}(4, k_v)$, where each $\pi_{i,v}$ is a tempered representation of some $\mathrm{GL}(n_i, k_v)$, $b_{1,v} > \cdots > b_{t,v}$, and $\pi_{0,v}$ is a tempered generic representation of some $\mathrm{GSp}(2m, k_v)$. (Note that there are very few possibilities for n_i and m as $n_1 + \cdots + n_t + m = 2$. We are allowing $m = 0$ and, by convention, $\mathrm{GSp}(0) = \mathrm{GL}(1)$.) Let ω_v denote the central character of π_v . Assume that π_v is the local component of the globally generic unitary cuspidal representation π of $\mathrm{GSp}(4, \mathbb{A})$ and let Π be its transfer to $\mathrm{GL}(4, \mathbb{A})$. Then the local component Π_v of Π at v is generic and of the form

$$\Pi_v \simeq \mathrm{Ind}(\pi_{1,v} |\det|^{b_{1,v}} \otimes \cdots \otimes \pi_{t,v} |\det|^{b_{t,v}} \otimes \Pi_{0,v} \otimes (\tilde{\pi}_{t,v} \otimes \omega_v) |\det|^{-b_{t,v}} \otimes \cdots \otimes (\tilde{\pi}_{1,v} \otimes \omega_v) |\det|^{-b_{1,v}}), \tag{9}$$

where $\Pi_{0,v}$ is a tempered generic representation of $\mathrm{GL}(2m, k_v)$ if $m > 0$.

Proof. Let us remark that, as in [CKPS04, § 7], one could define the notion of local transfer and obtain complete information about such transfers for a general irreducible admissible generic representation, whether a local component of a global representation or not. (In fact, the representation $\Pi_{0,v}$ would then be the local transfer of $\pi_{0,v}$.) However, we do not need the full extent of such results in this paper.

Recall that we already proved (cf. Remark 2.3) that each Π_v is generic and is full induced. Let $v \in S$ and consider π_v and Π_v as in the proposition. We first show that if ρ_v is any supercuspidal representation of $\mathrm{GL}(r, k_v)$, then

$$\gamma(s, \pi_v \times \rho_v, \psi_v) = \gamma(s, \Pi_v \times \rho_v, \psi_v). \tag{10}$$

The key here is the fact that there exists a unitary cuspidal representation ρ of $\mathrm{GL}(r, \mathbb{A})$ such that its local component at v is ρ_v and at all other finite places $w \neq v$ the local component ρ_w is unramified (cf. [Sha90, Proposition 5.1]). Now applying the converse theorem of Cogdell and Piatetski-Shapiro with $S' = S - \{v\}$ will give the result exactly as in the proof of [CKPS04, Proposition 7.2]. Moreover, by multiplicativity of γ -factors, we conclude that (10) also holds if ρ_v is a discrete series representation of $\mathrm{GL}(r, k_v)$.

Next, we claim that if π_v is tempered, then so is Π_v . Here, again the main tool is multiplicativity of γ -factors and the proof is exactly as in [CKPS04, Lemma 7.1]. This proves the proposition for the case $m = 2$. If $m = 1$, then the group $\mathrm{GSp}(2m) = \mathrm{GSp}(2)$ is the same as $\mathrm{GL}(2)$ and we set $\Pi_{0,v}$ to be $\pi_{0,v}$ itself. For $m = 0$ we need no choice of $\Pi_{0,v}$. Now, let $T = \{w_0\}$ consist of a single finite place $w_0 \neq v$ at which π_v is unramified and consider the representation Π' of $\mathrm{GL}(4, \mathbb{A})$ whose local

components are the same as Π outside of S and are the irreducible induced representations on the right-hand side of (9) when $v \in S$. We can now apply the converse theorem again to Π' and $T = \{w_0\}$ to conclude that Π' is a transfer of π . The key here is that the induced representations on the right-hand side of (9) have the right L -functions. Therefore, by uniqueness of the transfer we proved earlier, we have $\Pi'_v \simeq \Pi_v$ for $v \in S$. This completes the proof. \square

3. Applications

We first recall that the current formulation of the Ramanujan conjecture for generic cuspidal representations states that for any quasi-split group and any globally generic unitary cuspidal automorphic representation $\pi = \otimes_v \pi_v$, the local components π_v are tempered for all places v . As an application of our main theorem, we can prove two types of results in this direction: estimates toward this conjecture for the group $\mathrm{GSp}(4)$ as well as a weaker version of it for generic representations of this group.

3.1 Estimates toward Ramanujan

Following [CKPS04], we introduce the following notation in order to prove estimates. Let $\Pi = \otimes_v \Pi_v$ be a unitary cuspidal automorphic representation of $\mathrm{GL}(m, \mathbb{A}_k)$. For every place v the representation Π_v is unitary generic and can be written as a full induced representation

$$\Pi_v \simeq \mathrm{Ind}(\Pi_{1,v}|\det|^{a_{1,v}} \otimes \cdots \otimes \Pi_{t,v}|\det|^{a_{t,v}}) \tag{11}$$

with $a_{1,v} > \cdots > a_{t,v}$ and each $\Pi_{i,v}$ tempered [Vog78, Zel80].

DEFINITION 3.1. We say Π satisfies $H(\theta_m)$ with $\theta_m \geq 0$ if for all places v we have $-\theta_m \leq a_{i,v} \leq \theta_m$.

The classification of generic unitary dual of $\mathrm{GL}(m)$ (see [Tad86, Vog78]) trivially gives $H(\frac{1}{2})$. The best result currently known for a general number field k says that any unitary cuspidal representation of $\mathrm{GL}(m, \mathbb{A})$ satisfies $H(\frac{1}{2} - 1/(m^2 + 1))$ (see [LRS99]). When $k = \mathbb{Q}$ and $m \leq 4$, it is $H(\frac{1}{2} - 1/(1 + m(m + 1)/2))$. The same bound is also available for $m > 4$ for $k = \mathbb{Q}$ provided that one knows that the symmetric square L -function of Π is absolutely convergent for $\Re(s) > 1$ (cf. [KS03]), but this is only currently available for $m \leq 4$. When $m = 2$ we have the better bounds of $H(\frac{1}{9})$ for a general number field k (see [KS02]) and $H(\frac{7}{64})$ for $k = \mathbb{Q}$ (see [KS03]). The Ramanujan conjecture demands $H(0)$.

Similarly, if $\pi = \otimes_v \pi_v$ is a unitary generic cuspidal automorphic representation of $\mathrm{GSp}(2n, \mathbb{A}_k)$, then by [Mui01] and [Vog78], each π_v can be written as a full induced representation

$$\pi_v \simeq \mathrm{Ind}(\pi_{1,v}|\det|^{b_{1,v}} \otimes \cdots \otimes \pi_{t,v}|\det|^{b_{t,v}} \otimes \tau_v), \tag{12}$$

where each $\pi_{i,v}$ is a tempered representation of some $\mathrm{GL}(n_i, k_v)$ and τ_v is a tempered generic representation of some $\mathrm{GSp}(2m, k_v)$ with $n_1 + \cdots + n_t + m = n$.

DEFINITION 3.2. We say π satisfies $H(\theta_n)$ with $\theta_n \geq 0$ if for all places v we have $-\theta_n \leq b_{i,v} \leq \theta_n$.

The classification of generic unitary dual of $\mathrm{GSp}(4)$ (cf. [LMT04], for example) trivially gives the estimate $H(1)$. The Ramanujan conjecture demands $H(0)$ again. For a survey of results in this direction and their applications we refer to [Sar03, Sha04].

THEOREM 3.3. *Let k be a number field and assume that all unitary cuspidal representations of $\mathrm{GL}(4, \mathbb{A}_k)$ (respectively, $\mathrm{GL}(2, \mathbb{A}_k)$) satisfy $H(\theta_4)$ (respectively, $H(\theta_2)$) and $\theta_2 \leq \theta_4$. Then any globally generic unitary cuspidal representation π of $\mathrm{GSp}(4, \mathbb{A}_k)$ satisfies $H(\theta_4)$, and even the better bound of $H(\theta_2)$ if π transfers to a non-cuspidal representation of $\mathrm{GL}(4, \mathbb{A}_k)$ (cf. Theorem 2.4).*

Proof. Let Π be the functorial transfer of π to $\mathrm{GL}(4, \mathbb{A}_k)$. If v is an archimedean place of k , then the theorem is clear as in this case local functoriality is well understood through Langlands’ parametrization (cf. the proof of [AS06, Theorem 6.1], for example, for more details).

Let v be a non-archimedean place of k at which π_v is unramified. Then it follows from (2) that π_v is given by its Frobenius–Hecke (Satake) parameter, which is of the form

$$\mathrm{diag}(\chi_1(\varpi), \chi_2(\varpi), \chi_2^{-1}(\varpi)\chi_0(\varpi), \chi_1^{-1}(\varpi)\chi_0(\varpi)), \tag{13}$$

where ϖ denotes a uniformizer of k_v . If Π is cuspidal, then for $i = 1, 2$ we have by assumption

$$q_v^{-\theta_4} \leq |\chi_i(\varpi)| \leq q_v^{\theta_4}.$$

If Π is not cuspidal, then we have similar inequalities with θ_4 replaced by the even better estimate of θ_2 . As π_v is unitary, we have $|\chi_0(\varpi)| = 1$. Therefore, Frobenius–Hecke parameters of π_v also satisfy similar inequalities.

Next, assume that v is a place of k in S . Then, by Proposition 2.5, a similar argument as above works again. □

COROLLARY 3.4. *Let π be a globally generic unitary cuspidal representation of $\mathrm{GSp}(4, \mathbb{A}_k)$. Then π satisfies $H(\frac{15}{34})$. If π transfers to a non-cuspidal representation of $\mathrm{GL}(4, \mathbb{A}_k)$, then it satisfies $H(\frac{1}{9})$. If $k = \mathbb{Q}$, then we have the better estimates of $H(\frac{9}{22})$ and $H(\frac{7}{64})$, respectively.*

Proof. The proof is immediate if we combine Theorem 3.3 with the known results on estimates for the general linear groups mentioned above. □

COROLLARY 3.5. *The Ramanujan conjecture for unitary cuspidal representations of $\mathrm{GL}(4)$ and $\mathrm{GL}(2)$ imply the Ramanujan conjecture for the generic spectrum of $\mathrm{GSp}(4)$.*

3.2 The weak Ramanujan conjecture

Following [CP95, Ram97, Kim03] we recall the following definition.

DEFINITION 3.6. Let \mathbf{G} be a split reductive group over the number field k . Let $\pi = \bigotimes \pi_v$ be an automorphic representation of $\mathbf{G}(\mathbb{A}_k)$. We say that π is *weakly Ramanujan* if, given $\epsilon > 0$, there exists a set T of places of k containing the archimedean places and the non-archimedean ones with π_v ramified such that T has density zero and for $v \notin T$ the Frobenius–Hecke parameter $\mathrm{diag}(\lambda_{v,i})$ of π_v satisfies

$$\max_i \{|\lambda_{v,i}|, |\lambda_{v,i}^{-1}|\} \leq q_v^\epsilon.$$

Here, q_v denotes the cardinality of the residue field.

We are concerned with the cases of $\mathbf{G} = \mathrm{GL}(m)$ or $\mathbf{G} = \mathrm{GSp}(4)$ in this paper. We recall that (unitary) cuspidal representations of $\mathrm{GL}(m)$ for $m \leq 4$ are weakly Ramanujan (cf. [Ram97] and [Kim03, Propositions 3.7 and 6.3]).

Let π be a globally generic unitary cuspidal representation of $\mathrm{GSp}(4, \mathbb{A}_k)$. For any $v \notin T$ as above, let

$$\mathrm{diag}(a_{0,v}, a_{1,v}, a_{2,v}) \tag{14}$$

be the Frobenius–Hecke parameter of π_v (cf. (1)). Then, as in (13), the parameter of the local transfer Π_v is given by

$$\mathrm{diag}(a_{1,v}, a_{2,v}, a_{2,v}^{-1}a_{0,v}, a_{1,v}^{-1}a_{0,v}). \tag{15}$$

Moreover, $|a_{0,v}| = 1$ as π_v is unitary. Therefore, the above results about weak Ramanujan property of unitary cuspidal representations of $\mathrm{GL}(m)$ immediately imply the following.

THEOREM 3.7. *Let π be a globally generic unitary cuspidal representation of $\mathrm{GSp}(4, \mathbb{A}_k)$. Then π is weakly Ramanujan.*

3.3 Spinor L -function for $\mathrm{GSp}(4)$

As another application we get the following immediate corollary of our main result, Theorem 2.4.

PROPOSITION 3.8. *Let π be a globally generic unitary cuspidal representation of $\mathrm{GSp}(4, \mathbb{A}_k)$. Then the spinor L -function $L(s, \pi, \text{spin})$ is entire.*

Proof. Let Π be the transfer of π to $\mathrm{GL}(4, \mathbb{A})$. If Π is unitary cuspidal, then $L(s, \pi, \text{spin}) = L(s, \Pi)$ and if $\Pi = \Pi_1 \boxplus \Pi_2$ is the isobaric sum of two unitary cuspidal representations of $\mathrm{GL}(2, \mathbb{A})$, then we have $L(s, \pi, \text{spin}) = L(s, \Pi_1)L(s, \Pi_2)$. In either case the L -functions on the right-hand side are standard L -functions of unitary cuspidal representations of the general linear group and are entire. \square

Remark 3.9. This result has also been proved for a certain class of globally generic representations π by Takloo-Bighash in [Tak]. His methods are different from ours and are based on using an integral representation. T. Moriyama has proved certain cases of this result as well [Mor04, IM05].

ACKNOWLEDGEMENTS

We would like to thank James Cogdell for many helpful discussions and Peter Sarnak for his interest in this work. The first author would also like to thank Steve Gelbart, Dihua Jiang, Robert Langlands, Brooks Roberts and Ramin Takloo-Bighash for many helpful discussions during the course of this work.

REFERENCES

- Art04 J. Arthur, *Automorphic representations of $\mathrm{GSp}(4)$* , in *Contributions to automorphic forms, geometry, and number theory (Shalika volume)*, eds H. Hida, D. Ramakrishnan and F. Shahidi (Johns Hopkins University Press, Baltimore, MD, 2004), 65–81.
- AS06 M. Asgari and F. Shahidi, *Generic transfer for general spin groups*, *Duke Math. J.* **132** (2006), 137–190.
- CKPS04 J. Cogdell, H. Kim, I. Piatetski-Shapiro and F. Shahidi, *Functoriality for the classical groups*, *Publ. Math. Inst. Hautes Études Sci.* **99** (2004), 163–233.
- CP95 J. Cogdell and I. Piatetski-Shapiro, *Unitarity and functoriality*, *Geom. Funct. Anal.* **5** (1995), 164–173.
- GRS01 D. Ginzburg, S. Rallis and D. Soudry, *Generic automorphic forms on $\mathrm{SO}(2n+1)$: functorial lift to $\mathrm{GL}(2n)$, endoscopy, and base change*, *Internat. Math. Res. Notices* **14** (2001), 729–764.
- IM05 T. Ishi and T. Moriyama, *Spinor L -functions for generic cusp forms on $\mathrm{GSp}(2)$ belonging to principal series representations*, Preprint (2005).
- JPS83 H. Jacquet, I. Piatetskii-Shapiro and J. Shalika, *Rankin–Selberg convolutions*, *Amer. J. Math.* **105** (1983), 367–464.
- JS81a H. Jacquet and J. Shalika, *On Euler products and the classification of automorphic representations. I*, *Amer. J. Math.* **103** (1981), 499–558.
- JS81b H. Jacquet and J. Shalika, *On Euler products and the classification of automorphic forms. II*, *Amer. J. Math.* **103** (1981), 777–815.
- Kim01 H. Kim, *Residual spectrum of odd orthogonal groups*, *Internat. Math. Res. Notices* **17** (2001), 873–906.

- Kim02 H. Kim, *Applications of Langlands' functorial lift of odd orthogonal groups*, Trans. Amer. Math. Soc. **354** (2002), 2775–2796.
- Kim03 H. Kim, *Functoriality for the exterior square of GL_4 and the symmetric fourth of GL_2* (with Appendix 1 by D. Ramakrishnan and Appendix 2 by H. Kim and P. Sarnak), J. Amer. Math. Soc. **16** (2003), 139–183.
- KS02 H. Kim and F. Shahidi, *Cuspidality of symmetric powers with applications*, Duke Math. J. **112** (2002), 177–197.
- KS03 H. Kim and P. Sarnak, *Refined estimates towards the Ramanujan and Selberg conjectures*, J. Amer. Math. Soc. **16** (2003). Appendix 2 to [Kim03].
- Lan89 R. Langlands, *On the classification of irreducible representations of real algebraic groups*, in *Representation theory and harmonic analysis on semisimple Lie groups*, Mathematical Surveys and Monographs, vol. 31 (American Mathematical Society, Providence, RI, 1989), 101–170.
- LMT04 E. Lapid, G. Muić and M. Tadić, *On the generic unitary dual of quasisplit classical groups*, Internat. Math. Res. Notices **26** (2004), 1335–1354.
- LRS99 W. Luo, Z. Rudnick and P. Sarnak, *On the generalized Ramanujan conjecture for $\mathrm{GL}(n)$* , in *Automorphic forms, automorphic representations, and arithmetic*, Fort Worth, TX, 1996, Proceedings of Symposia in Pure Mathematics, vol. 66 (American Mathematical Society, Providence, RI, 1999), 301–310.
- Mor04 T. Moriyama, *Entireness of the spinor L -functions for certain generic cusp forms on $\mathrm{GSp}(2)$* , Amer. J. Math. **126** (2004), 899–920.
- Mui01 G. Muić, *A proof of Casselman-Shahidi's conjecture for quasi-split classical groups*, Canad. Math. Bull. **44** (2001), 298–312.
- Pia79 I. Piatetski-Shapiro, *Multiplicity one theorems*, in *Automorphic forms, representations and L -functions*, Oregon State University, Corvallis, OR, 1977, part 1, Proceedings of Symposia in Pure Mathematics, vol. 33 (American Mathematical Society, Providence, RI, 1979), 209–212.
- Pia97 I. Piatetski-Shapiro, *L -functions for GSp_4* . Olga Taussky-Todd: in memoriam, Pacific J. Math. **181** Special Issue (1997), 259–275.
- PS84 I. Piatetski-Shapiro and D. Soudry, *L and ε functions for $\mathrm{GSp}(4) \times \mathrm{GL}(2)$* , Proc. Natl Acad. Sci. USA **81** (1984), 3924–3927.
- PS87 I. Piatetski-Shapiro and D. Soudry, *On a correspondence of automorphic forms on orthogonal groups of order five*, J. Math. Pures Appl. (9) **66** (1987), 407–436.
- Ram97 D. Ramakrishnan, *On the coefficients of cusp forms*, Math. Res. Lett. **4** (1997), 295–307.
- Sar03 P. Sarnak, *Notes on the generalized Ramanujan conjectures*, Fields Institute Lectures, 2003, in *Harmonic analysis, the trace formula, and Shimura varieties*, Clay Mathematics Proceedings, vol. 4 (American Mathematical Society, Providence, RI, 2005), 659–685.
- Sha81 F. Shahidi, *On certain L -functions*, Amer. J. Math. **103** (1981), 297–355.
- Sha90 F. Shahidi, *A proof of Langlands' conjecture on Plancherel measures; complementary series for p -adic groups*, Ann. of Math. (2) **132** (1990), 273–330.
- Sha04 F. Shahidi, *On the Ramanujan conjecture for quasi-split groups*, Asian J. Math. **8** (2004), 813–836.
- Sou05 D. Soudry, *On Langlands functoriality from classical groups to GL_n* , Astérisque **298** (2005), 335–390.
- Tad86 M. Tadić, *Classification of unitary representations in irreducible representations of general linear group (non-Archimedean case)*, Ann. Sci. École Norm. Sup. (4) **19** (1986), 335–382.
- Tak R. Takloo-Bighash, *Spinor L -functions, theta correspondence, and Bessel coefficients*, Forum Math., to appear.
- Vog78 D. Vogan, *Gelfand–Kirillov dimension for Harish-Chandra modules*, Invent. Math. **48** (1978), 75–98.
- Zel80 A. Zelevinsky, *Induced representations of reductive p -adic groups II. On irreducible representations of $\mathrm{GL}(n)$* , Ann. Sci. École Norm. Sup. (4) **13** (1980), 165–210.

GENERIC TRANSFER FROM $\mathrm{GSp}(4)$ TO $\mathrm{GL}(4)$

Mahdi Asgari asgari@math.okstate.edu

School of Mathematics, Institute for Advanced Study, Princeton, NJ 08450, USA

Current address: Department of Mathematics, Oklahoma State University, Stillwater,
OK 74078-1058, USA

Freydoon Shahidi shahidi@math.purdue.edu

Department of Mathematics, Purdue University, West Lafayette, IN 47907, USA