

Menopause and Risk-Taking Behaviours: A Cross-Sectional, Online Survey

Dr Dan Reisel*, Ms Aini Kamal, Dr Sarah Glynne and Dr Louise Newson

Newson Health, Stratford-upon-Avon, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.232

Aims. Limited data suggest that negative mood symptoms in the menopause transition may be associated with a higher prevalence of alcohol misuse and other risk-taking behaviours in menopausal women. Excessive alcohol consumption can exacerbate menopausal symptoms, reduce quality of life and is associated with chronic morbidity that overlaps with the consequences of long-term oestrogen deficiency (such as osteoporosis and cardiovascular disease). The aim of this survey was to explore the impact of mental ill-health on alcohol consumption and gambling habits in menopausal women.

Methods. We constructed an anonymous survey consisting of multiple-choice and free-text questions. The survey was distributed online via social media channels on the 22 August 2023 and was open for 6 weeks. All perimenopausal and menopausal women were invited to participate. Responses were collected using the Qualtrics survey platform and analysed in Excel for descriptive statistics.

Results. 1,178 responses were submitted. One in three women reported drinking more alcohol during the perimenopause/menopause; 15% of women drink more than the recommended maximum of 14 units per week, and 24% (286) are spending up to £50 per week on alcohol. 70% (332) cited anxiety, stress, and/or depression as the reason for their increased alcohol consumption, whilst 29% (135) said they drank to alleviate menopause symptoms. Further, 5% (54) of respondents admitted gambling more since the onset of perimenopause/menopause; 43% (27) said it was due to anxiety, stress, and/or depression, whilst 13% (9) said they do so to help manage their menopause symptoms.

Conclusion. This anonymous, cross-sectional survey found evidence of an association between menopause and addiction. Increased awareness of this association should facilitate earlier recognition and more timely access to support and effective treatment for addiction, including hormone replacement therapy to treat menopausal symptoms that may underlie and/or exacerbate unhealthy lifestyle behaviours.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Neurodevelopmental Disorders and Their Association With Neurodegenerative Disorders: A Systematic Narrative Review

Dr Rebecca Restorick^{1,2*}, Dr Huw Dunstall³ and Dr Katja Umla-Runge²

¹NHS Borders, Selkirk, United Kingdom; ²Cardiff University, Cardiff, United Kingdom and ³Swansea Bay UHB, Swansea, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.233

Aims. Neurodevelopmental disorders (NDDs), such as dyslexia, dyspraxia, and dyscalculia affect cognitive function and therefore share symptomology with neurodegenerative disorders, such as

Alzheimer's disease, vascular dementia, and frontotemporal lobe dementia. The primary aim of this narrative systematic review is to ascertain if there is an association between NDDs and neurodegenerative disorders. Secondary aims are what the prevalence of NDDs within a dementia population is and what effect these early life learning disorders have on patients as they get older. It was hypothesised that NDDs would overestimate the severity of cognitive impairment, thereby increasing the severity of dementia staging, and impacting patient care.

Methods. Using a Population, Exposure, Comparator, Outcome, Setting, and Study design (PECOS) framework, keywords of "dementia", "dyslexia", "Dyspraxia/clumsy child syndrome/developmental apraxia/motor learning difficulty/disorder of attention and motor perception" and "dyscalculia/mathematical learning disability" were searched for on 4 databases (SCOPUS, OVID, Cochrane Central Register of Controlled Trials and Web of Science) from January 1, 1960 – June 10, 2022. Studies were included if they discussed both neurodegenerative and neurodevelopmental disorders or compared an intervention typically used in one disorder on the other (e.g., dementia intervention being used on neurodevelopmental disorder). Studies were excluded from grey literature articles, or if they only discussed a neurodevelopmental or neurodegenerative disorder without reference to the other, or if it included acquired, rather than neurodevelopmental dyslexia, dyscalculia, or dyspraxia.

Results. A total of 8 studies were included for narrative synthesis. The main finding was an association between dyslexia and both Alzheimer's disease and frontotemporal dementia. Many studies suggested this was due to a genetic phenotype that caused a vulnerability in the language regions of patients' cortices. There was also evidence of structural changes associated with NDDs and increased levels of grey and white matter atrophy in dementia subtypes, particularly in the language areas of the brain.

Conclusion. Due to screening and consequently formal diagnosis of neurodevelopmental disorders only recently coming into education systems, many adults currently attending memory clinics did not have a formal diagnosis. As there was limited research on dyspraxia and dementia, partly due to limited research into dyspraxia itself and without a standardized diagnostic tool for adolescents and adults, further research is needed in this area. The hypothesis of NDDs increasing the severity of dementia staging was also not supported by the literature results, and on the contrary, some studies suggested greater global preservation of cognitive function in patients with NDDs and dementia.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Decrypting the Thalamic Subnuclei and Functional Composites in Adolescents With Psychotic Experiences

Dr Sahar Riaz*, Michael O'Connor, Anurag Nasa, Dr Mary Cannon and Dr Darren Roddy

Royal College of Surgeons in Ireland, Dublin, Ireland

*Presenting author.

doi: 10.1192/bjo.2024.234

Aims. The thalamus, a dual grey matter formation within the diencephalon is thought to be involved in psychosis. It consists of distinct nuclei with specific functions. To date no study has investigated the volumes of the thalamic nuclei in young adolescents with Psychotic Experiences (PEs).