

Selenium inclusion decreases oxidative stress indicators and muscle injuries in sea bass larvae fed high-DHA microdiets

Mónica B. Betancor^{1*}, M^a José Caballero¹, Genciana Terova², Reda Saleh¹, Eyad Atalah¹, Tibiábin Benítez-Santana¹, J. Gordon Bell³ and Marisol Izquierdo¹

¹Grupo de Investigación en Acuicultura, University of Las Palmas de Gran Canaria, Instituto Universitario de Sanidad Animal, Trasmontaña s/n, 35413 Arucas, Las Palmas, Canary Islands, Spain

²Department of Biotechnology and Molecular Sciences, University of Insubria, Via Dunant 3, 21100 Varese, Italy

³Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, UK

(Submitted 20 June 2011 – Final revision received 23 November 2011 – Accepted 30 November 2011 – First published online 13 March 2012)

Abstract

The objective of the present study was to determine the effect of Se inclusion in high-DHA and vitamin E microdiets (5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight; 5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight supplemented with Se) in comparison with a control diet (1 g DHA/100 g dry weight and 150 mg vitamin E/100 g dry weight) on sea bass larval growth, survival, biochemical composition, malonaldehyde (MDA) content, muscle morphology and antioxidant enzymes (AOE), insulin-like growth factors (IGF) and myosin expression. For a given DHA and vitamin E dietary content, Se inclusion favoured larval total length and specific growth rate, and reduced the incidence of muscular lesions, MDA contents and AOE gene expression. In contrast, IGF gene expression was elevated in the 5/300 larvae, suggesting an increased muscle mitogenesis that was corroborated by the increase in mRNA copies of myosin heavy chain. The results of the present study denoted the beneficial effect of Se not only in preventing oxidative stress, as a glutathione peroxidase cofactor, but probably due to other as yet unknown physiological functions.

Key words: Sea bass larvae: Oxidative stress: DHA: Selenium

Free radicals and/or oxygen derivatives are continuously generated during regular cellular metabolism. At low concentrations, these reactive oxygen species (ROS) may be beneficial or even indispensable in processes such as defence against micro-organisms, contributing to phagocytic bactericidal activity. However, when an imbalance between ROS generation and ROS removal occurs, oxidative stress arises⁽¹⁾. The detrimental effects include oxidative damage to molecules of great biological importance, including lipids, proteins and DNA, causing alterations that produce a range of cellular damage which can ultimately lead to cell death⁽²⁾.

Several effective antioxidant systems prevent oxidative damage in fish. Among them, various antioxidant enzymes (AOE) prevent the cascade of oxidation reactions, intercepting and inactivating the reactive intermediates and closing the lipid-peroxidation catalytic cycle. This defence system includes enzymes such as catalase (CAT), superoxide dismutase (SOD)

and glutathione peroxidase (GPX)^(2,3). In addition to these enzymes, dietary micronutrients such as vitamins E and C as well as carotenoids have also been regarded as antioxidant defences in fish^(4–8). During larval stages, oxidation risks are particularly high, due to the increased metabolic rate, oxygen demand, water and long-chain PUFA (LC-PUFA) tissue contents found in fish larvae. Therefore, adequate supplementation of marine fish larvae with antioxidant elements is important when formulating diets in order to avoid *in vivo* lipid peroxidation. However, few studies have dealt with the effects of antioxidants in early fish feeding^(9–12). Among the other antioxidants, Se is an essential trace mineral in animal nutrition obtained partly from the surrounding water⁽¹³⁾, but mostly from the diet⁽¹⁴⁾. The importance of Se to oxidative stress involves its presence at the active site of the antioxidant enzyme GPX⁽¹⁵⁾, which reduces H₂O₂ at the expense of reduced glutathione⁽¹⁶⁾. In fish, studies have shown a synergistic action

Abbreviations: 1/150, 1 g DHA/100 g dry weight and 150 mg vitamin E/100 g dry weight diet; 5/300, 5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight diet; 5/300 + Se, 5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight diet supplemented with Se; AOE, antioxidant enzymes; CAT, catalase; dph, days post-hatching; DW, dry weight; GLM, general linear model; GPX, glutathione peroxidase; IGF-I, insulin-like growth factor I; IGF-II, insulin-like growth factor II; LC-PUFA, long-chain PUFA; MDA, malonaldehyde; MyHC, myosin heavy chain; ROS, reactive oxygen species; SGR, specific growth rate; SOD, superoxide dismutase; TBARS, thiobarbituric acid-reactive substances.

*Corresponding author: M. B. Betancor, fax +34 928 451143, email monica.betancor102@doctorandos.ulpgc.es

between vitamin E and Se^(17,18). Moreover, a Se deficiency may lead to reduced levels of tissue α -tocopherol in several fish species^(3,17–20). Additionally, a recent study in grouper (*Epinephelus malabaricus*) juveniles suggested a sparing effect between Se and vitamin E⁽²¹⁾.

One of the most important factors that can lead to oxidative stress in fish is their high requirement for LC-PUFA, which are high in diets and, subsequently, in fish tissue^(22,23). For instance, severe dystrophic lesions in the epaxial musculature of sea bass larvae have been related to the deleterious effect of oxidative stress due to the high ingestion of LC-PUFA, particularly DHA^(12,24). Muscle growth is an essential process during larval stages, as a massive increment of muscle fibres takes place from hatching to maturity⁽²⁵⁾ with damage to the musculature appearing to compromise larval growth^(12,24).

Muscle formation processes require the influence of growth factors and a sequence of cellular events that result in the regulation of myoblasts (myosatellite cells)^(26,27). Insulin-like growth factors I and II (IGF-I and IGF-II) are two myogenic regulatory factors which increase satellite cell proliferation and differentiation^(28,29). In various species of fish, it has been shown that hepatic^(30–33) and muscular^(32–34) IGF-I and IGF-II mRNA levels depend on feeding status. Differences in the regulation of myogenesis such as myosin isoform expression have also been observed during the earliest stages of development as well as during temperature acclimatisation^(35,36). However, nutritional regulation of the various components of the IGF signalling pathways in muscle growth in fish as well as myosin expression is not well studied.

The purpose of the present study was to investigate whether diets supplemented with Se (5 mg/kg dry weight (DW)) and vitamin E can protect sea bass larvae muscle from oxidative stress when high DHA levels are included in diets. To reach this objective, growth, survival, thiobarbituric acid-reactive substances (TBARS), fatty acid profile, α -tocopherol and Se contents, and mRNA expression levels of *CAT*, *SOD*, *GPX*, *IGF-I*, *IGF-II* and myosin heavy chain (*MyHC*) genes were determined in sea bass larvae fed diets with different LC-PUFA, Se and vitamin E contents.

Methods

Fish

All experiments were designed according to the Animal Welfare Ethics Committee guidelines of Las Palmas University. The experiment was carried out at the Grupo de Investigación en Acuicultura facilities (Telde, Canary Islands, Spain). Sea bass (*Dicentrarchus labrax*) larvae were obtained from natural spawnings from the Instituto de Acuicultura de Torre de la Sal (Castellón, Spain). Before the start of the feeding experiment, larvae were fed enriched (DHA Protein Selco®; INVE) yeast-fed rotifers until they reached 14 d post-hatching (dph). Larvae (total length 8.58 (SD 0.64) mm and dry body weight 0.36 (SD 0.0) mg) were randomly distributed into the experimental tanks at a density of 1000 larvae/tank and fed one of the experimental diets for 21 d. All tanks (170 litres light grey colour cylinder fibreglass tanks) were supplied with filtered sea water (34 g/l

salinity) at an increasing rate of 1.0–1.5 litres/min during the feeding trials. Water entered the tank from bottom to top; water quality was tested daily and no deterioration was observed. Water was continuously aerated (125 ml/min), attaining 5–8 g/l dissolved oxygen and saturation ranged between 60 and 80%. Water temperature ranged from 19.5 to 21.0°C.

Diets

For the feeding experiment, three isonitrogenous and isolipidic experimental microdiets (pellet size < 250 μ m) similar in their EPA content and different in DHA, Se and vitamin E content were formulated (Table 1) using concentrated fish oils INCROMEGA™ EPA500 and DHA500 (CRODA) as sources of EPA and DHA in TAG form and DL- α -tocopheryl acetate (Sigma-Aldrich) as a source of vitamin E. The diets were chosen based on previous trials^(12,24) and their names elected according to the level of dietary DHA and vitamin E content. A positive control diet was formulated to include 1 g DHA/100 g DW and 150 mg vitamin E/100 g DW (diet 1/150). The negative control diet consisted of 5 g DHA/100 g DW and 300 mg vitamin E/100 g DW (diet 5/300). The third diet had identical DHA and vitamin E content to the 5/300 diet, but was supplemented with Se in organic

Table 1. Formulation of the experimental diets

Ingredients	Experimental diets		
	1/150	5/300	5/300 + Se
Defatted squid powder (g/100 g)*	69.00	68.85	68.60
INCROMEGA™ EPA 500TG g/100 g (DW)†	2.80	1.80	1.80
INCROMEGA™ DHA 500TG g/100 g (DW)†	0.20	6.70	6.70
Oleic acid (%‡)	10.00	4.50	4.50
Soya lecithin§	2.00	2.00	2.00
Gelatin	3.00	3.00	3.00
Attractants¶	3.00	3.00	3.00
Taurine**	1.50	1.50	1.50
Vitamin premix‡‡	6.00	6.00	6.00
Mineral premix‡‡	2.50	2.50	2.50
Sel-Plex 2000® (g/100 g)§§	–	–	0.25
Vitamin E mg/100 g (DW)¶¶	–	0.15	0.15

DW, dry weight; 1/150, 1 g DHA/100 g DW and 150 mg vitamin E/100 g DW diet; 5/300, 5 g DHA/100 g DW and 300 mg vitamin E/100 g DW diet; 5/300 + Se, 5 g DHA/100 g DW and 300 mg vitamin E/100 g DW diet supplemented with Se.

* Riber and Son.

† Croda Chemicals Europe.

‡ Merck.

§ Acrofarma.

|| Panreac.

¶ Attractant premix supplied per 100 g diet: inosine-5-monophosphate, 500.0 mg; betaine, 660.0 mg; L-serine, 170.0 mg; L-phenylalanine, 250.0 mg; DL-alanine, 500.0 mg; L-sodium aspartate, 330.0 mg; L-valine, 250.0 mg; glycine, 170.0 mg.

** Sigma-Aldrich.

‡‡ Vitamin premix supplied per 100 g diet: cyanocobalamin, 0.03 mg; astaxanthin, 5.0 mg; folic acid, 5.4 mg; pyridoxine-HCl, 17.2 mg; thiamin, 21.7 mg; riboflavin, 72.5 mg; calcium-pantothenate, 101.5 mg; *p*-aminobenzoic acid, 145.0 mg; nicotinic acid, 290.1 mg; *myo*-inositol, 1450.9 mg; retinol acetate, 0.2 mg; ergocalciferol, 3.6 mg; menadione, 17.3 mg; α -tocopheryl acetate, 150.0 mg.

‡‡ Mineral premix supplied per 100 g diet: NaCl, 215.133 mg; MgSO₄·7H₂O, 677.545 mg; NaH₂PO₄·H₂O, 381.453 mg; K₂HPO₄, 758.949 mg; Ca(H₂PO₄)₂·H₂O, 671.610 mg; FeC₆H₅O₇, 146.884 mg; C₆H₅O₃·0.5Ca, 1617.210 mg; Al₂(SO₄)₃·6H₂O, 0.693 mg; ZnSO₄·7H₂O, 14.837 mg; CuSO₄·5H₂O, 1.247 mg; MnSO₄·H₂O, 2.998 mg; KI, 0.742 mg; CoSO₄·7H₂O, 10.706 mg.

§§ Sel-Plex 2000, 2000 mg Se/kg; Alltech.

¶¶ DL- α -Tocopheryl acetate; Sigma-Aldrich.

form (diet 5/300 + Se, Sel-Plex[®] 2000, 2000 mg/kg; Alltech). The protein source was derived from squid meal, naturally containing 14% lipid, defatted three consecutive times to allow complete control of the microdiet fatty acid profile. No other ingredients were defatted due to their poor lipid content. The squid meal was defatted with a chloroform:squid meal ratio of 3:1, rinsed and dried in an oven at 37°C until complete solvent evaporation. EPA500 and DHA500 were added in different quantities to the defatted meal (2.4% lipid content) to obtain the desired ratios. Oleic acid (Merck) was added to equalise the lipid content in each diet (Table 1) and soyabean lecithin (Acrofarma) was included as a source of phospholipids (Table 1). The microdiets were processed as described previously⁽³⁷⁾. Briefly, the squid powder and water-soluble components were mixed, followed by the lipid and fat-soluble vitamins and, before adding gelatine (Panreac), dissolved in warm water, as a binder. The paste was pelleted and oven dried at 38°C for 24 h. The pellets were ground and sieved to obtain particle size below 250 µm. To avoid peroxidation, the diets were stored under N₂ at -20°C until use. The diets were analysed for proximate and fatty acid composition on a dry basis and manually supplied; fourteen times/d at 45 min intervals from 09.00 to 19.00 hours. Daily feed supplied was 2, 2.5 and 3 g/tank during the first, second and third week of feeding, respectively.

Growth and survival

Final survival was determined by counting live larvae at the beginning and end of the experiment. Growth was determined by measuring dry body weight (105°C for 24 h) and total length (Profile Projector V-12A Tokyo; Nikon) of thirty fish/tank at the beginning, middle and end of the trial.

Biochemical analysis

All the remaining larvae in each tank were washed with distilled water, sampled and kept at -80°C for biochemical composition, TBARS, Se and vitamin E analysis after 12 h of starvation at the end of the trial. Moisture⁽³⁸⁾, protein⁽³⁸⁾ and lipid⁽³⁹⁾ contents of larvae and the diets were analysed.

Total lipid fatty acid analysis. Fatty acid methyl esters were obtained by transmethylation of total lipids as described by Christie⁽⁴⁰⁾. Fatty acid methyl esters were separated by GLC, quantified by flame ionisation detection (GC-14A; Shimadzu) under the conditions described previously⁽⁴¹⁾ and identified by comparison with previously characterised standards and GLC-MS.

Determination of vitamin E content. Vitamin E concentrations were determined in the diets and total larvae using HPLC at the University of Stirling (Scotland, UK). Samples were weighed, homogenised in pyrogallol and saponified as described by McMurray *et al.*⁽⁴²⁾ for the diets or according to Cowey *et al.*⁽⁴³⁾ for larval tissues. HPLC analysis was performed using a 150 × 4.60 mm reversed-phase Luna 5 µm C18 column (Phenomenex). The mobile phase was 98% methanol supplied at a flow rate of 1.0 ml/min. The effluent from the column was monitored at a wavelength of 293 nm and quantification

achieved by comparison with (+)-α-tocopherol (Poole) as the external standard.

Selenium determination. Total Se concentration was measured in total larvae and the diets, and analyses were carried out at the University of Stirling (Scotland, UK). Samples were acidified in a microwave digester (MarsXpress; CEM) with 5 ml of 69% pure HNO₃, then poured after digestion into a 10 ml volumetric flask and made up to volume with distilled water. A total of 0.4 ml of this solution were then added to a 10 ml sample tube, 10 µl of the internal standard (Ga and Sc, 10 ppm) included and 0.3 ml of methanol added. The tubes were made up to volume with distilled water and total Se measured by collision/reaction by inductively coupled plasma MS (Thermo Scientific) using argon and hydrogen as the carrier gas.

Measurement of thiobarbituric acid-reactive substances. TBARS from triplicate samples were determined using a method adapted from that used by Burk *et al.*⁽⁴⁴⁾. Approximately 20–30 mg of larval tissue per sample were homogenised in 1.5 ml of 20% TCA (w/v) containing 0.05 ml of 1% butylated hydroxytoluene in methanol. To this 2.95 ml of freshly prepared 50 mM-thiobarbituric acid solution were added before mixing and heating for 10 min at 100°C. After cooling, protein precipitates were removed by centrifugation (Sigma 4K15) at 2000 g and the supernatant was read in a spectrophotometer (Evolution 300; Thermo Scientific) at 532 nm. Absorbance was recorded against a blank at the same wavelength. The concentration of thiobarbituric acid-malonaldehyde (MDA), expressed as µmol MDA/g tissue was calculated using an extinction coefficient of 0.156 cm/µM.

Histopathological sampling

From each tank, thirty larvae were collected every 7 d from the beginning of the feeding trial, fixed in 10% buffered formalin for 1–2 d, dehydrated through graded alcohols, then xylene and finally embedded in paraffin wax. Then, six paraffin blocks containing five larvae/tank were sectioned at 3 µm, and stained with haematoxylin and eosin for histopathological evaluation⁽⁴⁵⁾.

From each tank, ten larvae were fixed for 24 h at 4°C in 2.5% glutaraldehyde in 0.2 M-phosphate buffer (pH 7.2). Samples were then rinsed in phosphate buffer and post-fixed for 1 h in 2% osmium tetroxide in 0.2 M-potassium ferrocyanide. Each larva was then embedded in an Epon/Araldite resin block. Thick serial transverse and longitudinal sections of the larvae were cut at 2 µm, stained with toluidine blue and examined by light microscopy⁽⁴⁶⁾. Thin sections were cut at 50 nm and stained with lead citrate before observing with a ZEISS EM 910 transmission electron microscope (Zeiss) at the Electron Microscope Service of the University of Las Palmas de Gran Canaria.

RNA extraction and quantitative RT-PCR

Molecular biology analysis was carried out at the University of Insubria (Varese, Italy). Total RNA was extracted from sea bass larvae (approximately 200 mg; pool per tank), using the PureYield RNA Midiprep System (Promega). The quantity and

purity of RNA was assessed by a spectrophotometer. Visualisation on 1% agarose gel stained with ethidium bromide showed that RNA was not degraded. After DNase treatment (Invitrogen), 3 µg of total RNA were reverse transcribed into complementary DNA in a volume of 12 µl, including 1 µl of oligo-dT16 primer (50 pmol) and 1 µl of 10 mM-deoxynucleotide triphosphates. This mix was heated at 65°C for 5 min and chilled on ice before 4 µl of 5 × reverse transcription buffer, 2 µl of 0.1 M-dithiothreitol, 1 µl RNase out and 1 µl of Moloney murine leukaemia virus was added. After incubation at 37°C for 50 min, the reaction was stopped by heating at 75°C for 15 min.

PCR primer sequences used for the PCR amplification of the complementary DNA of target genes were *CAT*, *SOD*, *GPX*, *IGF-I*, *IGF-II* and *MyHC*. To perform PCR, a 4 µl aliquot of complementary DNA was amplified using 25 µl GoTaq Green Master Mix (Promega) in 50 µl of final volume and 50 pmol of each designed primer.

A total of thirty-one PCR amplification cycles (eight touch-down) were performed for all primer sets, using an automated Thermal Cycler (MyCycler; BioRad). An aliquot of each sample was then subjected to electrophoresis on a 1% agarose gel in 1 × Tris-acetate-EDTA buffer (Bio-Rad) and bands were detected by ethidium bromide staining. Samples were run with a 100 bp–1.5 kb DNA ladder to control the molecular weight of DNA. The negative control (a reaction mixture without complementary DNA) confirmed the absence of genomic contamination. The PCR products from each primer set amplification were cloned using the pGEM[®]-T easy vector (Promega) and subsequently sequenced in both directions (T7 and SP6).

TaqMan[®] real-time RT-PCR was performed on a StepOne Real Time PCR System (Applied Biosystems) using Assays-by-DesignSM PCR primers (Applied Biosystems) and gene-specific fluorogenic probes. Primer sequences and Taqman[®] probes of the target genes were as follows:

Target gene: sea bass *CAT*

Forward primer: 5'-ATGGTGTGGACTTCTGGAG-3'

Reverse primer: 5'-GCTGAACAAGAAAGACACCTGATG-3'

Taqman[®] probe: 5'-CAGACACTCAGGCCTCA-3'

Target gene: sea bass *SOD*

Forward primer: 5'-TGGAGACCTGGGAGATGTAACCTG-3'

Reverse primer: 5'-TCTTGTCCGTGATGTCGATCTTG-3'

Taqman[®] probe: 5'-CAGGAGGAGATAACATTG-3'

Target gene: Sea bass *GPX*

Forward primer: 5'-AGTTAATCCGGAATTCGTGAG-3'

Reverse primer: 5'-AGCTTAGCTGTCAGGTCGTAAAC-3'

Taqman[®] probe: 5'-AATGGCTGGAAACGTG-3'

Target gene: Sea bass *IGF-I*

Forward primer: 5'-GCAGTTTGTGTGGAGAGAGA-3'

Reverse primer: 5'-GACCGCCGTGCATTGG-3'

Taqman[®] probe: 5'-CTGTAGGTTTACTGAAATAAAA-3'

Target gene: Sea bass *IGF-II*

Forward primer: 5'-TGCAGAGACGCTGTGTGG-3'

Reverse primer: 5'-GCCTACTGAAATAGAAGCCTCTGT-3'

Taqman[®] probe: 5'-CAAACCTGCAGCGCATCC-3'

Target gene: Sea bass *MyHC*

Forward primer: 5'-TGGAGAAGATGTGCCGTACTCT-3'

Reverse primer: 5'-CGTGTGATGATTTGACGGACATTT-3'

Taqman[®] probe: 5'-AACTGAGTGAAGTGAAGACC-3'

Data from TaqMan[®] PCR runs were collected using ABI's Sequence Detector Program. Cycle threshold values corresponded to the number of cycles at which fluorescence emission monitored in real time exceeded the threshold limit. The cycle threshold values were used to create standard curves to serve as a basis for calculating the absolute amounts of mRNA in total RNA. To reduce pipetting errors, master mixes were prepared to set up duplicate reactions (2 × 30 µl) for each sample.

Calculations

Larval survival was determined by comparing the number of larvae at the beginning of the trial with the larvae number measured in individual tanks at 35 dph to which the average number of larvae sampled from the tanks during the trial was added. Percentage survival could then be calculated for each tank to get a mean and standard deviation per treatment. The incidence of muscular lesions was calculated as the percentage of injured larvae per tank compared with the total larvae observed, with standard deviation referring to deviation among the tanks. Specific growth rate (SGR) was calculated as

$$\text{SGR} = (\ln W_1 - \ln W_0) \times 100 / t_2 - t_1,$$

where W_0 and W_1 are the initial and final DW (tank means), respectively, and $t_2 - t_1$ is the time interval in days between the beginning and end of the experimental trial (21 d).

Statistical analysis

Survival, growth and molecular biology data were tested for normality and homogeneity of variances with Levene's test. Where necessary, data were log transformed before further statistical analysis. The χ^2 test was employed for incidence of muscular lesions and TBARS content. Survival, growth and biochemical analysis data were treated using one-way ANOVA and molecular biology results were treated using a general linear model (GLM). Means were compared by Duncan's test. Results are presented as means and standard deviations. The tank was considered as the experimental unit, except for the estimation of the incidence of muscular lesions, where each individual larva was considered as a unit. For percentage data (final survival), arcsine transformation was performed before the analysis. For the analysis of one-way ANOVA, the following GLM was used:

$$Y_{ij} = \mu + \alpha_i + \epsilon_{ij},$$

where Y_{ij} is the mean value of the tank, μ is the mean population, α_i is the fixed effect of the diet and ϵ_{ij} is the residual error. For the analysis of molecular biology data, a two-variable GLM was employed to analyse possible interactions between treatment and time:

$$Y_{ijk} = \mu + \alpha_i + \delta_j + (\alpha\delta)_{ij} + \epsilon_{ijk},$$

where Y_{ijk} is the mean value of the tank, μ is the mean population, α_i is the fixed effect of the diet, δ_j is the fixed effect of the time, $(\alpha\delta)_{ij}$ is the interaction between diet and time and ϵ_{ijk} is the residual error. Significance was accepted at

$P \leq 0.05$. Statistical analysis was performed using SPSS software (SPSS for Windows 14.0; SPSS, Inc., 2005).

Results

The diet containing about 1% DHA (1/150) showed a higher monoenoic fatty acid level than the diets containing 5% DHA (5/300 and 5/300 + Se) due to a higher oleic acid content in the former diet (Table 2). Elevation of dietary DHA (5/300 diet) increased *n*-3 and *n*-3 LC-PUFA fatty acid contents, as well as the *n*-3:*n*-6 ratio. Vitamin E levels were more than two times higher in the diets containing 300 mg/100 g compared with the control diet (1/150) (Table 3). Se contents differed among the dietary treatments, with a higher level of this mineral found in the diet supplemented with Se compared with the others (Table 3).

All the experimental diets were well accepted by the larvae. Dietary increase of vitamin E or Se did not significantly affect larval survival ($P=0.158$; Table 5). The highest larval growth, in terms of total length, was found in the larvae fed the positive control diet containing the lowest vitamin E and DHA contents (diet 1/150). However, increases in vitamin E and DHA (diet 5/300) significantly reduced the larval growth ($P=0.001$), whereas Se inclusion (diet 5/300 + Se) significantly increased this parameter ($P=0.02$; Table 5). The average SGR was higher ($P=0.024$) in the larvae fed the 1/150 diet, although no differences were found in the larvae fed the 5/300 + Se diet.

In terms of fatty acid composition, larvae fed the 5/300 diet resulted in a higher concentration of *n*-3 and *n*-3 LC-PUFA, reflecting the higher content of these components in the diets (Table 4). However, larvae fed the 1/150 diet showed a higher retention rate of DHA (279.26%) and EPA (68.24%) compared with larvae fed the 5/300 (73.48 and 43.93%, respectively) and 5/300 + Se diets (77.00 and 44.75%, respectively) and also a higher content of 18:1*n*-9, displaying levels similar to those found in the former microdiet. In contrast, the 5/300 larvae showed a higher retention rate of arachidonic acid (about 50%) compared with the 1/150 larvae (29.83%).

The level of lipid oxidation, as indicated by the MDA content ($\mu\text{mol/g}$ larval tissues), was significantly higher ($P=0.001$) in the larvae fed the diets with the highest DHA content. Nevertheless, the inclusion of Se showed a beneficial effect preventing the formation of H_2O_2 , as denoted by the decrease in MDA levels (Table 5). The lowest peroxidation level was observed in the larvae fed the 1/150 diet. Despite the increase in dietary α -tocopherol in the 5/300 and 5/300 + Se diets, the content of this nutrient in larvae did not significantly increase in comparison with larvae fed the 1/150 diet ($P=0.601$; Table 5). Therefore, the retention rate of dietary vitamin E in the larvae fed the 5/300 (17.87%) and 5/300 + Se (17.38%) diets was lower than the 1/150 larvae (44.68%). Sea bass larvae fed the diets supplemented with Se showed a significantly higher ($P=0.001$) content of this mineral compared with the other larvae, this level being 1.7 times higher than the larvae fed the 1/150 diet and 2.4 times higher than the 5/300 larvae (Table 5). Thus, although Se contents in the 1/150 and 5/300 diets were similar, the larvae fed the latter diet showed lower Se levels together with the highest TBARS value.

Table 2. Main fatty acids (% total fatty acids) of the experimental diets fed to sea bass larvae

	Diets		
	1/150	5/300	5/300 + Se
14:0	1.54	1.26	0.78
14:1 <i>n</i> -7	0.15	0.25	0.06
14:1 <i>n</i> -5	0.22	0.35	0.09
15:0	0.28	0.43	0.15
15:1 <i>n</i> -5	0.02	0.14	ND
16:0 ISO	0.14	0.23	0.07
16:0	7.86	5.59	5.15
16:1 <i>n</i> -7	3.59	2.26	2.00
16:1 <i>n</i> -5	0.19	0.23	0.11
16:2 <i>n</i> -4	0.32	0.39	0.25
17:0	1.21	0.82	0.65
16:3 <i>n</i> -3	0.08	0.12	0.07
16:4 <i>n</i> -3	0.09	0.13	0.10
18:0	1.29	2.29	2.21
18:1 <i>n</i> -9 + <i>n</i> -7	55.70	31.12	29.67
18:1 <i>n</i> -5	0.72	0.46	0.45
18:2 <i>n</i> -9	0.25	0.13	0.12
18:2 <i>n</i> -6	7.40	6.99	6.87
18:2 <i>n</i> -4	0.46	0.28	0.26
18:3 <i>n</i> -6	0.11	0.11	0.10
18:3 <i>n</i> -4	0.13	0.10	0.09
18:3 <i>n</i> -3	0.72	0.83	0.82
18:4 <i>n</i> -3	0.83	0.94	1.00
18:4 <i>n</i> -1	0.08	0.08	0.09
20:0	0.10	0.31	0.29
20:1 <i>n</i> -9 + <i>n</i> -7	1.10	1.53	1.45
20:1 <i>n</i> -5	0.05	0.12	0.11
20:2 <i>n</i> -9	0.05	0.04	0.02
20:2 <i>n</i> -6	0.09	0.21	0.21
20:3 <i>n</i> -6	0.09	0.13	0.13
20:4 <i>n</i> -6	0.71	1.57	1.50
20:3 <i>n</i> -3	0.07	0.18	0.16
20:4 <i>n</i> -3	0.32	0.52	0.55
20:5 <i>n</i> -3	8.66	11.04	12.20
22:1 <i>n</i> -11	0.17	0.51	0.50
22:1 <i>n</i> -9	0.08	0.25	0.25
22:4 <i>n</i> -6	0.02	0.19	0.20
22:5 <i>n</i> -6	0.19	1.75	1.78
22:5 <i>n</i> -3	0.32	1.29	1.43
22:6 <i>n</i> -3	4.58	24.55	27.95
Saturated	12.28	10.70	9.22
Monoenoics	61.99	37.23	34.69
<i>n</i> -3	15.68	39.61	44.27
<i>n</i> -6	8.61	11.14	10.85
<i>n</i> -9	57.19	33.10	31.54
<i>n</i> -3 LC-PUFA	13.96	37.58	42.28
ARA:EPA	0.08	0.14	0.12
EPA:DHA	1.89	0.45	0.44
Oleic:DHA	12.15	1.27	1.06
Oleic: <i>n</i> -3 LC-PUFA	3.99	0.83	0.70
<i>n</i> -3: <i>n</i> -6	1.82	3.56	4.08

1/150, 1 g DHA/100 g dry weight and 150 mg vitamin E/100 g dry weight diet; 5/300, 5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight diet; 5/300 + Se, 5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight diet supplemented with Se; ND, not determined; LC-PUFA, long-chain PUFA; ARA, arachidonic acid.

Histopathological examinations revealed the presence of lesions affecting the larval axial musculature. These lesions showed the typical features of necrotic degeneration of muscle, characterised by marked eosinophilia, loss of striations and adjacent nucleus. The incidence of muscular lesions increased with an increase in DHA dietary content (Table 5).

Table 3. Gross composition, α -tocopherol and selenium content in the experimental diets fed to sea bass larvae

(Mean values and standard deviations)

	1/150		5/300		5/300 + Se	
	Mean	SD	Mean	SD	Mean	SD
Protein (%)	66.79	0.52	67.77	0.09	66.24	0.36
Ash (%)	4.49	0.11	4.76	0.12	4.87	0.13
Moisture (%)	10.31	0.46	9.99	0.28	9.09	0.27
Lipids (% DW)	14.98	0.56	15.80	0.02	15.97	0.51
α -Tocopherol (μ g/g DW)	1410.12	38.77	3033.01	43.33	3217.37	14.45
Se (μ g/mg)	1.54	0.12	1.33	0.26	6.27	0.26

DW, dry weight; 1/150, 1 g DHA/100 g DW and 150 mg vitamin E/100 g DW diet; 5/300, 5 g DHA/100 g DW and 300 mg vitamin E/100 g DW diet; 5/300 + Se, 5 g DHA/100 g DW and 300 mg vitamin E/100 g DW diet supplemented with Se.

However, inclusion of Se was shown to reduce this incidence to almost half.

In semithin sections, more detailed features of these muscular lesions could be observed. In the severely damaged fibres, a coagulation of the muscular proteins could be observed as a darkening of the surrounding fibres due to hypercontraction (Fig. 1(a)). In initial, mild stages of the condition, an increase in the presence of vacuoles within the fibres was observed, together with the loss of the shape of muscular fibres, alteration of sarcoplasmic membranes and variation in the diameter of fibres (Fig. 1(b)).

Transmission electron microscopy showed muscle degeneration with the presence of hydropic and autophagic vacuoles, considered secondary lysosomes, within some affected fibres in the larvae fed the 5/300 diet (Fig. 2(a)). Altered mitochondria, observed as swollen double membrane organelles, were seen in the affected fibres and presented loss of the cristae (Fig. 2(a)) in contrast to the normal ones (Fig. 2(b)). Additionally, satellite cells were observed to be attached to the existing damaged muscle fibres under the basal lamina (Fig. 2(c)).

The general pattern of antioxidant enzyme gene expression in all groups of sea bass larvae was characterised by a rapid increase between 14 and 26 dph, followed by a decrease back to the levels slightly higher than those observed at 14 dph, by 29 dph (Fig. 3(a)–(c)). The only exception to this trend was observed in GPX gene expression, where sea bass larvae fed the 1/150 diet had a lower mRNA level of this enzyme at 35 dph in comparison with that at 14 dph (Fig. 3(c)). CAT gene expression was higher in the larvae fed the diets containing a high content of DHA at all sampling points, although no statistical differences were observed (Fig. 3(a)). However, GLM analysis showed differences between all the three treatments ($P=0.001$; Fig. 3(a); Table 6). The SOD mRNA copy number was significantly higher ($P=0.004$) at 35 dph in the 5/300 larvae (Fig. 3(b)). GPX expression level was highest in the larvae fed the diets containing a high level of DHA compared with larvae fed low DHA levels. Nevertheless, the larvae fed the diets supplemented with Se showed a lower number of GPX mRNA copies, comparable with the larvae fed the 1/150 diet (Fig. 3(c)).

Regarding the IGF genes, IGF-I mRNA copy number increased from 14 to 26 dph in all treatments, showing

a decrease at day 35 in the 1/150 and 5/300 + Se larvae. In contrast, the 5/300 larvae showed an increasing IGF-I expression levels throughout the experimental trial, with significantly higher levels at 35 dph ($P=0.006$; Fig. 3(d)). No significant differences were observed among the treatments taking into account the whole experimental period (Fig. 3(d); Table 6). The mRNA levels of IGF-II followed a similar increasing pattern in the 5/300 and 5/300 + Se larvae from 14 to 26 dph, in contrast to the 1/150 larvae. At 26 dph, the 5/300 + Se larvae showed a marked decrease in IGF-II expression, whereas in the 5/300 larvae, a steady increase could be observed. In the larvae fed the 1/150 diet, a decrease in IGF-II expression could be observed at all sampling points (Fig. 3(e)).

MyHC expression levels were elevated in the larvae fed the 5/300 diet, with a lower expression of this gene mRNA copy number was found at 26 dph, when Se was included in the diet ($P=0.007$). At 35 dph, no statistical differences were observed when Se was added to the 5/300 diet, with the lowest values in the 1/150 larvae (Fig. 3(f)). GLM analysis showed significant differences between the 5/300 larvae and the larvae fed the other dietary treatments ($P=0.001$; Fig. 3(f); Table 6).

Interactions were found in the gene expression between dietary treatment and time within the experimental trial for SOD and IGF-I (Table 6), indicating that the increase in the expression of these genes may be induced for the larval stage and the effect of the diet. However, IGF-I showed no differences in the expression among the treatments or during the whole experimental trial (Table 6).

Discussion

The aim of the present study was to evaluate the oxidative status of sea bass larvae when Se and vitamin E were included in the diet, at high DHA level inclusion. Previously, Betancor *et al.*^(12,24) provided evidence of the appearance of muscular dystrophy in sea bass larvae when fed high DHA, implying that an excessive production of free radicals was present. Furthermore, the same authors showed that an increase in vitamin E alone could not prevent its adverse effects. Se and vitamin E have different but complementary biochemical functions which may allow these nutrients to interact physiologically⁽⁴⁷⁾.

Table 4. Main fatty acid compositions of total lipids from sea bass larvae fed the experimental diets for 21 d (% total fatty acids)

(Mean values and standard deviations)

	Initial		1/150		5/300		5/300 + Se		F
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
14:0	1.08	0.01	0.93	0.06	0.76	0.03	0.71	0.03	7.092
14:1n-7	0.33	0.01	0.08	0.01	0.08	0.03	0.09	0.04	0.472
14:1n-5	0.07	0.00	0.07	0.02	0.04	0.01	0.06	0.00	1.407
15:0	0.18	0.12	0.63	0.29	0.98	1.22	0.22	0.05	2.126
15:1n-5	0.08	0.00	0.12	0.10	0.09	0.06	0.06	0.00	0.634
16:0	16.4	3.31	17.51	2.44	17.61	0.25	16.56	0.42	0.152
16:1n-7	9.67	1.75	2.02	0.14	1.60	0.06	1.56	0.00	9.531
16:1n-5	0.54	0.06	0.25	0.01	0.25	0.01	0.29	0.10	1.788
16:2n-6	0.92	0.23	0.30	0.00	0.36	0.04	0.37	0.02	1.025
16:2n-4	0.89	0.17	0.93	0.40	0.96	0.15	0.81	0.06	0.248
17:0	0.98	0.11	0.91 ^a	0.10	0.80 ^a	0.03	0.75 ^b	0.00	15.532
16:3n-3	0.06	0.00	0.12	0.01	0.14	0.03	0.12	0.03	0.764
16:3n-1	0.07	0.01	0.10	0.02	0.54	0.10	0.64	0.05	0.589
16:4n-3	0.41	0.16	0.62	0.35	0.44	0.10	0.56	0.03	0.573
18:0	7.34	0.72	11.66	3.41	12.29	0.36	11.37	0.39	0.265
18:1n-9	16.14	2.01	26.35 ^a	4.87	20.92 ^b	0.51	19.36 ^b	0.32	11.452
18:1n-7	6.27	0.48	4.85	0.29	4.44	0.39	3.98	0.29	0.900
18:1n-5	0.58	0.17	0.62	0.24	0.49	0.07	0.54	0.02	0.626
18:2n-6	3.75	1.14	4.23 ^a	0.08	3.90 ^b	0.17	3.75 ^b	0.08	11.072
18:2n-4	0.19	0.09	0.04	0.05	0.06	0.01	0.07	0.00	1.126
18:3n-6	0.17	0.08	0.43	0.01	0.38	0.04	0.32	0.02	0.399
18:3n-4	0.09	0.01	0.07	0.04	0.06	0.02	0.08	0.00	0.225
18:3n-3	0.8	0.27	0.32 ^b	0.05	0.44 ^{a,b}	0.04	0.49 ^a	0.03	13.717
18:4n-3	0.22	0.09	0.29	0.06	0.29	0.13	0.37	0.04	2.413
20:0	0.18	0.08	0.38	0.19	0.47	0.01	0.47	0.08	0.647
20:1n-9 + n-7	1.97	0.51	1.83	0.00	1.77	0.06	1.77	0.21	0.063
20:1n-5	0.49	0.26	0.26	0.15	0.13	0.01	0.16	0.00	1.268
20:2n-6	0.71	0.38	0.50	0.13	0.65	0.08	0.67	0.07	1.881
20:3n-6	0.24	0.12	0.08	0.01	0.08	0.00	0.08	0.01	0.263
20:4n-6	3.34	0.72	2.38 ^b	0.04	3.07 ^a	0.22	3.03 ^a	0.03	9.277
20:3n-3	0.09	0.00	0.12	0.09	0.15	0.04	0.14	0.01	0.304
20:4n-3	0.34	0.11	0.14	0.00	0.16	0.01	0.16	0.00	0.125
20:5n-3	7.60	0.33	5.91 ^a	1.18	4.85 ^b	0.20	5.46 ^{a,b}	0.36	0.063
22:1n-11	0.14	0.09	0.46 ^a	0.30	0.19 ^b	0.08	0.23 ^b	0.01	177.782
22:1n-9	0.33	0.17	0.26	0.15	0.26	0.06	0.23	0.08	1.548
22:5n-6	0.58	0.28	1.09	0.09	1.26	0.07	1.40	0.03	1.074
22:5n-3	1.58	0.33	0.64	0.26	0.57	0.06	0.69	0.02	0.347
22:6n-3	14.22	4.36	12.79 ^b	0.37	18.04 ^{a,b}	1.19	21.80 ^a	0.94	11.845
Saturated	26.17	5.43	32.01	6.29	32.91	1.54	30.08	0.76	0.333
Monoenoics	36.62	3.61	36.73 ^a	4.51	30.20 ^b	0.55	28.26 ^b	0.39	23.716
n-3	25.39	2.48	20.94 ^c	2.23	25.11 ^b	1.43	29.78 ^a	1.25	26.669
n-6	9.70	1.22	8.45	0.97	9.80	0.47	9.74	0.04	1.562
n-9	18.81	4.97	28.44 ^a	4.72	23.08 ^b	0.39	21.58 ^c	0.04	12.277
n-3 LC-PUFA	23.74	5.94	12.92 ^c	7.45	23.78 ^b	1.42	28.25 ^a	1.30	73.731
Oleic:DHA	1.13	0.21	1.76 ^a	0.08	1.16 ^b	0.10	0.89 ^c	0.05	46.781
Oleic:n-3 LC-PUFA	0.68	0.06	2.58 ^a	1.86	0.88 ^b	0.07	0.69 ^b	0.04	12.042
n-3:n-6	2.62	0.71	2.51	0.55	2.56	0.05	3.06	0.12	0.776

1/150, 1 g DHA/100 g dry weight and 150 mg vitamin E/100 g dry weight diet; 5/300, 5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight diet; 5/300 + Se, 5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight diet supplemented with Se; LC-PUFA, long-chain PUFA.

^{a,b,c} Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$).

Se contents in the 5/300 + Se diet were adjusted according to the levels found in copepods^(48,49), natural live prey of marine fish larvae, although they were higher than those recommended by the National Research Council⁽⁵⁰⁾ for juveniles of other fish species. Since marine fish larvae have a rapid growth rate, it is possible that they may well have a higher requirement than the juveniles which have been used in most of the requirement studies quoted by the National Research Council. In addition, the Se level did not seem to be excessive, as the larvae fed this diet did not show reduced growth in comparison with the

larvae fed the same vitamin E and DHA contents. Reduced growth is one of the first symptoms occurring when excessive levels of Se are fed to fish^(11,21,51-53). Moreover, the Se source used in the present study was derived from yeast, making it less likely to be toxic as Se toxicity is highly dependent on its speciation⁽⁵⁴⁾, with mineral Se being more toxic than organic Se^(51,55).

A dose-dependent effect of dietary vitamin E on larval tissue concentration was not observed in the present study, with the highest vitamin E content found in the larvae fed the lowest

Table 5. Sea bass larvae performance and levels of lipid peroxidation products (thiobarbituric acid-reactive substances (TBARS)), vitamin E (α -tocopherol) and selenium content of sea bass larvae at the beginning and after eating the experimental diets for 21 d (Mean values and standard deviations)

	Diets							
	Initial		1/150		5/300		5/300 + Se	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Results of the dietary trial								
Larval total length (mm)	8.58	0.64	12.60 ^a	0.93	10.89 ^c	1.24	11.35 ^b	1.29
Larval dry weight (mg)	0.36	0.00	1.46	0.47	0.94	0.05	1.08	0.10
SGR	–	–	6.79 ^a	1.15	4.55 ^b	0.27	5.24 ^{a,b}	0.44
Survival (%)	–	–	60.51	9.10	48.42	4.00	49.03	8.02
Incidence of muscular lesions (%)	–	–	15.7 ^b	14.14	52.63 ^a	15.93	27.6 ^{a,b}	6.94
Vitamin E (α -tocopherol) (μ g/g dry weight)	111.45	43.26	630.24	12.39	542.10	80.51	559.23	88.58
TBARS (nmol/g dry weight)	62.85	0.61	166.62 ^c	25.08	2402.15 ^a	67.91	282.29 ^b	92.48
Se (μ g/mg dry weight)	1.38	0.10	1.58 ^b	0.12	1.11 ^c	0.31	2.65 ^a	0.27

1/150, 1 g DHA/100 g dry weight and 150 mg vitamin E/100 g dry weight diet; 5/300, 5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight diet; 5/300 + Se, 5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight diet supplemented with Se; SGR, specific growth rate.

^{a,b,c} Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$).

level of vitamin E (150 mg/100 g DW). These results are in contrast with previous reports where vitamin E concentrations in fish were linked to dietary input^(56,57). It is also noteworthy that vitamin E levels were influenced by the dietary DHA ratio, being lower in the larvae fed the diets containing the higher amount of DHA (5/300 diet). This indicates that more vitamin E was being utilised as an antioxidant in the larvae fed higher DHA levels to protect tissue lipids from an increased oxidation risk. Consequently, α -tocopherol was accumulated at lower amounts in larval tissues. These results match those of previous reports where vitamin E concentration in juvenile or adult fish was lower when high contents of n -3 LC-PUFA were included in the diets⁽⁵⁷⁾. In addition, the low DHA retention rate observed in the 5/300 larvae could probably be due to the high *in vivo* lipid oxidation induced by this fatty acid.

In agreement with this, a dietary DHA increase resulted in higher levels of MDA, whereas Se supplementation improved protection against peroxidation by decreasing TBARS values. Moreover, Se incorporation rate was very low in the 5/300 + Se larvae (42.26%) in contrast to the 1/150 (100%) and 5/300 (83.46%) larvae, suggesting that this mineral was being used at the active sites of the antioxidant enzyme GPX⁽¹⁵⁾. The synergism between tocopherol and Se has previously been observed in rainbow trout (*Oncorhynchus mykiss*) and

salmon (*Salmo salar*) using diets deficient in vitamin E, Se or both^(17,18). In the present study, tocopherol levels were high enough to avoid a deficiency in this nutrient, suggesting that vitamin E addition as the sole antioxidant is not sufficient enough to control lipid peroxidation when high levels of DHA are included in fish larval diets. Therefore, the role of tocopherol as an effective antioxidant depends on the extent of oxidative stress in fish and is thus related to the degree of unsaturation of dietary fatty acids.

The availability of the literature on the activities of AOE in fish is mainly focused on pollutant detoxification^(58,59) or developmental aspects^(9,60). A few reports exist concerning the effect of dietary components on their activity and gene expression during early developmental stages of marine fish larvae⁽⁶¹⁾. The results from the present study demonstrate that there is an increase in the expression of specific antioxidant genes in sea bass larvae exposed to oxidative stress in order to neutralise the generated ROS. Moreover, when sea bass larvae were exposed to high dietary DHA contents (5%), the induction of antioxidant enzyme genes coincided with increases in MDA levels. Accordingly, studies in Manchurian trout (*Brachymystax lenok*) larvae⁽⁶²⁾ revealed that high dietary lipid levels produced elevated MDA levels, inducing an antioxidant response noticeable by an increase in the activity of AOE.




Fig. 1. Semithin micrographs of (a) longitudinal and (b) transversal sections (400 \times) from larvae fed the diet 5/300 (5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight) showing (a) coagulation of muscular proteins in the affected fibre (\rightarrow) and hypercontraction of the surrounding muscular fibres (*). (b) Mild affected fibres showed loss of the polyhedral structure, abundant vacuoles (*) and dilatation of sarcoplasmic membranes (\rightarrow).




Fig. 2. Electromicrographs of transversal sections of sea bass larvae fed the diet 5/300 (5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight). (a) Damaged muscle fibre showing autophagic (AV) and hydropic vacuoles (HV) and swollen mitochondria (arrow; 8000 ×). (b) Unaffected fibre where normal mitochondria can be observed (→; 8000 ×). (c) Presence of a satellite cell (SC) with a mitochondrion (*) between two damaged muscle fibres, with the presence of vacuoles and degenerated mitochondria (→; 5000 ×). MF, normal myofilaments; disMF, disarrayed myofilaments.

In the present study, an initial increase in the expression of each AOE was observed in all treatments at 26 dph, including the control group. Fernández-Díaz *et al.*⁽⁶³⁾ found that the administration of inert diets to *Solea senegalensis* larvae produced increased CAT and SOD activity compared with larvae fed with *Artemia*. Therefore, the observed initial increase in expression could be due to the use of inert food.

In an attempt to dismutate superoxide anions and to decompose H₂O₂, increases in SOD, CAT and GPX expression were detected in fish larvae fed a high-DHA and Se-free diet. Similarly, exposure to high-DHA diets caused a significant increase in CAT and GPX in the larvae fed the Se-supplemented diets. Given that increases in SOD activity were less significant in Se-supplemented larvae, it can be concluded that H₂O₂ formation declined or that CAT activity was sufficient to remove

H₂O₂. These results agree with Monteiro *et al.*⁽⁶⁴⁾ who observed that Se supplementation had a protective effect against oxidative stress caused by methyl parathion in *Brycon cephalus*, as denoted by a decrease in CAT and SOD activity. In contrast with these authors, Se supplementation did not increase the GPX level, or decreased GPX caused by methyl parathion. In mammals, it is likely that maintaining the activity of known selenoproteins, including GPX, is not the mechanism by which Se acts since it appears to be saturated at normal nutritional intakes. Thus, supranutritional levels of Se are required to reduce the incidence of human and animal diseases⁽⁶⁵⁾. Therefore, it appears that other selenoproteins could be implicated in tissue antioxidant defence mechanisms. Among all selenoproteins, selenoprotein P seems to play an important role as an antioxidant defence in mammals⁽⁶⁶⁾ by associating with




Fig. 3. (a) Catalase (CAT), (b) superoxide dismutase (SOD), (c) glutathione peroxidase (GPX), (d) insulin-like growth factor I (IGF-I), (e) insulin-like growth factor II (IGF-II) and (f) myosin heavy chain (MyHC) expression levels measured by real-time PCR in *Dicentrarchus labrax* larvae when fed the diets 1/150 (◆; 1 g DHA/100 g dry weight and 150 mg vitamin E/100 g dry weight), 5/300 (■; 5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight) or 5/300 + Se (●; 5 g DHA/100 g dry weight and 300 mg vitamin E/100 g dry weight supplemented with Se). mRNA copy number of each gene was normalised as a ratio to 100 ng total RNA. Values are means, with standard deviations represented by vertical bars. ^{a,b}Mean values with unlike letters were significantly different in gene expression among the treatments at given sampling points.

Table 6. Effects of the dietary treatment, time and their interaction on the global gene expression

	D	T	D × T
<i>CAT</i>	**	**	NS
<i>SOD</i>	**	**	*
<i>GPX</i>	**	**	NS
<i>IGF-I</i>	NS	NS	*
<i>IGF-II</i>	**	*	NS
<i>MyHC</i>	**	**	NS

D, diet; T, time; *CAT*, catalase; *SOD*, superoxide dismutase; *GPX*, glutathione peroxidase; *IGF-I*, insulin-like growth factor I; *IGF-II*, insulin-like growth factor II; *MyHC*, myosin heavy chain.

* $P \leq 0.05$, ** $P \leq 0.01$.

endothelial cells. Previous studies in zebrafish (*Danio rerio*) indicate that selenoprotein P is utilised to a larger extent than in human subjects, as it is encoded by two genes and has seventeen selenocysteine residues, the largest number of selenocysteine residues found in any known protein^(67,68). In consequence, the action of selenoprotein P or any other selenoproteins could be critical as an antioxidant defence against lipid H_2O_2 in sea bass larvae when Se requirements are covered. Thus, further studies are required to clarify the antioxidant mode of Se action in marine fish larvae.

As observed in our previous studies^(12,24), high dietary DHA levels caused pathological changes in sea bass larvae muscle. However, in the present study, inclusion of Se proved to be efficient in controlling the damage caused by ROS, reducing the incidence of muscle injury to almost half as compared with the 5/300 diet. Moreover, certain properties of muscle may render it especially susceptible to ROS injury⁽¹⁾. For instance, muscle is prone to oxidative injury as a result of increased electron flux due to its requirement and ability to undertake rapid and coordinated changes in energy supply and oxygen flux during contraction⁽⁶⁹⁾. There is also a very high concentration of myoglobin in the muscle, and it is known that such a haem-containing protein may confer greater sensitivity to free radical-induced damage by the conversion of H_2O_2 to a more reactive species⁽⁷⁰⁾. Furthermore, the requirement of skeletal muscle membranes for phospholipids containing large proportions of PUFA may render those membranes particularly susceptible to oxidative stress⁽⁷¹⁾. Finally, low Se accumulation in muscular tissues will make this tissue more susceptible to oxidative damage^(64,72). However, interaction of LC-PUFA with other cellular components should be taken into account to complete the scenario. In this sense, studies using juvenile salmon fed with high-EPA and DHA diets showed loss of mitochondrial β -oxidation, reduced lipid deposition and apoptosis in white adipose tissue, indicating that high supplementation rates of these LC-PUFA may lead to oxidative stress⁽⁷³⁾.

IGF-I and IGF-II are polypeptides well known for promoting proliferation and differentiation in many vertebrates with nutritional status having a profound effect on the IGF system in fish⁽⁷⁴⁾. However, most nutritional studies have focused on the effects of food restriction^(32,33,75), dietary protein or carbohydrate content^(76,77) and probiotics⁽⁷⁸⁾, whereas little information is known about the effect of lipids on this system.

Moreover, no information exists about the effect of oxidative stress on IGF in fish larvae. In the present study, an increase in IGF-I and IGF-II in the larvae fed the 5/300 diet was observed, especially when no Se was added, suggesting oxidative stress may play a role in the expression of these growth factors. Accordingly, the larvae fed the highest content of DHA showed a higher incidence of muscular lesions and the presence of abundant satellite cells. Satellite cells are able to regenerate damaged muscle by forming new myofibres by fusing to existing muscle fibres or fusing together⁽⁷⁹⁾. It is known that to control the satellite cell population, growth factors are required⁽⁸⁰⁾. In mammals, IGF-I appears to utilise multiple signalling pathways in the regulation of the satellite cell pool such as the mitogen-activated protein or phosphatidylinositol-3-OH kinase^(81,82). In agreement with this, Pozios *et al.*⁽⁸³⁾ showed that IGF-II and IGF-I potently activate cell proliferation and DNA synthesis in embryonic zebrafish cells via mitogen-activated protein and phosphatidylinositol-3-OH kinase, suggesting that the increase in IGF-I mRNA copies observed in the present study in the larvae with the highest incidence of muscular lesions could be due to the regeneration process carried out by satellite cells. Similarly, the mitogenic effect of IGF in fish has also been described in cultured muscle cells from rainbow trout⁽⁸⁴⁾.

Late markers of myogenesis include the myofibrillar protein MyHC. By monitoring the expression patterns of this marker gene, the effect that nutritional status has on muscle growth can be determined⁽²⁹⁾. Regeneration of fish muscle has only rarely been described. Rowleron *et al.*⁽⁸⁵⁾ demonstrated a vigorous regeneration in juvenile sea bream (*Sparus aurata*) after mechanical injury, with myosin expression in regenerating fibres resembling that seen in newly produced fibres in post-larval white muscle. In the present study, a higher expression of myosin was observed in the 5/300 larvae, especially when Se was not added to the diets with a positive correlation observed between the incidence of muscular lesions and myosin mRNA copies at 35 dph ($y = 2 \times 10^{-6}x - 29.567$; $R^2 = 0.9133$).

In the present study, the high levels of AOE and MDA content observed in the 5/300 larvae demonstrate an adaptive response in attempting to neutralise the generated ROS. Moreover, a reactive response was observed by the increase in IGF and MyHC expression in larval tissues, suggesting regenerative processes in the injured muscle. Organic Se proved to enhance the cell antioxidant capacity, protecting muscle, as shown by the decrease in the incidence of muscular lesions, MDA content and AOE expression. Therefore, when high levels of LC-PUFA are included in sea bass larvae microdiets, an adequate combination of dietary α -tocopherol and Se must be included to avoid the appearance of oxidative stress in larval tissues and favour culture performance.

Acknowledgements

M. B. B. was supported by a grant given by the Cabildo Insular de Gran Canaria. The present study was funded by the Spanish Ministry of Sciences and Education (AGL2009-14661). The authors are thankful to Dr Matthew Sprague for his assistance in proof reading the manuscript. Acknowledgments are due to

Instituto de Acuicultura de Torre la Sal that kindly provided the sea bass eggs, Croda Company and Alltech that provided DHA and EPA 50 and the Se yeast, respectively. The authors disclose no conflicts of interest. M. B. B. carried out the experiment, performed all analyses, participated in the interpretation of the results and drafted the manuscript. M. J. C. and M. I. participated in the animal experimental design, interpretation of the results and the preparation of the manuscript. R. S. and T. B.-S. participated in the animal feeding and sampling. E. A. collaborated to formulate and prepare the experimental microdiets. G. T. supervised the molecular biology analysis and J. G. B. supervised the vitamin E and Se analysis.

References

- Rando TA (2002) Oxidative stress and the pathogenesis of muscular dystrophies. *Am J Phys Med Rehab* **81**, 175–186.
- Halliwell B & Gutteridge JMC (1996) Lipid peroxidation: a radical chain reaction. In *Free Radicals in Biology and Medicine*, pp. 188–266 [B Halliwell and JMC Gutteridge, editors]. Oxford: Clarendon Press.
- Bell JG, Cowey CB, Adron JW, *et al.* (1987) Some effects of selenium deficiency on enzyme activities and indices of tissue peroxidation in Atlantic salmon parr (*Salmo salar*). *Aquaculture* **65**, 43–54.
- Wilhelm-Filho D (1996) Fish antioxidant defenses – a comparative approach. *Braz J Med Biol Res* **29**, 1735–1742.
- Lygren B, Hamre K, Waagbø R, *et al.* (1999) Effect of dietary pro- and antioxidants on some protective mechanisms and health parameters in Atlantic salmon. *J Aquat Anim Health* **11**, 211–221.
- Bell JG, McEvoy J, Tocher DR, *et al.* (2000) Depletion of α -tocopherol and astaxanthin in Atlantic salmon (*Salmo salar*) affects autoxidative defence and fatty acid metabolism. *J Nutr* **130**, 1800–1808.
- Díaz ME, Furné M, Trenzado CE, *et al.* (2010) Antioxidant defences in the first life phases of the sturgeon *Acipenser naccarii*. *Aquaculture* **307**, 123–129.
- Hamre K (2010) Metabolism, interactions, requirements and functions of vitamin E in fish. *Aquacult Nutr* **17**, 98–115.
- Mourente G, Tocher DR, Díaz E, *et al.* (1999) Relationships between antioxidants, antioxidant enzyme activities and lipid peroxidation products during early development in *Dentex dentex* eggs and larvae. *Aquaculture* **179**, 309–324.
- Atalah E (2008) Importance of the proportions of dietary polyunsaturated fatty acids and antioxidants in larval development of marine fish. PhD Thesis, University of Las Palmas de Gran Canaria.
- Penglase S, Nordgreen A, Van der Meeren T, *et al.* (2010) Increasing the level of selenium in rotifers (*Brachionus plicatilis* ‘Cayman’) enhances the mRNA expression and activity of glutathione peroxidase in cod (*Gadus morhua* L.) larvae. *Aquaculture* **306**, 259–269.
- Betancor MB, Atalah E, Caballero MJ, *et al.* (2011) α -Tocopherol in weaning diets for European sea bass (*Dicentrarchus labrax*) improves survival and reduces tissue damage caused by excess dietary DHA contents. *Aquacult Nutr* **13**, e112–e122.
- Lall SP & Bishop FJ (1977) Studies on mineral and protein utilization by Atlantic salmon (*Salmo salar*) grown in sea water. Fisheries and Marine Service, Environment Canada, Ottawa, ON, Technical Report No. 688, pp. 16.
- Halver JE (2002) The vitamins. In *Fish Nutrition*, 2nd ed., pp. 62–132 [JE Halver and R Hardy, editors]. New York: Academic Press.
- Felton SP, Landolt ML & Grace R (1996) Effects of selenium dietary enhancement on hatchery-reared coho salmon, *Oncorhynchus kisutch* (Walbaum), when compared with wild coho: hepatic enzymes and seawater adaptation evaluated. *Aquacult Res* **27**, 135–142.
- Arteel GE & Sies H (2001) The biochemistry of selenium and the glutathione system. *Environ Toxicol Phar* **10**, 153–158.
- Poston HA, Combs GF Jr & Leibovitz L (1976) Vitamin E and selenium interrelations in the diet of Atlantic salmon (*Salmo salar*): gross, histological and biochemical deficiency signs. *J Nutr* **106**, 892–904.
- Bell JG & Cowey CB (1985) Roles of vitamin E and selenium in the prevention of pathologies related to fatty acid oxidation in salmonids. In *Nutrition and Feeding in Fish*, pp. 333–347 [CB Cowey, AM Mackie and JG Bell, editors]. London: Academic Press.
- Bell JG, Pirie BJS, Adron JW, *et al.* (1986) Some effects of selenium deficiency on glutathione peroxidase (EC 1.11.1.9) activity and tissue pathology in rainbow trout (*Salmo gairdneri*). *Br J Nutr* **55**, 305–311.
- Gatlin DM, Poe WE & Wilson RF (1986) Effects of singular and combined dietary deficiencies of selenium and vitamin E on fingerling channel catfish (*Ictalurus punctatus*). *J Nutr* **116**, 1061–1067.
- Lin YH & Shiau SY (2009) Mutual sparing of dietary requirements for alpha-tocopherol and selenium in grouper, *Epinephelus malabaricus*. *Aquaculture* **294**, 242–245.
- McEvoy LA, Naess T, Bell JG, *et al.* (1998) Lipid and fatty acid composition of normal and malpigmented Atlantic halibut (*Hippoglossus hippoglossus*) fed enriched *Artemia*: a comparison with fry fed wild copepods. *Aquaculture* **163**, 237–250.
- Evjemo JO, Reitan KL & Olsen Y (2003) Copepods and live food organisms in the larval rearing of halibut larvae (*Hippoglossus hippoglossus* L.) with special emphasis on the nutritional value. *Aquaculture* **227**, 191–210.
- Betancor MB, Caballero MJ, Benítez-Santana T, *et al.* (2012) Oxidative status and histological changes in sea bass larvae muscle in response to high dietary content of DHA. *J Fish Dis* (In the press).
- Johnston IA, Abercromby M, Vieira VL, *et al.* (2004) Rapid evolution of muscle fibre number in post-glacial populations of Arctic charr *Salvelinus alpinus*. *J Exp Biol* **207**, 4343–4360.
- Allen RE & Boxhorn LK (1989) Regulation of skeletal muscle satellite cell proliferation and differentiation by transforming growth-factors beta, insulin-like growth factor I and fibroblast growth factor. *J Cell Physiol* **138**, 311–315.
- Johnston IA (1999) Muscle development and growth: potential implications for flesh quality in fish. *Aquaculture* **177**, 99–115.
- Goldspink G, Wilkes D & Ennion S (2001) Myosin expression during ontogeny, post-hatching growth and adaptation. In *Muscle Development and Growth*, pp. 43–66 [IA Johnston, editor]. San Diego, CA: Academic Press.
- Bower NI, Li X, Taylor R, *et al.* (2008) Switching to fast growth: the insulin-like growth factor (IGF) system in skeletal muscle of Atlantic salmon. *J Exp Biol* **211**, 3859–3870.
- Duan C & Plisetskaya EM (1993) Nutritional regulation of insulin-like growth factor-I mRNA expression in salmon tissues. *J Endocrinol* **139**, 243–252.
- Matthews SJ, Kinhult AK, Hoeben P, *et al.* (1997) Nutritional regulation of insulin-like growth factor I mRNA expression in barramundi, *Lates calcarifer*. *J Mol Endocrinol* **18**, 273–276.
- Terova G, Rimoldi S, Chini V, *et al.* (2007) Cloning and expression analysis of insulin-like growth factor I and II in

- liver and muscle of sea bass (*Dicentrarchus labrax*, L.) during long-term fasting and refeeding. *J Fish Biol* **70**, 219–233.
33. Hevrøy EM, Azpeleta C, Shimizu M, *et al.* (2011) Effects of short-term starvation on ghrelin, GH–IGF system and IGF-binding proteins in Atlantic salmon. *Fish Physiol Biochem* **37**, 217–232.
 34. Chauvigné F, Gabillard JC, Weil C, *et al.* (2003) Effect of refeeding on IGFI, IGFII, IGF receptors, FGF2, FGF6, and myostatin mRNA expression in rainbow trout myotomal muscle. *Gen Comp Endocrinol* **132**, 209–215.
 35. Watabe S (1999) Myogenic regulatory factors and muscle differentiation during ontogeny in fish. *J Fish Biol* **55**, 1–18.
 36. Silva P, Rowlerson AM, Valente LMP, *et al.* (2008) Muscle differentiation in blackspot seabream (*Pagellus bogaraveo*, Brunnich): histochemical and immunohistochemical study of the fibre types. *Tissue Cell* **40**, 447–458.
 37. Liu J, Caballero MJ, Izquierdo MS, *et al.* (2002) Necessity of dietary lecithin and eicosapentaenoic acid for growth, survival, stress resistance and lipoprotein formation in gilthead sea bream *Sparus aurata*. *Fish Sci* **68**, 1165–1172.
 38. AOAC (1995) *Official Methods of Analysis*, 16th ed. Washington, DC: AOAC International.
 39. Folch J, Lees M & Sloane Stanley GH (1957) A simple method for the isolation and purification of total lipides from animal tissues. *J Biol Chem* **226**, 497–509.
 40. Christie WW (1982) *Lipid Analysis*. Oxford: Pergamon Press.
 41. Izquierdo MS, Arakawa T, Takeuchi T, *et al.* (1992) Effect of *n*-3 HUFA levels in *Artemia* on growth of larval Japanese flounder (*Paralichthys olivaceus*). *Aquaculture* **105**, 73–82.
 42. McMurray CH, Blanchflower WJ & Rice DA (1980) Influence of extraction techniques on determination of α -tocopherol in animal feedstuffs. *J Assoc Off Anal Chem* **63**, 1258–1261.
 43. Cowey CB, Adron JW, Walton MJ, *et al.* (1981) Tissue distribution, uptake and requirement for α -tocopherol of rainbow trout (*Salmo gairdneri*) fed diets with a minimal content of unsaturated fatty acids. *J Nutr* **111**, 1556–1567.
 44. Burk RF, Trumble MJ & Lawrence RA (1980) Rat hepatic cytosolic GSH-dependent enzyme protection against lipid peroxidation in the NADPH microsomal lipid peroxidation system. *Biochim Biophys Acta* **618**, 35–41.
 45. Martoja R & Martoja Pearson M (1970) *Técnicas de Histología Animal (Animal Histology Techniques)*. Barcelona: Toray-Masson, SA.
 46. Hoffman EO, Flores TR, Coover J, *et al.* (1983) Polychrome stains for high resolution light microscopy. *Lab Med* **14**, 779–781.
 47. Combs GF Jr & Scott ML (1974) Dietary requirements for vitamin E and selenium measured at the cellular level in the chick. *J Nutr* **104**, 1292–1296.
 48. Hamre K, Srivastava A, Rønnestad I, *et al.* (2008) Several micronutrients in the rotifer *Brachionus* sp. may not fulfil the nutritional requirements of marine fish larvae. *Aquac Nutr* **14**, 51–60.
 49. Van der Meer T, Olsen RE, Hamre K, *et al.* (2008) Biochemical composition of copepods for evaluation of feed quality in production of juvenile marine fish. *Aquaculture* **274**, 375–397.
 50. National Research Council (NRC) (1993) *Nutrient Requirements of Fish*. Washington, DC: National Academy Press pp. 62–63.
 51. Hilton JW, Hodson PV & Slinger SJ (1980) The requirement and toxicity of selenium in rainbow trout (*Salmo gairdneri*). *J Nutr* **110**, 2527–2535.
 52. Lin YH & Shiau SY (2005) Dietary selenium requirement of grouper, *Epinephelus malabaricus*. *Aquaculture* **250**, 356–363.
 53. Jaramillo F, Peng L & Gatlin D (2009) Selenium nutrition of hybrid striped bass (*Morone chrysops* \times *M. saxatilis*) bioavailability, toxicity and interaction with vitamin E. *Aqua Nutr* **15**, 160–165.
 54. Tinggi U (2003) Essentiality and toxicity of selenium and its status in Australia: a review. *Toxicol Lett* **137**, 103–110.
 55. Rider SA, Davies SJ, Jha AN, *et al.* (2010) Bioavailability of co-supplemented organic and inorganic zinc and selenium sources in white fishmeal-based rainbow trout (*Oncorhynchus mykiss*) diet. *J Anim Physiol Anim Nutr* **94**, 99–110.
 56. Kiron V, Puangkaew J, Ishizaka K, *et al.* (2004) Antioxidant status and nonspecific immune response in rainbow trout (*Oncorhynchus mykiss*) fed two levels of vitamin E along with three lipid sources. *Aquaculture* **234**, 361–379.
 57. Puangkaew J, Kiron V, Satoh S, *et al.* (2005) Antioxidant defense of rainbow trout (*Oncorhynchus mykiss*) in relation to dietary *n*-3 highly unsaturated fatty acids and vitamin E contents. *Comp Biochem Physiol C* **140**, 187–196.
 58. Ji H, Li J & Liu P (2011) Regulation of growth performance and lipid metabolism by dietary *n*-3 highly unsaturated fatty acids in juvenile grass carp, *Ctenopharyngodon idellus*. *Comp Biochem Physiol B* **159**, 49–56.
 59. Kim S, Ji K, Lee S, *et al.* (2011) Perfluorooctane sulfonic acid (PFOS) exposure increases cadmium toxicity in early life stage of zebrafish (*Danio rerio*). *Environ Toxicol Chem* **38**, 870–877.
 60. Peters LD & Livingstone DR (1996) Antioxidant enzyme activities in embryologic and early larval stages of turbot. *J Fish Biol* **49**, 986–997.
 61. Tovar-Ramírez D, Mazurais D, Gatesoupe JF, *et al.* (2010) Dietary probiotics live yeast modulate antioxidant enzyme activities and gene expression of sea bass (*Dicentrarchus labrax*) larvae. *Aquaculture* **300**, 142–147.
 62. Zhang H, Mu Z, X LM, *et al.* (2009) Dietary lipid level induced antioxidant response in Manchurian trout, *Brachymystax lenok* (Pallas) larvae. *Lipids* **44**, 643–654.
 63. Fernández-Díaz C, Kopecka I, Canavete JP, *et al.* (2006) Variations on development and stress resistance in *Solea senegalensis* larvae fed on live and microencapsulated diets. *Aquaculture* **251**, 573–584.
 64. Monteiro DA, Rantin FT & Kalinin AL (2009) The effects of selenium on oxidative stress biomarkers in the freshwater characid fish matrinxã, *Brycon cephalus* (Günther, 1869) exposed to organophosphate insecticide Folisuper 600 BR[®] (methyl parathion). *Comp Biochem Physiol C* **149**, 40–49.
 65. Brown KM & Arthur JR (2001) Selenium, selenoproteins and human health: a review. *Public Health Nutr* **4**, 593–599.
 66. Burk RF, Hill KE, Boeglin ME, *et al.* (1997) Selenoprotein P associates with endothelial cells in rat tissues. *Histochem Cell Biol* **108**, 11–15.
 67. Tubajeva RM, Ransom DG, Harney JW, *et al.* (2000) Expression and characterization of nonmammalian selenoprotein P in the zebrafish, *Danio rerio*. *Genes Cells* **5**, 897–903.
 68. Kryukov GV & Gladyshev VN (2000) Selenium metabolism in zebrafish: multiplicity of selenoprotein genes and expression of a protein containing 17 selenocysteine residues. *Genes Cells* **5**, 1049–1060.
 69. Haycock JW, Jones P, Harris JB, *et al.* (1996) Differential susceptibility of human skeletal muscle proteins to free radical induced oxidative damage: a histochemical, immunocytochemical and electron microscopical study *in vitro*. *Acta Neuropathol* **92**, 331–340.

70. Ostdal H, Skibsted LH & Andersen HJ (1997) Formation of long-lived protein radicals in the reaction between H₂O₂-activated metamyoglobin and others proteins. *Free Rad Biol Med* **23**, 754–761.
71. Murphy ME & Kehrer JP (1989) Oxidative stress and muscular dystrophy. *Chem Biol Interact* **69**, 101–178.
72. Elia AC, Prearo M, Pacini N, *et al.* (2011) Effects of selenium on growth, accumulation and antioxidant response in juvenile carp. *Ecotoxicol Environ Saf* **74**, 166–173.
73. Todorčević M, Kjær M, Djaković N, *et al.* (2009) *n*-3 HUFAs affect fat deposition, susceptibility to oxidative stress, and apoptosis in Atlantic salmon visceral adipose tissue. *Comp Biochem Physiol B* **152**, 135–143.
74. Duan C (1998) Nutritional and developmental regulation of insulin-like growth factors in fish. *J Nutr* **128**, 306S–314S.
75. Moriyama S, Swanson P, Nishi M, *et al.* (1994) Development of a homologous radioimmunoassay for coho salmon insulin-like growth factor-I. *Gen Comp Endocrinol* **96**, 149–161.
76. Enes P, Sánchez-Gurmaches J, Navarro I, *et al.* (2010) Role of insulin and IGF-I on the regulation of glucose metabolism in European sea bass (*Dicentrarchus labrax*) fed with different dietary carbohydrate levels. *Comp Biochem Physiol A* **157**, 346–353.
77. Pérez-Sánchez J, Martipalanca H & Kaushik SJ (1995) Ration size and protein intake affect circulating growth hormone concentration, hepatic growth binding and plasma insulin-like growth factor I immunoreactivity in a marine teleost, the gilthead sea bream (*Sparus aurata*). *J Nutr* **125**, 546–552.
78. Carnevali O, de Vivo L, Sulpizio R, *et al.* (2006) Growth improvement by probiotic in European sea bass juveniles (*Dicentrarchus labrax* L.), with particular attention to IGF-1, myostatin and cortisol gene expression. *Aquaculture* **258**, 430–438.
79. Schultz E & McCormick KM (1994) Skeletal muscle satellite cells. *Rev Physiol Biochem Pharmacol* **123**, 213–257.
80. Seale P & Rudnicki MA (2000) A new look at the origin, function and ‘stem-cell’ status of muscle satellite cells. *Dev Biol* **218**, 115–124.
81. Coolican SA, Samuel DS, Ewton DZ, *et al.* (1997) The mitogenic and myogenic actions of insulin-like growth factors utilize distinct signalling pathways. *J Biol Chem* **272**, 6653–6662.
82. Semsarian C, Wu MJ, Ju YK, *et al.* (1999) Skeletal muscle hypertrophy is mediated by a Ca²⁺-dependent calcineurin signaling pathway. *Nature* **400**, 576–581.
83. Pozios KC, Ding J, Degger B, *et al.* (2001) IGFs stimulate zebrafish cell proliferation by activating MAP kinase and PI3-kinase-signalling pathways. *Am J Physiol* **280**, R1230–R1239.
84. Castillo J, Codina M, Martínez ML, *et al.* (2004) Metabolic and mitogenic effects of IGF-I and insulin on muscle cells of rainbow trout. *Am J Physiol Integr Comp Physiol* **286**, 935–941.
85. Rowlerson A, Radaelli G, Mascarello F, *et al.* (1997) Regeneration of skeletal muscle in two teleost fish: *Sparus aurata* and *Brachydanio rerio*. *Cell Tissue Res* **289**, 311–322.