

Systematic Review

Metabolic benefits of dietary prebiotics in human subjects: a systematic review of randomised controlled trials

Nicole J. Kellow^{1*}, Melinda T. Coughlan^{2,3} and Christopher M. Reid¹

¹*Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive Medicine, Monash University, Alfred Medical Research and Education Precinct, Melbourne, Victoria 3004, Australia*

²*Glycation, Nutrition and Metabolism Laboratory, Baker IDI Heart and Diabetes Institute, Melbourne, Victoria 8008, Australia*

³*Department of Medicine, Central Clinical School, Monash University, Alfred Medical Research and Education Precinct, Melbourne, Victoria 3004, Australia*

(Submitted 30 July 2013 – Final revision received 7 October 2013 – Accepted 8 October 2013 – First published online 13 November 2013)

Abstract

Complex relationships exist between the gut microflora and their human hosts. Emerging evidence suggests that bacterial dysbiosis within the colon may be involved in the pathogenesis of the metabolic syndrome, type 2 diabetes and CVD. The use of dietary prebiotic supplements to restore an optimal balance of intestinal flora may positively affect host metabolism, representing a potential treatment strategy for individuals with cardiometabolic disorders. The present review aimed to examine the current evidence supporting that dietary prebiotic supplementation in adults has beneficial effects on biochemical parameters associated with the development of metabolic abnormalities including obesity, glucose intolerance, dyslipidaemia, hepatic steatosis and low-grade chronic inflammation. Between January 2000 and September 2013, eight computer databases were searched for randomised controlled trials published in English. Human trials were included if at least one group received a dietary prebiotic intervention. In the present review, twenty-six randomised controlled trials involving 831 participants were included. Evidence indicated that dietary prebiotic supplementation increased self-reported feelings of satiety in healthy adults (standardised mean difference -0.57 , 95% CI -1.13 , -0.01). Prebiotic supplementation also significantly reduced postprandial glucose (-0.76 , 95% CI -1.41 , -0.12) and insulin (-0.77 , 95% CI -1.50 , -0.04) concentrations. The effects of dietary prebiotics on total energy intake, body weight, peptide YY and glucagon-like peptide-1 concentrations, gastric emptying times, insulin sensitivity, lipids, inflammatory markers and immune function were contradictory. Dietary prebiotic consumption was found to be associated with subjective improvements in satiety and reductions in postprandial glucose and insulin concentrations. Additional evidence is required before recommending prebiotic supplements to individuals with metabolic abnormalities. Large-scale trials of longer duration evaluating gut microbial growth and activity are required.

Key words: Prebiotics: Gut microflora: Human subjects: Metabolic disorders

The composition and possible health effects of human gut micro-organisms have been the focus of renewed interest since the development of metagenomic techniques enabling the identification and characterisation of intestinal bacteria that cannot be cultured. In addition, the discovery of differences in gut microbial composition between lean and obese individuals⁽¹⁾ and people with and without type 2 diabetes^(2,3) has highlighted the potential role played by the

colonic microflora and their fermentation products in the pathogenesis of host metabolic health and disease.

Although the number and diversity of bacterial species within an individual's gastrointestinal tract remain relatively constant throughout life, it is possible to stimulate the proliferation of specific micro-organisms known to have beneficial health effects by manipulating the host diet. Prebiotics are defined as non-digestible plant-derived carbohydrates that

Abbreviations: GLP, glucagon-like peptide; GPR, G protein-coupled receptor; HOMA-IR, homeostasis model assessment for insulin resistance; LPS, lipopolysaccharide; NASH, non-alcoholic steatohepatitis; SMD, standardised mean difference.

* **Corresponding author:** N. J. Kellow, fax +61 3 990 30556, email nicolekellow@live.com.au

act as a fermentation substrate within the colon, stimulating the preferential growth and activity of a limited number of microbial species that confer health benefits on the host⁽⁴⁾. Carbohydrates with an established prebiotic effect include inulin-type fructans (inulin, oligofructose and fructo-oligosaccharides) and galactans (galacto-oligosaccharides)⁽⁵⁾, known to promote the proliferation of beneficial lactic acid-producing species such as bifidobacteria and lactobacilli⁽⁶⁾.

Gut bacteria play an important role in the development of the host immune system⁽⁷⁾ and modulation of inflammatory processes⁽⁸⁾, extraction of energy from the host diet⁽⁹⁾, fermentation of dietary fibres to produce SCFA⁽¹⁰⁾, alteration of human glucose and fatty acid metabolism⁽¹¹⁾, regulation of intestinal permeability⁽¹²⁾, production of vitamins⁽¹³⁾ and promotion of mineral absorption by the host⁽¹⁴⁾. They may also be involved in the modification of the secretion of gut hormones to enhance satiety and improve gastrointestinal function⁽¹⁵⁾. Dietary prebiotic supplements capable of favourably altering the composition of the intestinal microflora might represent a potential therapeutic strategy for the prevention and treatment of metabolic abnormalities widespread in modern society.

The present review aimed to examine the current evidence supporting dietary prebiotic supplementation in adults on biochemical parameters associated with the development of metabolic abnormalities such as obesity, glucose intolerance, dyslipidaemia, non-alcoholic fatty liver disease and low-grade chronic inflammation.

Methods

A computer search of databases such as MEDLINE, CINAHL, Embase, Current Contents, PubMed, Cochrane Central Register of Controlled Trials, Cochrane Database of Systematic Reviews and AMED was undertaken for the period between 1 January 2000 and 30 September 2013. Databases were not searched before 2000, to exclude studies utilising non-molecular, culture-dependent techniques for the characterisation of intestinal bacteria. Reference lists of all identified studies were hand-searched for relevant trials. The following search terms were used: (1) (prebiotic* OR fructan* OR oligofructose OR inulin OR fructooligosaccharide* OR galactooligosaccharide*) and (gut OR obes* OR diabet* OR lipid*

OR hepat* OR immune* OR metaboli*); (2) limit 1 to year = '2000–2003'; (3) limit 2 to humans. Trials were included if they were published in English and involved human participants aged ≥18 years and at least one group of participants were randomised to receive a dietary prebiotic intervention. For the purposes of the present review, a prebiotic intervention was defined as one that contained inulin, oligofructose, fructo-oligosaccharides or galacto-oligosaccharides. Additional plant-derived carbohydrates such as arabinoxylan and β-glucan were excluded from the search, as, although demonstrated to have prebiotic effects^(16,17), these compounds require further research before being formally classified as prebiotics.

Dietary prebiotic intervention studies of less than 24 h duration were excluded from the present review, as the growth of colonic microflora is unlikely to be affected in this brief time period⁽¹⁸⁾. Nutritional intervention studies involving the administration of probiotics (beneficial live micro-organisms) or synbiotics (a combination of pre- and probiotics) were also excluded. Trials involving prebiotic supplementation in people with disease conditions such as HIV and inflammatory bowel disease were considered to be outside the scope of the present review and were therefore excluded. The methodological quality of all the included trials was assessed by two authors independently using the Heyland Methodological Quality Score⁽¹⁹⁾ (Table 1). This checklist rates primary research based on the use of allocation concealment during randomisation, intention-to-treat analysis, double-blinding, patient selection with minimal risk of bias, comparability of intervention and control groups at baseline, 100% participant follow-up, clearly described treatment protocol and well-defined outcome measurements. Trials scoring ≥8 out of a possible 14 points were considered to be of high methodological quality. Disagreements between authors in assigning methodological quality scores were resolved by discussion until consensus was reached.

Trials measuring similar outcomes were subjected to a random-effects model meta-analysis using RevMan 5.1 (The Cochrane Collaboration, Copenhagen 2011). Treatment effects and 95% CI were calculated using the Hedges (adjusted-g) standardised mean difference (SMD), to enable the comparison of effect sizes between trials using different

Table 1. Heyland Methodological Quality Score⁽¹⁹⁾

Criterion	Score		
	0	1	2
Randomisation	Not applicable	Not concealed or not sure	Concealed randomisation
Analysis	Other	Not applicable	Intention to treat
Blinding	Not blinded	Single blind	Double blind
Patient selection	Selected patients or unable to tell	Consecutive eligible patients	Not applicable
Comparability of groups at baseline	No or not sure	Yes	Not applicable
Extent of follow-up	Less than 100 %	100 %	Not applicable
Treatment protocol	Poorly described	Reproducibly described	Not applicable
Co-interventions applied equally across groups	Not described	Described but not equal or not sure	Well described and all equal
Outcomes	Not described	Partially described	Objectively defined

The Heyland Methodological Quality Score for individual studies is based on nine quality criteria. The maximum possible score is 14 with studies scoring ≥8 considered to be of high methodological quality.

outcome measures. SMD values of 0.2, 0.5 and 0.8 were considered to represent small, moderate and large effect sizes, respectively⁽²⁰⁾. Limited numbers of studies investigating comparable outcomes, small sample sizes and heterogeneity among trial subjects, disease conditions, prebiotic supplements, intervention duration and outcome measures limited the majority of data synthesis to a narrative analysis.

Results

Description of the selected trials

A total of 1130 citations were originally identified at the time of the initial database search and were selected to be included in the review based on the predefined inclusion criteria (Fig. 1). In the present review, twenty-nine articles reporting on twenty-six randomised controlled trials involving 831 participants were ultimately included^(21–49). The characteristics of the included trials are outlined in Table 2. Of the twenty-six trials included in the present review, thirteen trials included only healthy participants, five trials included only overweight or obese participants, one trial included only overweight participants with the metabolic syndrome, two trials included only participants with type 2 diabetes, two trials included only participants with hypercholesterolaemia, one trial included only participants with non-alcoholic steatohepatitis (NASH), one trial included only participants with gastro-oesophageal reflux disease and one trial included only elderly participants diagnosed with mild malnutrition or at risk of becoming malnourished. The duration of intervention ranged from 2 d to 28 weeks and the participants were aged 19–99 years. A variety of post-intervention outcome measures were reported including self-reported hunger and satiety ratings, total body weight, BMI, waist circumference,

energy intake, gastric emptying times, concentrations of appetite-regulating hormones (ghrelin, cholecystokinin, peptide YY and glucagon-like peptide (GLP)-1), concentrations of lipids (total cholesterol, LDL, HDL, TAG, Lp(a) and NEFA), indicators of glucose homeostasis (glucose, insulin, glucagon, homeostasis model assessment for insulin resistance (HOMA-IR), HbA1c and fructosamine), inflammatory markers (TNF- α , C-reactive protein and IL), indices of immune function (natural killer cell activity and T-cell activation), and parameters associated with oxidative stress (total radical-trapping antioxidant parameter (TRAP), photosensitive chemiluminescence, total antioxidant capacity, superoxide dismutase and malondialdehyde) and liver function (aspartate aminotransferase). All the trials were of high methodological quality as assessed by the Heyland Methodological Quality Score. Methodological strengths of the trials included double-blinding utilised in the majority of the studies and random allocation of participants to intervention and control groups or treatment sequence. Methodological limitations of most of the trials included small sample sizes and short study duration. Some cross-over studies did not have a washout period or did not stipulate the duration of their washout period.

Outcomes associated with body weight

Of the five trials investigating the effect of dietary prebiotic supplementation on self-reported quantitative ratings of satiety, three demonstrated improvements in subjective satiety measurements in healthy participants consuming prebiotics in comparison with controls^(22,23,49). After the meta-analysis (n 52), the pooled SMD for satiety was -0.57 (95% CI -1.13 , -0.01 ; $P < 0.05$), indicating a statistically significant effect favouring prebiotic supplementation over placebo (Fig. 2). Inclusion of two trials finding no change in satiety




Fig. 1. Flow chart showing the progression of trials through each stage of the selection process. RCT, randomised controlled trial. (A colour version of this figure can be found online at <http://www.journals.cambridge.org/bjn>).

Table 2. Summary of published human intervention randomised controlled trials examining the relationship between dietary prebiotic intake and metabolic health

Study	Participants	Study design/blinding	Dietary prebiotic intervention	Effect of dietary prebiotic supplements on metabolic outcomes
Bunout <i>et al.</i> (2002) ⁽²¹⁾	Chile, <i>n</i> 43 normal-weight and overweight elderly adults: sex not stated (mean age 75.5 years; mean BMI 27 kg/m ²)	Parallel RCT double-blinded HMQS: 11	Random assignment to either a 28-week prebiotic-supplemented diet (6 g FOS/d) or a 28-week placebo-supplemented diet (6 g maltodextrin/d)	↔ IL-4 and interferon-gamma ↔ Secretory IgA
Cani <i>et al.</i> (2006) ⁽²²⁾	Belgium, <i>n</i> 10 healthy adults: five males and five females (mean age 27 years; mean BMI 22.3 kg/m ²)	Cross-over RCT subjects blinded HMQS: 9	Random assignment to either a 2-week prebiotic-supplemented diet (16 g oligofructose/d) or a 2-week placebo-supplemented diet (16 g maltodextrin/d) before cross-over. Washout period: 2 weeks	↑ Satiety after breakfast and dinner ↓ Hunger after dinner ↓ Energy intake after dinner ↓ Total energy intake
Cani <i>et al.</i> (2009) ⁽²³⁾	Belgium, <i>n</i> 10 healthy adults: five males and five females (mean age 26 years; mean BMI 21.6 kg/m ²)	Parallel RCT double-blinded HMQS: 10	Random assignment to either a 2-week prebiotic-supplemented diet (16 g chicory-derived fructan/d) or a 2-week placebo-supplemented diet (16 g maltodextrin/d)	↑ Plasma GLP-1 ↑ Plasma peptide YY ↓ Postprandial plasma glucose ↓ Hunger ↔ Satiety
Causey <i>et al.</i> (2000) ⁽²⁴⁾	USA, <i>n</i> 12 adult males with mild hypercholesterolaemia (age range 27–49 years)	Cross-over RCT double-blinded HMQS: 8	Random assignment to either a 3-week prebiotic-supplemented diet (20 g inulin/d in low-fat ice cream) or a 3-week placebo-supplemented diet (regular low-fat ice cream containing sucrose). Washout period: nil	↓ TAG ↑ Postprandial plasma glucagons
Daubioul <i>et al.</i> (2005) ⁽²⁵⁾	Belgium, <i>n</i> 7 adult males with NASH (mean age 55 years; mean BMI 29.1 kg/m ²)	Cross-over RCT double-blinded HMQS: 8	Random assignment to either an 8-week prebiotic-supplemented diet (16 g oligofructose/d) or an 8-week placebo-supplemented diet (16 g maltodextrin/d) before cross-over. Washout period: 5 weeks	↓ Serum AST ↔ Serum insulin ↔ Serum TAG
Dehghan <i>et al.</i> (2013) ⁽²⁶⁾ Pourghassem Gargari <i>et al.</i> (2013) ⁽²⁷⁾	Iran, <i>n</i> 49 women with type 2 diabetes (mean age 48.3 years; mean BMI 30.8 kg/m ² ; time after DM diagnosis > 6 months; mean HbA1c levels 8.3 %)	Parallel RCT double-blinded HMQS: 9	Random assignment to either an 8-week prebiotic-supplemented diet (10 g inulin/d) or an 8-week placebo-supplemented diet (10 g maltodextrin/d)	↓ Fasting glucose ↓ Energy intake and HbA1c ↓ CRP and TNF-α ↓ Lipopolysaccharide ↓ Malondialdehyde ↑ TAC and ↑ SOD activity ↔ Fasting insulin and HOMA-IR ↔ Glutathione peroxidase activity
De Luis <i>et al.</i> (2011) ⁽²⁸⁾	Spain, <i>n</i> 30 obese adults: twelve males and eighteen females (mean age 51 years; mean BMI 39.2 kg/m ²)	Parallel RCT double-blinded HMQS: 13	Random assignment to either 4-week prebiotic + ALA-supplemented cookies (2 cookies/d containing 2 g inulin + 3.1 g FOS + 3.2 g ALA) or 4-week placebo cookies (2 control cookies/d)	↓ Total cholesterol, LDL and CRP (only males) ↔ Total body weight, HOMA, TAG and HDL
Dewulf <i>et al.</i> (2013) ⁽²⁹⁾	Belgium, <i>n</i> 30 obese women (mean age 47.5 years; mean BMI 35.9 kg/m ²)	Parallel RCT double-blinded HMQS: 12	Random assignment to either a 3-month prebiotic-supplemented diet (16 g/d of inulin/oligofructose 50:50 mix) or a 3-month placebo-supplemented diet (16 g maltodextrin/d). All the subjects received dietary advice for weight reduction	↓ Glucose after OGTT ↔ Insulin after OGTT ↔ Fasting glucose and fasting insulin ↔ HbA1c and HOMA ↔ Lipid levels ↔ CRP
Genta <i>et al.</i> (2009) ⁽³⁰⁾	Argentina, <i>n</i> 35 obese women (mean age 40.5 years; mean BMI 33.5 kg/m ²)	Parallel RCT double-blinded HMQS: 10	Random assignment to either 17-week consumption of Yacon syrup (approximately 12.5 g FOS/d), <i>n</i> 20, or 17-week consumption of placebo syrup, <i>n</i> 15	↓ Total body weight, ↓ BMI ↓ Waist circumference ↓ Fasting serum insulin ↓ HOMA and serum LDL ↑ Satiety (only qualitative) ↔ Fasting serum glucose ↔ Serum total cholesterol, HDL and TAG

Table 2. Continued

Study	Participants	Study design/blinding	Dietary prebiotic intervention	Effect of dietary prebiotic supplements on metabolic outcomes
Giacco <i>et al.</i> (2004) ⁽³¹⁾	Italy, <i>n</i> 30 adults with plasma cholesterol concentrations between 5.17 and 7.76 mmol/l: twenty males and ten females (mean age 45.5 years; mean BMI 26.6 kg/m ²)	Cross-over RCT double-blinded HMQS: 10	Random assignment to either a 2-month prebiotic-supplemented diet (10.6 g short-chain FOS/d) or a 2-month placebo-supplemented diet (7.5 g maltodextrin/d) before cross-over. Washout period: not stated	↑ Fasting plasma Lp(a) ↓ Postprandial serum insulin ↔ Fasting plasma cholesterol and TAG ↔ Postprandial glucose, NEFA and TAG
Lecerf <i>et al.</i> (2012) ⁽³²⁾	France, <i>n</i> 59 healthy adults: twenty-six males and thirty-three females (mean age 20 years; mean BMI 21 kg/m ²)	Parallel RCT double-blinded HMQS: 10	Random assignment to a 4-week xylo-oligosaccharide supplement (5 g XOS/d) or a 4-week xylo-oligosaccharide + inulin supplement (1 g XOS + 3 g inulin/d) or 4-week placebo (4 g maltodextrin/d)	↓ Lipopolysaccharide ↓ IL-1β and TNF-α expression ↑ IL-13 and IL-10 expression
Letexier <i>et al.</i> (2003) ⁽³³⁾	France, <i>n</i> 8 healthy adults: four males and four females (age range 23–32 years; BMI range 19–25 kg/m ²)	Cross-over RCT double-blinded HMQS: 10	Random assignment to either a 3-week prebiotic-supplemented diet (10 g inulin/d) or a 3-week placebo-supplemented diet (10 g maltodextrin/d) before cross-over. Washout period: 4 months	↓ Hepatic lipogenesis ↓ Plasma TAG ↔ Total cholesterol, LDL and HDL ↔ Glucose, NEFA, insulin and glucagon
Lomax <i>et al.</i> (2012) ⁽³⁴⁾	UK, <i>n</i> 43 normal-weight and overweight adults: eleven males and thirty-two females (mean age 55 years; mean BMI 25 kg/m ²)	Parallel RCT double-blinded HMQS: 11	Random assignment to either a 4-week prebiotic-supplemented diet (8 g fructans/d) or a 4-week placebo-supplemented diet (8 g maltodextrin/d)	↔ Total body weight ↔ T-cell activation ↔ NK cell activity ↔ Cytokine production ↔ Monocyte respiratory burst ↔ Basal hepatic glucose production ↔ Fasting plasma glucose and insulin ↔ Fasting lipids, Lp(a) and NEFA ↔ ApoA1 and apoB
Luo <i>et al.</i> (2000) ⁽³⁵⁾	France, <i>n</i> 10 adults with type 2 diabetes: six males and 4 females (mean age 57 years; mean BMI 28 kg/m ² ; mean time after DM diagnosis 11 years; mean HbA1c levels 7.7%)	Cross-over RCT double-blinded HMQS: 10	Random assignment to either 4-week prebiotic-supplemented cookies (20 g FOS/d) or 4-week control-supplemented cookies (sucrose) before cross-over. Washout period: nil	↔ Total body weight ↔ T-cell activation ↔ NK cell activity ↔ Cytokine production ↔ Monocyte respiratory burst ↔ Basal hepatic glucose production ↔ Fasting plasma glucose and insulin ↔ Fasting lipids, Lp(a) and NEFA ↔ ApoA1 and apoB
Parnell <i>et al.</i> (2009) ⁽³⁶⁾	Canada, <i>n</i> 39 overweight and obese adults with BMI > 25 kg/m ² : seven males and thirty-two females (mean age 40 years; mean BMI 30 kg/m ²)	Parallel RCT double-blinded HMQS: 10	Random assignment to either a 12-week prebiotic-supplemented diet (21 g oligofructose/d) or a 12-week placebo-supplemented diet (7.9 g maltodextrin/d)	↓ Body weight and fat mass ↓ Energy intake ↓ Postprandial ghrelin ↓ Postprandial insulin ↔ Postprandial glucose ↔ Postprandial peptide YY and GLP-1
Peters <i>et al.</i> (2009) ⁽³⁷⁾	The Netherlands, <i>n</i> 21 normal-weight and overweight adults: five males and sixteen females (mean age 53 years; mean BMI 25.9 kg/m ²)	Cross-over RCT double-blinded HMQS: 10	Random assignment to 2 d consumption of breakfast prebiotic meal-replacement bar (8 g FOS) or breakfast prebiotic + barley meal-replacement bar (8 g FOS + 8 g barley) or breakfast barley meal-replacement bar (8 g barley) or breakfast control meal-replacement bar (oats)	↔ Appetite ↔ Satiety ↔ Energy intake
Piche <i>et al.</i> (2003) ⁽³⁸⁾	France, <i>n</i> 9 adults with gastro-oesophageal reflux disease: five males and four females (mean age 52 years).	Cross-over RCT double-blinded HMQS: 10	Random assignment to either a 1-week prebiotic-supplemented low-residue diet (19.8 g FOS/d) or a 1-week placebo-supplemented low-residue diet (sucrose) before cross-over. Washout period: minimum 3 weeks	↑ Postprandial plasma GLP-1 ↔ Postprandial peptide YY ↔ Postprandial cholecystokinin
Russo <i>et al.</i> (2008) ⁽³⁹⁾ Russo <i>et al.</i> (2010) ⁽⁴⁰⁾	Italy, <i>n</i> 15 healthy males (mean age 19 years; mean BMI 22.8 kg/m ²)	Cross-over RCT double-blinded HMQS: 9	Random assignment to either a 5-week prebiotic-supplemented diet (11% inulin-enriched pasta) or a 5-week placebo-supplemented diet (control wheat pasta) before cross-over. Washout period: 8 weeks	↓ Lp(a) and TAG ↑ HDL ↓ Total cholesterol:HDL ratio ↓ Fasting glucose and fructosamine ↓ HbA1c and HOMA-IR ↓ Gastric emptying

Metabolic benefits of dietary prebiotics

Table 2. Continued

Study	Participants	Study design/blinding	Dietary prebiotic intervention	Effect of dietary prebiotic supplements on metabolic outcomes
Russo <i>et al.</i> (2011) ⁽⁴¹⁾ Russo <i>et al.</i> (2012) ⁽⁴²⁾	Italy, <i>n</i> 20 healthy males (mean age 19 years; mean BMI 22.8 kg/m ²)	Cross-over RCT double-blinded HMQS: 9	Random assignment to either a 5-week prebiotic-supplemented diet (11 % inulin-enriched pasta) or a 5-week placebo-supplemented diet (control wheat pasta) before cross-over. Washout period: 8 weeks	↑ Neurotensin and somatostatin ↑ Plasma GLP-2 ↓ Gastric emptying ↓ Small-intestinal permeability ↓ Serum zonulin ↔ Corticotropin-releasing factor
Schiffirin <i>et al.</i> (2007) ⁽⁴³⁾	Switzerland, <i>n</i> 74 elderly adults with mild malnutrition: eighteen males and fifty-six females (mean age 84 years; mean BMI 25 kg/m ²)	Parallel RCT double-blinded HMQS: 12	Random assignment to either a 12-week prebiotic-supplemented drink (2–4 g FOS/d) or a 12-week identical drink (without FOS)	↓ IL-6 mRNA and TNF-α mRNA ↓ Soluble CD14
Seidel <i>et al.</i> (2007) ⁽⁴⁴⁾	Germany, <i>n</i> 38 males: twenty smokers and eighteen non-smokers (mean age 27 years; mean BMI 23.2 kg/m ²)	Parallel RCT double-blinded HMQS: 9	Subjects participated in a pre-randomisation run-in period (consumed at least 200 g wheat-rye bread/d for 5 weeks), followed by randomisation to an intervention period (nineteen participants consumed at least 200 g prebiotic bread/d for 5 weeks and nineteen participants consumed at least 200 g prebiotic + antioxidant bread/d), followed by a post-intervention period (all thirty-eight participants received a standard diet for 1 week). Intervention breads contained 4 g inulin/100 g	↑ TRAP ↑ PCL ↑ CD19 ↓ ICAM-1 ↓ CD3 + NK + ↑ CD3 + HLA-DR + (activated T cells) ↔ Body weight
Tovar <i>et al.</i> (2012) ⁽⁴⁵⁾	Mexico, <i>n</i> 110 overweight and obese women (age range 18–50 years; BMI ≥25 kg/m ²)	Parallel RCT not blinded HMQS: 8	Random assignment to 12-week partial meal replacement or 12-week partial meal replacement + prebiotic (10 g inulin/d) or 12-week prebiotic (10 g inulin/d) or 12-week control (no meal replacement or inulin). All the subjects received a low-energy diet	↓ TAG ↔ Total cholesterol, HDL and glucose
Verhoef <i>et al.</i> (2011) ⁽⁴⁶⁾	The Netherlands, <i>n</i> 29 normal-weight and overweight adults: nine males and twenty females (mean age 28 years; mean BMI 24.8 kg/m ²)	Cross-over RCT double-blinded HMQS: 9	Random assignment to either a 13 d prebiotic-supplemented diet (10 g FOS/d or 16 g FOS/d) or a 13 d placebo-supplemented diet (16 g maltodextrin) before cross-over. Washout period: 2 weeks	Outcomes for 16 g FOS/d intervention v. placebo: ↔ Appetite ↔ Satiety ↔ Energy intake ↔ Plasma GLP-1 ↑ Plasma peptide YY ↑ NK cell activity ↑ IL-10 production ↓ IL-6, IL-1β and TNF-α production ↔ Total cholesterol and HDL
Vulevic <i>et al.</i> (2008) ⁽⁴⁷⁾	UK, <i>n</i> 44 normal-weight and overweight elderly adults: sixteen males and twenty-eight females (age range 64–79 years; BMI range 22–31 kg/m ²)	Cross-over RCT double-blinded HMQS: 10	Random assignment to either a 10-week prebiotic-supplemented diet (5.5 g GOS/d) or a 10-week placebo-supplemented diet (5.5 g maltodextrin) before cross-over. Washout period: 4 weeks	↓ Plasma CRP and insulin ↓ Total cholesterol and TAG ↓ TC:HDL ratio
Vulevic <i>et al.</i> (2013) ⁽⁴⁸⁾	UK, <i>n</i> 45 overweight and obese adults with the metabolic syndrome: sixteen males and twenty-nine females (mean age 45 years; mean BMI 31 kg/m ²)	Cross-over RCT double-blinded HMQS: 10	Random assignment to either a 12-week prebiotic-supplemented diet (5.5 g GOS/d) or a 12-week placebo-supplemented diet (5.5 g maltodextrin) before cross-over. Washout period: 4 weeks	↔ IL-6, IL-10, IL-8 and TNF-α ↑ Mean fullness and minimum fullness
Whelan <i>et al.</i> (2006) ⁽⁴⁹⁾	UK, <i>n</i> 11 healthy adults: five males and six females (mean age 28 years; mean BMI 23.5 kg/m ²)	Cross-over RCT double-blinded HMQS: 10	Random assignment to either a 2-week prebiotic-supplemented liquid enteral formula (approximately 18 g pea fibre + 10 g FOS/d) or a 2-week placebo liquid enteral formula (standard formula) before cross-over. Washout period: 4 weeks	↑ Minimum satiety

RCT, randomised controlled trial; HMQS, Heyland Methodological Quality Score, where trials scoring ≥8 out of 14 points are considered to be of high methodological quality; FOS, fructo-oligosaccharide; GLP, glucagon-like peptide; NASH, non-alcoholic steatohepatitis; AST, aspartate aminotransferase; DM, diabetes mellitus; CRP, C-reactive protein; TAC, total antioxidant capacity; SOD, superoxide dismutase; HOMA-IR, homeostasis model assessment for insulin resistance; ALA, α-linolenic acid; OGTT, oral glucose tolerance test; Lp(a), lipoprotein (a); XOS, xylo-oligosaccharide; NK, natural killer; CD, cluster of differentiation; TRAP, total radical-trapping antioxidant parameter; PCL, photosensitive chemiluminescence; ICAM-1, intracellular adhesion molecule-1; GOS, galacto-oligosaccharide; TC, total cholesterol; ↓, significantly lower than that in the comparison diet group after intervention; ↑, significantly higher than that in the comparison diet group after intervention; ↔, no significant difference between the prebiotic-supplemented diet and control diet groups after intervention; HLA-DR, human leucocyte antigen-D-related.

after prebiotic consumption^(37,46) was not possible in the meta-analysis, as they did not report study data and did not provide results when contacted by the reviewers. An additional trial carried out in obese subjects was also excluded from the meta-analysis because it reported only qualitative improvements in satiety⁽³⁰⁾. Of the five trials measuring energy intakes in normal-weight and overweight participants and those with type 2 diabetes, three found a significant reduction in total energy consumption during the prebiotic intervention when compared with placebo^(22,26,36). However, the reduction in energy intake lost statistical significance after the meta-analysis (n 208) yielded a pooled SMD of -0.51 (95% CI $-1.20, 0.19$; $P=0.16$). The duration of one trial finding no difference in energy intake between control and intervention groups was 2 d⁽³⁷⁾, and the trials finding reduced energy intake by the intervention groups lasted a minimum of 2 weeks. Available evidence supported that dietary prebiotic supplementation for at least 2 weeks' duration increases circulating peptide YY concentrations in normal-weight and overweight adults^(23,36,46), but the effect was not statistically significant after the meta-analysis (n 100), with a combined SMD of -0.96 (95% CI $-1.98, 0.06$; $P=0.07$). Of four high-quality trials, two found increased GLP-1 concentrations after prebiotic supplementation in healthy and overweight subjects^(23,38). The increase in GLP-1 concentrations was not significant after the meta-analysis (n 117), with a pooled SMD of -0.32 (95% CI $-0.87, 0.23$; $P=0.25$). Each of the trials reported significant reductions in ghrelin concentrations⁽³⁶⁾ and increased GLP-2 concentrations⁽⁴²⁾ in subjects consuming dietary prebiotics. Contradictory results were reported by five trials examining the effect of prebiotic intervention on body weight. Significant reductions in body weight after prebiotic supplementation in comparison with placebo were reported by two trials^(30,36), while no change in body weight was observed in three trials^(26,28,44). Trials of longer duration (12–17 weeks) were more likely to observe reductions in body weight than shorter trials lasting 4–8 weeks. The meta-analysis (n 191) indicated a non-significant

reduction in body weight after prebiotic supplementation, with a pooled SMD of -0.48 (95% CI $-1.19, 0.23$; $P=0.19$).

Outcomes associated with glucose homeostasis

Of the four studies measuring the effect of prebiotic supplementation on postprandial glucose concentrations, two reported significant reductions in glycaemia in normal-weight and obese participants^(23,29). Following the meta-analysis (n 131), the pooled SMD for postprandial glucose concentrations was -0.76 (95% CI $-1.41, -0.07$; $P<0.05$), indicating a statistically significant effect supporting that prebiotic consumption results in the reduction of postprandial glucose concentrations (Fig. 3). Of three studies, two reported significant reductions in postprandial insulin concentrations following prebiotic intervention in overweight and hypercholesterolaemic subjects^(31,36). Meta-analysis of these trials (n 121) indicated a statistically significant reduction in postprandial insulin concentrations, with a combined SMD of -0.77 (95% CI $-1.50, -0.04$; $P<0.05$) (Fig. 4). Significant delays in gastric emptying times in healthy males consuming prebiotic supplements were found in two trials carried out by the same study group^(40,41). Studies investigating fasting glucose and fasting insulin concentrations and insulin resistance (HOMA-IR) reported conflicting results. Significant reductions in HbA1c levels in healthy participants after only 5 weeks of prebiotic supplementation⁽⁴⁰⁾ and in women with type 2 diabetes after 8 weeks⁽²⁶⁾ were found by two trials, while no change in HbA1c levels in obese women after prebiotic supplementation lasting 3 months was found by another trial⁽²⁹⁾.

Outcomes associated with cardiovascular and hepatic health

There was insufficient evidence to support that prebiotic supplementation reduces total cholesterol or LDL concentrations in healthy, obese or dyslipidaemic individuals, with the




Fig. 2. Effects of dietary prebiotic supplementation on self-reported satiety. Forest plot of standardised mean differences (SMD, 95% CI) for individual and pooled trials.




Fig. 3. Effects of dietary prebiotic supplementation on postprandial glucose concentrations. Forest plot of standardised mean differences (SMD, 95 % CI) for individual and pooled trials.

majority of studies finding no change in the concentrations of these lipids after intervention. Of the eleven trials investigating the effect of prebiotic supplementation on circulating TAG concentrations, five reported significant reductions in healthy, overweight or hypercholesterolaemic individuals compared with controls^(24,33,39,45,48). However, the remaining six trials that failed to detect changes in TAG concentrations were also carried out in healthy, overweight or hypercholesterolaemic subjects^(25,28–31,35). These trials were subjected to meta-analysis (n 402), resulting in a non-significant pooled SMD for TAG concentrations of -0.11 (95 % CI $-0.31, 0.08$; $P=0.26$) (Fig. 5). A significant reduction in serum aspartate aminotransferase concentrations was reported by one small trial carried out in people with NASH⁽²⁵⁾.

Outcomes associated with inflammation and immune function

Of the four trials investigating the impact of dietary prebiotic supplementation on circulating C-reactive protein (a biochemical marker of inflammation) concentrations, three found significant reductions in overweight and obese adults and

women with type 2 diabetes in comparison with controls^(26,28,48). Meta-analysis of these trials (n 181) indicated a non-significant reduction in C-reactive protein concentrations after prebiotic supplementation, however, with a pooled SMD of -0.85 (95 % CI $-2.11, 0.42$; $P=0.19$). Studies measuring the production of pro-inflammatory cytokines (TNF- α and IL) and immune cell activity (T-cell activation and natural killer cell activation) yielded contradictory results. Significant increases in the measures of antioxidant status (total antioxidant capacity⁽²⁷⁾, total radical-trapping antioxidant parameter and photosensitive chemiluminescence⁽⁴⁴⁾) were found by two studies, and a decrease in small-intestinal permeability⁽⁴²⁾ following prebiotic interventions was reported by one trial. Significant reductions in circulating lipopolysaccharide (LPS) concentrations after dietary prebiotic supplementation in healthy adults⁽³²⁾ and women with type 2 diabetes⁽²⁶⁾ were identified by two studies.

Discussion

Simple, safe and effective interventions are urgently needed to prevent and treat obesity and its associated co-morbidities. The human gut microbiota and its metabolites influence host




Fig. 4. Effects of dietary prebiotic supplementation on postprandial insulin concentrations. Forest plot of standardised mean differences (SMD, 95 % CI) for individual and pooled trials.




Fig. 5. Effects of dietary prebiotic supplementation on TAG concentrations. Forest plot of standardised mean differences (SMD, 95 % CI) for individual and pooled trials.

physiology, energy homeostasis, inflammatory processes and immune function both locally and within distal tissues. The use of dietary prebiotic supplements to promote the selective proliferation of beneficial intestinal microbes might represent an important nutritional strategy in the management of metabolic abnormalities and chronic disease.

Prebiotics and overweight/obesity

The SCFA acetate, propionate and butyrate are produced as by-products of bacterial prebiotic fermentation in the colon. In addition to representing a source of energy for the host, these SCFA play a number of beneficial roles including the maintenance of human intestinal health and modulation of metabolic and immune processes. SCFA are the only known ligands for two G protein-coupled receptors, GPR41 and GPR43, which are expressed in a variety of gastrointestinal cells and stimulate the secretion of hormones involved in the regulation of energy intake and expenditure. Binding of SCFA to GPR41 increases the production of peptide YY and GLP-1, hormones that reduce appetite, delay gastric emptying and increase insulin sensitivity⁽⁵⁰⁾. SCFA also promote the differentiation of intestinal L-cells, contributing to increased endogenous GLP-1 production⁽⁵¹⁾.

In animal studies, dietary supplementation of the SCFA butyrate has been found to prevent diet-induced obesity and

improve insulin sensitivity with a concomitant increase in energy expenditure and fatty acid oxidation and an increase in mitochondrial respiration⁽⁵²⁾. In mice, the selective growth of certain *Lactobacillus* species in the colon has been found to reduce body fat storage through the up-regulation of *Fiaf* (fasting-induced adipose factor) gene expression and inhibition of lipoprotein lipase^(53,54). Indeed, several animal studies have demonstrated the protective effects of prebiotics on the development of obesity and insulin resistance^(55,56); however, more robust human studies are required to confirm the protective effects of prebiotics on these pathways in human physiology.

The present review found consensus among three of the five high-quality trials supporting that the daily consumption of a prebiotic supplement for a minimum of 2 weeks increases satiety cues in healthy adults. However, these findings were based on self-reports from relatively small numbers of subjects (*n* 81) and prebiotic supplementation failed to result in significant weight reduction. Weight reduction was unlikely to be observed in these trials due to the short duration of their prebiotic interventions (2d–2 weeks). The addition of pea fibre to the prebiotic supplement confounded one trial reporting an increase in satiety after 2 weeks of prebiotic consumption, making it difficult to draw conclusions about the action of either type of fibre individually⁽⁴⁹⁾. Of the two trials that did not detect any change in satiety after prebiotic supplementation,

one involved an intervention period of 2 d⁽³⁷⁾, which may have been insufficient time to modify the growth and activity of intestinal bacteria to influence changes in host physiology. Increases in breath hydrogen production in the intervention group, indicative of enhanced intestinal bacterial fermentation, were detected by one of the trials investigating the effects of prebiotic supplementation on satiety sensations⁽²³⁾. However, no trials analysed the stool samples of participants for changes in microbial growth; therefore, alterations in hunger and fullness reported by the subjects in these studies may have occurred independently of any changes in gut microbial fermentation. Prebiotics are soluble fibres capable of modifying the intestinal transit of food due to their water-binding and bulking capacity⁽⁵⁷⁾. Indeed, study participants consuming prebiotics within a single meal have reported increased levels of satiety, well before any changes in colonic bacterial growth could have taken place^(58,59). Trials quantitatively evaluating the effect of prebiotic consumption on satiety and gut bacterial growth in overweight and obese individuals are now required.

Of the five high-quality trials, two provided consistent evidence favouring dietary prebiotic consumption for at least 2 weeks' duration for the reduction of total energy intake in normal-weight and overweight individuals and in women with type 2 diabetes. However, the pooled reduction in energy consumption was not statistically significant after the meta-analysis. A longer prebiotic supplementation period lasting 12 weeks was required before participants' reduced energy intake resulted in significant weight reduction⁽³⁶⁾. Where body composition was evaluated, the weight lost was predominantly fat mass rather than lean tissue or fluid⁽³⁶⁾. Trials of extended duration are now needed to determine whether dietary prebiotic consumption is a safe and effective therapeutic option for long-term weight and body fat reduction or whether physiological adaptations by the host eventually compensate for this energy imbalance to minimise weight loss.

The majority of trials investigating the effect of dietary prebiotic supplementation on the regulation of intestinal peptide (peptide YY) and incretin (GLP-1) secretion reported significant increases in the production of these molecules after 2 weeks, but the combined changes were not significant after the meta-analysis. The unique role played by prebiotics and specific bacteria in gut hormone kinetics requires further investigation, as non-prebiotic dietary fibres have also been reported to be associated with increased SCFA, peptide YY and GLP-1 production in human feeding studies^(60,61).

Prebiotics and glucose intolerance

Reduced levels of bifidobacteria and lactobacilli and increased gastrointestinal permeability are found in mice consuming a diet high in saturated fat when compared with those consuming a standard diet. The provision of dietary prebiotic supplements subsequently restores the growth of these beneficial bacterial species and improves the integrity of the gut barrier^(11,62). Animal studies have shown a causal link between the consumption of a high-fat diet and increased intestinal levels of LPS-containing bacteria, or concentrations

of circulating LPS, and the development of obesity and insulin resistance⁽⁶³⁾. LPS is the major component of the outer membrane of Gram-negative bacteria and is composed of a hydrophobic lipid (lipid A), a hydrophilic core oligosaccharide and a repeating hydrophilic polysaccharide side chain (O-antigen). In the setting of a high-fat diet, LPS is able to translocate from the intestine into the host circulation, resulting in 'metabolic endotoxaemia'⁽⁶⁴⁾. LPS stimulates the overproduction of reactive oxygen species and pro-inflammatory cytokines by macrophages, resulting in subclinical systemic inflammation, weight gain and insulin resistance development^(65,66). Human subjects with type 2 diabetes have been found to possess serum endotoxin levels that are 2-fold higher than those observed in non-diabetic controls⁽⁶⁷⁾. Metabolic endotoxaemia is also positively correlated with total energy intake and fasting insulin concentrations in the non-diabetic population⁽⁶⁸⁾. In mice with high-fat diet-induced metabolic endotoxaemia, nutritional supplementation with prebiotics restores intestinal levels of Gram-positive bacteria, improves glucose tolerance and reduces circulating concentrations of LPS and pro-inflammatory cytokines⁽⁵⁶⁾.

Prebiotics and their fermentation products have been shown to reduce gastrointestinal permeability by a variety of mechanisms. The SCFA butyrate is involved in the maintenance of gut epithelial integrity by acting as the principal fuel for colonocytes and promoting the transcription of tight junction proteins between gastrointestinal cells⁽⁶⁹⁾. Butyrate also reduces gastrointestinal permeability by enhancing the activation of the peroxisomal proliferator-activated receptor gamma (*PPAR* γ) gene, a nuclear factor receptor involved in the attenuation of inflammation in colonic epithelial cells^(70,71). Prebiotic-induced changes in gut microbiota also increase the endogenous production of GLP-2, which enhances gut barrier function by promoting the proliferation of crypt cells^(12,72).

The present review found general agreement among trials supporting that the consumption of dietary prebiotic supplements reduces postprandial glucose and insulin concentrations in healthy and overweight individuals. Pooled reductions in postprandial glucose and insulin concentrations were statistically significant after the meta-analysis. High-quality randomised controlled trials conducted in subjects with either impaired glucose tolerance⁽⁷³⁾ or type 2 diabetes⁽⁷⁴⁾ have also found reduced postprandial serum insulin concentrations after the consumption of arabinoxylan (a potential prebiotic fibre). Whether these results were mediated by alterations in intestinal bacterial growth or activity is unclear, as stool samples were not analysed in these trials. Significant delays in gastric emptying times after prebiotic supplementation in healthy males were found by two studies. However, these trials were conducted by the same research group, and it is unclear whether some subjects participated in both the studies. Therefore, further independent research is required before definitive conclusions can be drawn about the effects of prebiotic consumption on gastric emptying. The findings of studies investigating fasting glucose and fasting insulin concentrations and insulin resistance (HOMA-IR) after prebiotic supplementation were contradictory. Long-term prebiotic intervention studies in people with pre-diabetes

or the metabolic syndrome are now required to determine whether prebiotics confer some protection against the future development of type 2 diabetes in high-risk individuals.

Prebiotics and dyslipidaemia

The abundance of particular bacterial species in the gut has been shown to be positively correlated with serum total cholesterol and LDL-cholesterol concentrations in subjects with CVD⁽⁷⁵⁾. It has also been hypothesised that bacteria found in atherosclerotic plaques may have originated from the gastrointestinal tract, as the DNA of specific micro-organisms can be found in both the colon and coronary atheroma of the same individual⁽⁷⁶⁾. As the development of CVD involves multiple pro-inflammatory pathways, it is plausible that pathogenic microbes may potentiate inflammation within atherosclerotic plaques by delivering macrophages to the arterial wall and stimulating their production of reactive oxygen species and cytokines or their conversion to foam cells⁽⁷⁷⁾.

Propionate, a SCFA product of prebiotic fermentation, may play a significant role in the modification of hepatic lipid metabolism. In the liver, propionate is a possible substrate for gluconeogenesis and may contribute to the inhibition of cholesterol synthesis by altering the activity of 3-hydroxy-3-methylglutaryl-CoA reductase⁽⁵⁾. In addition, prebiotic supplementation might attenuate cholesterol and TAG production by stimulating the synthesis of *cis*-9, *trans*-11-conjugated linoleic acid from PUFA by beneficial bacterial species. This isoform of *cis*-9, *trans*-11-conjugated linoleic acid has been shown to reduce cholesterol and TAG concentrations in animal studies⁽⁷⁸⁾, but the results of human trials are less conclusive. Gut microbes are also an essential requirement for the production of secondary bile acids in the colon. These bile acids are de-conjugated and are therefore unavailable for enterohepatic recirculation. As a result, the liver is forced to produce additional bile acids from circulating cholesterol⁽⁷⁹⁾.

The findings of human intervention studies investigating the effect of dietary prebiotic supplementation on circulating total and LDL-cholesterol concentrations were contradictory. Of the two studies reporting significant reductions in total cholesterol concentrations, one found a reduction in only male subjects and was complicated by the use of an intervention containing both prebiotics and α -linolenic acid⁽²⁸⁾. α -Linolenic acid may have contributed to the cholesterol-lowering effect in this instance. There is limited evidence to support that prebiotic supplementation reduces total or LDL-cholesterol concentrations in hypercholesterolaemic individuals, as the only two trials conducted in participants with hypercholesterolaemia found no significant changes in total cholesterol, LDL-cholesterol or HDL-cholesterol concentrations^(24,31). However, these trials involved short-term prebiotic intervention periods (3–8 weeks' duration) and studies of longer duration are therefore required.

The present review found conflicting evidence describing the effect of prebiotic supplementation on circulating TAG concentrations in healthy, overweight/obese and hypercholesterolaemic subjects. In addition to a prebiotic supplement, one study⁽⁴⁵⁾ provided a low-energy diet co-intervention

to all the trial participants, making it difficult to establish whether the TAG-lowering effect was associated with prebiotic-induced intestinal microbial changes alone or whether weight reduction or a reduced fat intake together with the action of the modified microflora produced a synergistic effect. Additional systematic reviews exploring this topic have also reported equivocal conclusions. The use of prebiotics for the reduction of TAG concentrations in humans regardless of health condition was favoured by one review of trials published between 1995 and 2005⁽⁸⁰⁾, with the majority of trials being conducted in normolipidaemic individuals. The other meta-analysis of trials published between 1999 and 2010 supported the TAG-lowering effects of inulin in only hypercholesterolaemic subjects⁽⁷⁵⁾, but the reduction in TAG concentrations lost significance when results from both hyperlipidaemic and normolipidaemic subjects were combined. The present review also found a non-significant reduction in TAG concentrations after the meta-analysis of pooled trials. Future trials must simultaneously quantify lipid concentrations and gut bacterial growth and activity to determine whether prebiotic-induced modulation of the intestinal flora contributes to the reduction of serum TAG concentrations.

Prebiotics and non-alcoholic steatohepatitis

NASH is an asymptomatic disease characterised by fatty infiltration of the liver and inflammation, which can eventually lead to fibrosis, cirrhosis, portal hypertension, hepatocellular carcinoma and liver failure⁽⁸¹⁾. Obesity, dyslipidaemia, insulin resistance and diabetes have frequently been reported to be associated with the development of NASH. Increased plasma endotoxaemia, overproduction of inflammatory cytokines and excessive oxidative stress within hepatic cells are also thought to contribute to the pathogenesis of NASH. The use of dietary prebiotic supplements to restore an optimal microbial balance within the gastrointestinal tract of individuals with NASH may assist in the reduction of TAG accumulation in the liver, attenuate inflammation and promote hepatic secretion of lipoproteins such as VLDL⁽¹⁵⁾. The translocation of Gram-negative bacteria from the intestine into the circulation has been reported to be associated with an increased severity of cirrhosis⁽⁸²⁾. By maintaining gut barrier function and reducing bacterial translocation, prebiotics may be effective in the management of liver disease complications⁽⁸³⁾.

Studies exploring the effects of dietary prebiotic consumption on metabolic parameters in individuals with NASH are scarce. In the present review, one small trial involving seven adult males was included, which found a significant reduction in serum aspartate aminotransferase concentrations after prebiotic supplementation⁽²⁵⁾. This finding was supported by a larger randomised controlled trial (*n* 66), which administered a synbiotic (fructo-oligosaccharide + *Bifidobacterium longum*) to participants for 24 weeks⁽⁸⁴⁾. In addition to a significant reduction in serum aspartate aminotransferase concentrations, researchers found a reduction in the concentrations of circulating cytokines (TNF- α) and markers of inflammation (C-reactive protein), reduced concentrations of serum LDL-cholesterol and

endotoxins, an improvement in insulin sensitivity (HOMA-IR) and a reduction in hepatic steatosis (determined by liver biopsy). More research is required in this potentially very promising area of study.

Prebiotics and immune cell dysfunction

Gut microbiota, innate immune function and metabolism are inextricably linked, with early pathological processes occurring at the molecular level (subclinical inflammation, immune cell activation, increased oxidative and endoplasmic reticulum stress, altered production of vascular adhesion molecules and advanced glycation end products) contributing to the eventual development of metabolic disturbances such as hyperlipidaemia, atherosclerosis, insulin resistance and weight gain. Colonic bacteria and their prebiotic fermentation products may play a key role in the modulation of immune function by both increasing host resistance to infection and down-regulating inappropriate immune responses in the case of allergic reactions or chronic inflammatory conditions⁽⁵⁾. By maintaining the integrity of the gastrointestinal barrier, prebiotics reduce the invasion of pathogenic intestinal bacteria and their products (including LPS) into the circulation, preventing downstream immune cell activation. Prebiotics are thought to encourage increased intestinal mucin production, protecting the intestinal wall from bacterial adherence and invasion⁽⁸⁵⁾. SCFA produced as a by-product of bacterial prebiotic fermentation interact with GPR41 and GPR43 receptors on neutrophils and inhibit NF- κ B activation, reducing the production of pro-inflammatory cytokines⁽⁸⁶⁾. Additional bacterial fermentation products such as polysaccharide A and peptidoglycan exert anti-inflammatory effects on the host immune system⁽⁸⁷⁾.

There is insufficient evidence at present to recommend dietary prebiotics for the modulation of immune function to improve cardiometabolic health. There are very few human trials available, and most have reported contradictory findings. Although individual studies have found significant increases in the measures of antioxidant capacity and reductions in small-intestinal permeability and circulating LPS concentrations after prebiotic interventions, further studies are required to verify these results. Inulin exhibits antioxidant properties independent of altering gut bacterial growth and is able to scavenge a number of reactive oxygen species, which may help to reduce lipid peroxidation in the stomach⁽⁸⁸⁾. Future studies must distinguish between health benefits derived solely from the consumption of soluble fibres and those associated with the growth and activity of beneficial gut microbes. Future intervention studies exploring the effect of dietary prebiotics on immune function need to be conducted in healthy individuals who are subsequently exposed to an immune challenge.

Conclusions

Although animal studies have provided convincing evidence to support the beneficial role of prebiotics in metabolic health, the results of human trials to date have been less conclusive. Research involving laboratory animals enables the provision of tightly controlled diets, whereas studies involving

free-living humans are complicated by the variety of foods consumed by individuals from day to day. Some human studies have been complicated by the use of nutritional supplements containing prebiotics in combination with additional health-promoting components such as live bacteria, antioxidants and other dietary fibres, making it difficult to attribute changes in metabolism to prebiotics alone. To rule out cardiometabolic benefits associated with concomitant nutrients, prebiotic supplements in their pure form must be used in future trials.

In addition to bifidobacteria and lactobacilli, dietary prebiotics modulate the growth of numerous other gastrointestinal micro-organisms, the identity and function of which have not yet been fully characterised. Different species of bifidobacteria also have a variety of functions, which require further elucidation. Prebiotics are likely to undergo cross-fermentation by other microbial species of unknown benefit to the host. Bacterial analyses of human stool samples provide information only about the micro-organisms inhabiting the colon and are unlikely to accurately reflect the microbial composition of the proximal intestine. Responses to dietary prebiotics are variable in humans, with bifidogenic potential being affected by an individual's age, body weight, antibiotic use, dietary macronutrient intake, physical activity and baseline levels of colonic bifidobacteria^(89,90). More research is required to determine host lifestyle behaviours capable of promoting intestinal normobiosis and to establish the optimal prebiotic dose required to maximise health benefits.

In conclusion, the present review found convincing evidence from short-term high-quality human trials supporting the use of dietary prebiotics as a potential therapeutic intervention for the regulation of appetite and the reduction of circulating postprandial glucose and insulin concentrations. Further studies are needed to correlate these findings with changes in the growth and function of specific gut bacteria. There is insufficient evidence at present to recommend dietary prebiotics for reducing energy intake and body weight, increasing gastric peptide YY and GLP-1 secretion, improving insulin sensitivity, lowering lipid levels and modulating immune function. Long-term prospective trials investigating primary metabolic end points are now required.

Acknowledgements

N. J. K. was the recipient of National Health and Medical Research Council (NHMRC) Postgraduate Public Health Scholarship APP1039709. N. J. K. thanks Mr Brendan Kellow for IT assistance.

N. J. K. was supported by NHMRC Postgraduate Public Health Scholarship APP1039709. The NHMRC had no role in the design and analysis of this work or in the writing of this article.

The authors' contributions are as follows: N. J. K. designed the review, conducted the literature search, extracted and analysed the data and drafted the manuscript; M. T. C. and C. M. R. provided comments on the manuscript. All the authors read and approved the final manuscript.

None of the authors has any conflicts of interest to declare.

References

1. Diamant M, Blaak EE & de Vos WM (2011) Do nutrient–gut–microbiota interactions play a role in human obesity, insulin resistance and type 2 diabetes? *Obes Rev* **12**, 272–281.
2. Wu X, Ma C, Han L, *et al.* (2010) Molecular characterisation of the faecal microbiota in patients with type II diabetes. *Curr Microbiol* **61**, 69–78.
3. Larsen N, Vogensen FK, van den Berg FW, *et al.* (2010) Gut microbiota in human adults with type 2 diabetes differs from non-diabetic adults. *PLoS ONE* **5**, e9085.
4. Gibson GR, Probert HM, Loo JV, *et al.* (2004) Dietary modulation of the human colonic microbiota: updating the concept of prebiotics. *Nutr Res Rev* **17**, 259–275.
5. Roberfroid M, Gibson GR, Hoyles L, *et al.* (2010) Prebiotic effects: metabolic and health benefits. *Br J Nutr* **104**, Suppl. 2, S1–S63.
6. Ramirez-Farias C, Slezak K, Fuller Z, *et al.* (2009) Effect of inulin on the human gut microbiota: stimulation of *Bifidobacterium adolescentis* and *Faecalibacterium prausnitzii*. *Br J Nutr* **101**, 541–550.
7. Chung H & Kasper DL (2010) Microbiota-stimulated immune mechanisms to maintain gut homeostasis. *Curr Opin Immunol* **22**, 455–460.
8. Ding S, Chi MM, Scull BP, *et al.* (2010) High-fat diet: bacteria interactions promote intestinal inflammation which precedes and correlates with obesity and insulin resistance in mouse. *PLoS ONE* **5**, e12191.
9. Backhed F, Manchester JK, Semenkovich CF, *et al.* (2007) Mechanisms underlying the resistance to diet-induced obesity in germ-free mice. *Proc Natl Acad Sci U S A* **104**, 979–984.
10. Havenaar R (2011) Intestinal health functions of colonic microbial metabolites: a review. *Benef Microbes* **2**, 103–114.
11. Cani PD, Neyrinck AM, Fava F, *et al.* (2007) Selective increases of bifidobacteria in gut microflora improve high-fat-diet-induced diabetes in mice through a mechanism associated with endotoxaemia. *Diabetologia* **50**, 2374–2383.
12. Cani PD, Possemiers S, Van de Wiele T, *et al.* (2009) Changes in gut microbiota control inflammation in obese mice through a mechanism involving GLP-2-driven improvement of gut permeability. *Gut* **58**, 1091–1103.
13. Hill MJ (1997) Intestinal flora and endogenous vitamin synthesis. *Eur J Cancer Prev* **6**, Suppl. 1, S43–S45.
14. Scholz-Ahrens KE, Ade P, Marten B, *et al.* (2007) Prebiotics, probiotics, and synbiotics affect mineral absorption, bone mineral content, and bone structure. *J Nutr* **137**, 838S–846S.
15. Musso G, Gambino R & Cassader M (2011) Interactions between gut microbiota and host metabolism predisposing to obesity and diabetes. *Annu Rev Med* **62**, 361–380.
16. Neyrinck AM & Delzenne NM (2010) Potential interest of gut microbial changes induced by non-digestible carbohydrates of wheat in the management of obesity and related disorders. *Curr Opin Clin Nutr Metab Care* **13**, 722–728.
17. Cloetens L, Ulmius M, Johansson-Persson A, *et al.* (2012) Role of dietary beta-glucans in the prevention of the metabolic syndrome. *Nutr Rev* **70**, 444–458.
18. Kootte RS, Vrieze A, Holleman F, *et al.* (2012) The therapeutic potential of manipulating gut microbiota in obesity and type 2 diabetes mellitus. *Diabetes Obes Metab* **14**, 112–120.
19. Heyland DK, Novak F, Drover JW, *et al.* (2001) Should immunonutrition become routine in critically ill patients? A systematic review of the evidence. *JAMA* **286**, 944–953.
20. White IR & Thomas J (2005) Standardized mean differences in individually-randomized and cluster-randomized trials, with applications to meta-analysis. *Clin Trials* **2**, 141–151.
21. Bunout D, Hirsch S, Pia de la Maza M, *et al.* (2002) Effects of prebiotics on the immune response to vaccination in the elderly. *JPEN J Parenter Enteral Nutr* **26**, 372–376.
22. Cani PD, Joly E, Horsmans Y, *et al.* (2006) Oligofructose promotes satiety in healthy human: a pilot study. *Eur J Clin Nutr* **60**, 567–572.
23. Cani PD, Lecourt E, Dewulf EM, *et al.* (2009) Gut microbiota fermentation of prebiotics increases satietogenic and incretin gut peptide production with consequences for appetite sensation and glucose response after a meal. *Am J Clin Nutr* **90**, 1236–1243.
24. Causey JL (2000) Effects of dietary inulin on serum lipids, blood glucose and the gastrointestinal environment in hypercholesterolemic men. *Nutr Res* **20**, 191–201.
25. Daubioul CA, Horsmans Y, Lambert P, *et al.* (2005) Effects of oligofructose on glucose and lipid metabolism in patients with nonalcoholic steatohepatitis: results of a pilot study. *Eur J Clin Nutr* **59**, 723–726.
26. Dehghan P, Gargari BP, Jafar-Abadi MA, *et al.* (2013) Inulin controls inflammation and metabolic endotoxemia in women with type 2 diabetes mellitus: a randomized-controlled clinical trial. *Int J Food Sci Nutr* (Epublication ahead of print version 24 September 2013).
27. Pourghassem Gargari B, Dehghan P, Aliasgharzadeh A, *et al.* (2013) Effects of high performance inulin supplementation on glycemic control and antioxidant status in women with type 2 diabetes. *Diabetes Metab J* **37**, 140–148.
28. de Luis DA, de la Fuente B, Izaola O, *et al.* (2011) Double blind randomized clinical trial controlled by placebo with an alpha linoleic acid and prebiotic enriched cookie on risk cardiovascular factor in obese patients. *Nutr Hosp* **26**, 827–833.
29. Dewulf EM, Cani PD, Claus SP, *et al.* (2013) Insight into the prebiotic concept: lessons from an exploratory, double blind intervention study with inulin-type fructans in obese women. *Gut* **62**, 1112–1121.
30. Genta S, Cabrera W, Habib N, *et al.* (2009) Yacon syrup: beneficial effects on obesity and insulin resistance in humans. *Clin Nutr* **28**, 182–187.
31. Giacco R, Clemente G, Luongo D, *et al.* (2004) Effects of short-chain fructo-oligosaccharides on glucose and lipid metabolism in mild hypercholesterolaemic individuals. *Clin Nutr* **23**, 331–340.
32. Lecerf JM, Depeint F, Clerc E, *et al.* (2012) Xylo-oligosaccharide (XOS) in combination with inulin modulates both the intestinal environment and immune status in healthy subjects, while XOS alone only shows prebiotic properties. *Br J Nutr* **108**, 1847–1858.
33. Letexier D, Diraison F & Beylot M (2003) Addition of inulin to a moderately high-carbohydrate diet reduces hepatic lipogenesis and plasma triacylglycerol concentrations in humans. *Am J Clin Nutr* **77**, 559–564.
34. Lomax AR, Cheung LV, Tuohy KM, *et al.* (2012) β -2-Fructans have a bifidogenic effect in healthy middle-aged human subjects but do not alter immune responses examined in the absence of an *in vivo* immune challenge: results from a randomised controlled trial. *Br J Nutr* **108**, 1818–1828.
35. Luo J, Van Yperselle M, Rizkalla SW, *et al.* (2000) Chronic consumption of short-chain fructooligosaccharides does not affect basal hepatic glucose production or insulin resistance in type 2 diabetics. *J Nutr* **130**, 1572–1577.
36. Parnell JA & Reimer RA (2009) Weight loss during oligofructose supplementation is associated with decreased ghrelin and increased peptide YY in overweight and obese adults. *Am J Clin Nutr* **89**, 1751–1759.

37. Peters HP, Boers HM, Haddeman E, *et al.* (2009) No effect of added beta-glucan or of fructooligosaccharide on appetite or energy intake. *Am J Clin Nutr* **89**, 58–63.
38. Piche T, des Varannes SB, Sacher-Huvelin S, *et al.* (2003) Colonic fermentation influences lower esophageal sphincter function in gastroesophageal reflux disease. *Gastroenterology* **124**, 894–902.
39. Russo F, Chimienti G, Riezzo G, *et al.* (2008) Inulin-enriched pasta affects lipid profile and Lp(a) concentrations in Italian young healthy male volunteers. *Eur J Nutr* **47**, 453–459.
40. Russo F, Riezzo G, Chiloiro M, *et al.* (2010) Metabolic effects of a diet with inulin-enriched pasta in healthy young volunteers. *Curr Pharm Des* **16**, 825–831.
41. Russo F, Clemente C, Linsalata M, *et al.* (2011) Effects of a diet with inulin-enriched pasta on gut peptides and gastric emptying rates in healthy young volunteers. *Eur J Nutr* **50**, 271–277.
42. Russo F, Linsalata M, Clemente C, *et al.* (2012) Inulin-enriched pasta improves intestinal permeability and modifies the circulating levels of zonulin and glucagon-like peptide 2 in healthy young volunteers. *Nutr Res* **32**, 940–946.
43. Schiffrin EJ, Thomas DR, Kumar VB, *et al.* (2007) Systemic inflammatory markers in older persons: the effect of oral nutritional supplementation with prebiotics. *J Nutr Health Aging* **11**, 475–479.
44. Seidel C, Boehm V, Vogelsang H, *et al.* (2007) Influence of prebiotics and antioxidants in bread on the immune system, antioxidative status and antioxidative capacity in male smokers and non-smokers. *Br J Nutr* **97**, 349–356.
45. Tovar AR, Caamano Mdel C, Garcia-Padilla S, *et al.* (2012) The inclusion of a partial meal replacement with or without inulin to a calorie restricted diet contributes to reach recommended intakes of micronutrients and decrease plasma triglycerides: a randomized clinical trial in obese Mexican women. *Nutr J* **11**, 44.
46. Verhoeve SP, Meyer D & Westerterp KR (2011) Effects of oligofructose on appetite profile, glucagon-like peptide 1 and peptide YY3-36 concentrations and energy intake. *Br J Nutr* **106**, 1757–1762.
47. Vulevic J, Drakoularakou A, Yaqoob P, *et al.* (2008) Modulation of the fecal microflora profile and immune function by a novel *trans*-galactooligosaccharide mixture (B-GOS) in healthy elderly volunteers. *Am J Clin Nutr* **88**, 1438–1446.
48. Vulevic J, Juric A, Tzortzis G, *et al.* (2013) A mixture of *trans*-galactooligosaccharides reduces markers of metabolic syndrome and modulates the fecal microbiota and immune function of overweight adults. *J Nutr* **143**, 324–331.
49. Whelan K, Efthymiou L, Judd PA, *et al.* (2006) Appetite during consumption of enteral formula as a sole source of nutrition: the effect of supplementing pea-fibre and fructooligosaccharides. *Br J Nutr* **96**, 350–356.
50. Flint HJ, Scott KP, Louis P, *et al.* (2012) The role of the gut microbiota in nutrition and health. *Nat Rev Gastroenterol Hepatol* **9**, 577–589.
51. Cani PD, Hoste S, Guiot Y, *et al.* (2007) Dietary non-digestible carbohydrates promote L-cell differentiation in the proximal colon of rats. *Br J Nutr* **98**, 32–37.
52. Gao Z, Yin J, Zhang J, *et al.* (2009) Butyrate improves insulin sensitivity and increases energy expenditure in mice. *Diabetes* **58**, 1509–1517.
53. Backhed F, Ding H, Wang T, *et al.* (2004) The gut microbiota as an environmental factor that regulates fat storage. *Proc Natl Acad Sci U S A* **101**, 15718–15723.
54. Aronsson L, Huang Y, Parini P, *et al.* (2010) Decreased fat storage by *Lactobacillus paracasei* is associated with increased levels of angiopoietin-like 4 protein (ANGPTL4). *PLoS ONE* **5**, e13087.
55. Parnell JA & Reimer RA (2012) Prebiotic fiber modulation of the gut microbiota improves risk factors for obesity and the metabolic syndrome. *Gut Microbes* **3**, 29–34.
56. Everard A, Lazarevic V, Derrien M, *et al.* (2011) Responses of gut microbiota and glucose and lipid metabolism to prebiotics in genetic obese and diet-induced leptin-resistant mice. *Diabetes* **60**, 2775–2786.
57. Nyman M (2002) Fermentation and bulking capacity of indigestible carbohydrates: the case of inulin and oligofructose. *Br J Nutr* **87**, Suppl. 2, S163–S168.
58. Perrigue MM, Monsivais P & Drewnowski A (2009) Added soluble fiber enhances the satiating power of low-energy-density liquid yogurts. *J Am Diet Assoc* **109**, 1862–1868.
59. Tarini J & Wolever TM (2010) The fermentable fibre inulin increases postprandial serum short-chain fatty acids and reduces free-fatty acids and ghrelin in healthy subjects. *Appl Physiol Nutr Metab* **35**, 9–16.
60. Costabile A, Klinder A, Fava F, *et al.* (2008) Whole-grain wheat breakfast cereal has a prebiotic effect on the human gut microbiota: a double-blind, placebo-controlled, crossover study. *Br J Nutr* **99**, 110–120.
61. Freeland KR, Wilson C & Wolever TM (2010) Adaptation of colonic fermentation and glucagon-like peptide-1 secretion with increased wheat fibre intake for 1 year in hyperinsulinaemic human subjects. *Br J Nutr* **103**, 82–90.
62. Serino M, Luche E, Gres S, *et al.* (2012) Metabolic adaptation to a high-fat diet is associated with a change in the gut microbiota. *Gut* **61**, 543–553.
63. Cani PD, Amar J, Iglesias MA, *et al.* (2007) Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes* **56**, 1761–1772.
64. Cani PD, Osto M, Geurts L, *et al.* (2012) Involvement of gut microbiota in the development of low-grade inflammation and type 2 diabetes associated with obesity. *Gut Microbes* **3**, 279–288.
65. Nakamura YK & Omaye ST (2012) Metabolic diseases and pro- and prebiotics: Mechanistic insights. *Nutr Metab (Lond)* **9**, 60.
66. Liang H, Hussey SE, Sanchez-Avila A, *et al.* (2013) Effect of lipopolysaccharide on inflammation and insulin action in human muscle. *PLOS ONE* **8**, e63983.
67. Creely SJ, McTernan PG, Kusminski CM, *et al.* (2007) Lipopolysaccharide activates an innate immune system response in human adipose tissue in obesity and type 2 diabetes. *Am J Physiol Endocrinol Metab* **292**, E740–E747.
68. Amar J (2010) Energy intake is associated with endotoxemia induces adipose inflammation and insulin resistance in humans. *Diabetes* **59**, 9.
69. Delzenne NM & Cani PD (2011) Interaction between obesity and the gut microbiota: relevance in nutrition. *Annu Rev Nutr* **31**, 15–31.
70. Eun CS, Han DS, Lee SH, *et al.* (2006) Attenuation of colonic inflammation by PPARgamma in intestinal epithelial cells: effect on Toll-like receptor pathway. *Dig Dis Sci* **51**, 693–697.
71. Lewis K, Lutgendorff F, Phan V, *et al.* (2010) Enhanced translocation of bacteria across metabolically stressed epithelia is reduced by butyrate. *Inflamm Bowel Dis* **16**, 1138–1148.
72. Cani PD (2012) Crosstalk between the gut microbiota and the endocannabinoid system: impact on the gut barrier function and the adipose tissue. *Clin Microbiol Infect* **18**, Suppl. 4, 50–53.
73. Garcia AL, Otto B, Reich SC, *et al.* (2007) Arabinoxylan consumption decreases postprandial serum glucose, serum insulin and plasma total ghrelin response in subjects with impaired glucose tolerance. *Eur J Clin Nutr* **61**, 334–341.

74. Lu ZX, Walker KZ, Muir JG, *et al.* (2004) Arabinoxylan fibre improves metabolic control in people with type II diabetes. *Eur J Clin Nutr* **58**, 621–628.
75. Guo Z, Liu XM, Zhang QX, *et al.* (2012) Effects of inulin on the plasma lipid profile of normolipidemic and hyperlipidemic subjects: a meta-analysis of randomized controlled trials. *Clin Lipidol* **7**, 215–222.
76. Koren O, Spor A, Felin J, *et al.* (2011) Human oral, gut, and plaque microbiota in patients with atherosclerosis. *Proc Natl Acad Sci U S A* **108**, Suppl. 1, 4592–4598.
77. Wang Z, Klipfell E, Bennett BJ, *et al.* (2011) Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature* **472**, 57–63.
78. Baddini Feitoza A, Fernandes Pereira A, Ferreira da Costa N, *et al.* (2009) Conjugated linoleic acid (CLA): effect modulation of body composition and lipid profile. *Nutr Hosp* **24**, 422–428.
79. Fava F, Lovegrove JA, Gitau R, *et al.* (2006) The gut microbiota and lipid metabolism: implications for human health and coronary heart disease. *Curr Med Chem* **13**, 3005–3021.
80. Brighenti F (2007) Dietary fructans and serum triacylglycerols: a meta-analysis of randomized controlled trials. *J Nutr* **137**, 2552S–2556S.
81. Henao-Mejia J, Elinav E, Jin C, *et al.* (2012) Inflammation-mediated dysbiosis regulates progression of NAFLD and obesity. *Nature* **482**, 179–185.
82. Sheth AA & Garcia-Tsao G (2008) Probiotics and liver disease. *J Clin Gastroenterol* **42**, Suppl. 2, S80–S84.
83. Gratz SW, Mykkanen H & El-Nezami HS (2010) Probiotics and gut health: a special focus on liver diseases. *World J Gastroenterol* **16**, 403–410.
84. Malaguarnera M, Vacante M, Antic T, *et al.* (2012) *Bifidobacterium longum* with fructo-oligosaccharides in patients with non alcoholic steatohepatitis. *Dig Dis Sci* **57**, 545–553.
85. Schley PD & Field CJ (2002) The immune-enhancing effects of dietary fibres and prebiotics. *Br J Nutr* **87**, Suppl. 2, S221–S230.
86. Macfarlane GT & Macfarlane S (2011) Fermentation in the human large intestine: its physiologic consequences and the potential contribution of prebiotics. *J Clin Gastroenterol* **45**, S120–S127.
87. Maslowski KM & Mackay CR (2011) Diet, gut microbiota and immune responses. *Nat Immunol* **12**, 5–9.
88. Stoyanova S, Geuns J, Hideg E, *et al.* (2011) The food additives inulin and stevioside counteract oxidative stress. *Int J Food Sci Nutr* **62**, 207–214.
89. Russell WR, Gratz SW, Duncan SH, *et al.* (2011) High-protein, reduced-carbohydrate weight-loss diets promote metabolite profiles likely to be detrimental to colonic health. *Am J Clin Nutr* **93**, 1062–1072.
90. Walton GE, van den Heuvel EG, Kusters MH, *et al.* (2012) A randomised crossover study investigating the effects of galacto-oligosaccharides on the faecal microbiota in men and women over 50 years of age. *Br J Nutr* **107**, 1466–1475.