

Foods provided at shelters during a heavy rain disaster: comparison of weighed food records between different numbers of days

Research Article

Cite this article: Sato H, Sudo N, Nagao-Sato S, and Tsuboyama-Kasaoka N (2025) Foods provided at shelters during a heavy rain disaster: comparison of weighed food records between different numbers of days. *British Journal of Nutrition* **133**: 253–261. doi: [10.1017/S0007114524003088](https://doi.org/10.1017/S0007114524003088)

Received: 23 April 2024

Revised: 18 November 2024

Accepted: 24 November 2024

First published online: 29 November 2024

Keywords:

Administrative dietitian; Evacuation shelter; Natural disaster; Variation; Weighed food record

Abbreviations:

MHLW, Ministry of Health, Labour and Welfare; RV, reference value; WFR, weighed food record

Corresponding author:

Sudo Noriko; Email: sudo.noriko@ocha.ac.jp

Hiroka Sato¹ , Noriko Sudo² , Sayaka Nagao-Sato^{1,3} and Nobuyo Tsuboyama-Kasaoka⁴

¹Department of Food and Nutritional Science, Division of Life Sciences, Graduate School of Humanities and Sciences, Ochanomizu University, 2-1-1 Otsuka, Bunkyo City, Tokyo 112-8610, Japan; ²Natural Science Division, Faculty of Core Research, Ochanomizu University, 2-1-1, Otsuka, Bunkyo City, Tokyo 112-8610, Japan; ³Faculty of Health and Welfare Department of Nutrition, Takasaki University of Health and Welfare, 37-1, Nakaorimachi, Takasaki City, Gunma, 370-0033, Japan and ⁴Laboratory of Disaster Nutrition and Information, National Institutes of Biomedical Innovation, Health and Nutrition, 3-17, Senrioka shinmachi, Settsu City, Osaka 566-0002, Japan

Abstract

Dietitians working at evacuation shelters conduct weighed food records (WFR) for multiple days for dietary assessment. Because the menus in evacuation shelters do not change much from day to day, this study examined whether 1- and 2-d WFR are sufficient for dietary assessment at shelters and identified dietary components that can influence the number of assessment days. Overall, twenty-six WFR were collected from ten shelters in Kumamoto Prefecture, and the amounts of energy; protein; vitamins B₁, B₂ and C and salt were calculated. Correlation analysis and paired sample tests were conducted to examine significant differences between 'one- and two-consecutive- or non-consecutive-day WFR' and 'three-consecutive-day WFR', which were set as the standard in this study. Additionally, the (CV for the categories by meal and dish were calculated to examine the variables that affected the large variations. As a result, 1-d WFR had significant positive correlations with the standard; thus, it could be used for the triage of shelters requiring nutrition assistance as a substitute for 3-d WFR. Two-consecutive-day and non-consecutive-day WFR showed a stronger correlation with the standard compared with the 1-d WFR. For energy and nutrients and dish categories, ready-to-eat foods had larger CV than boxed meals or foods from hot meal services. Whenever the meals included ready-to-eat foods, a two-non-consecutive-day WFR is recommended considering large between-day variations. Salty soup or beverages affected the variation of some nutrients. Our result would help municipalities to consider the number of WFR during emergency.

Japan frequently experiences natural disasters, such as earthquakes and typhoons⁽¹⁾. To prepare for such situations, local government stockpile foods for evacuees⁽²⁾. Within few days following a disaster onset, hot meals or order boxed meals are to be distributed to the evacuees⁽³⁾. Carbohydrate-based foods, such as instant noodles and bread, were previously reported to be in excess⁽⁴⁾, and the consequential increased salt intake in evacuation shelters was associated with the risk of hypertension among the evacuees⁽⁵⁾.

To sustain evacuees' health, the nutritional supply of meals served in evacuation shelters must be estimated. The Ministry of Health, Labor and Welfare (MHLW) released the notice 'Nutritional Reference Values for Feeding at Evacuation Shelters' (reference values (RV))⁽⁶⁾ for the prefectural governments of the affected areas. In general, registered dietitians and dietitians who provide support in evacuation shelters conduct dietary assessments of the provided meals⁽⁷⁾. The survey results are assessed based on the RV and are used as scientific bases to change the contents of the meals provided at the shelters for nutritional improvement⁽²⁾.

Among various dietary assessment methods, the weighed food record (WFR) is considered the gold standard because it provides a high degree of accuracy in assessing food and nutrient intakes, without estimating the amount of food and recalling previous diets^(8,9). This method was conducted in evacuation shelters in the post-disaster period in Japan^(10,11). Unlike dietary assessments in normal settings, WFR in shelters exhibit unique characteristics. First, an individual's nutrient intake is not measured; in evacuation shelters, one meal is measured before distribution⁽¹¹⁾ because it is provided to everyone regardless of their specific needs. Therefore, the nutrient content of one meal is measured regardless of the number of evacuees in the shelter. The aim is mainly to identify the shelters with the greatest need for improvement of meals rather than the individual nutritional assessment. Second, the meal contents following disasters tend to have less variety^(12,13). Consequently, food does not change much from one day to the next. Nishimura⁽¹⁴⁾ found that the monotonous menu of hot meal services was served by self-defence forces. Although local governments conducted WFR for a maximum of 3 d in July 2020 Heavy

Rainfall (original data from the previous study⁽¹¹⁾), the difference may not be large between 3-d and fewer-day WFR.

Although the WFR is the most precise method for dietary assessment, it is time-consuming and could be burdensome for dietitians in emergency settings. In addition, to the best of our knowledge, no study has determined the number of days required for WFR in evacuation shelters. Thus, this study aimed to examine whether 1- and 2-d WFR for dietary assessment at shelters could replace the 3-d WFR conducted in disaster settings. In addition, the meal and dish categories that contributed to the potential variations were examined.

Experimental methods

Study background

Uniquely, Japan has consistently followed its annual National Health and Nutrition Survey since 1946 by applying WFR. Unlike Western countries that use a 24-h dietary recall for their national nutrition surveys^(15–18), the WFR method is widely used by Japanese dietitians who work for local governments, not only for the National Health and Nutrition Survey but also for their periodic prefectural health and nutrition surveys⁽¹⁹⁾.

During emergencies, it takes more time to order supplies from disaster-affected areas. Additionally, the concerned municipalities face difficulties in quickly procuring their basic supplies. Therefore, central government provide a 'push-type relief', including foods and beverages, without waiting for specific request from disaster-stricken municipalities⁽²⁰⁾. Moreover, foods are excessively delivered to specific areas because of media coverage and information spread through social networking services⁽²¹⁾. Consequently, municipal agents cannot presume when and what type of foods they would receive beforehand. To collect promptly dietary information, WFR is also applied during emergencies for an instant dietary support since no additional training is required. Consequently, the use of alternative methods, such as the 24-h dietary recall, solely for emergency situations is not considered efficient.

Study setting

Kumamoto Prefecture, located in the southwest part of Japan, was lashed by heavy rainfalls in July 2020 (from 3 to 4 July 2020)⁽²²⁾. This prefecture mainly has electronics, transportation machinery and production machinery industries⁽²³⁾. In this prefecture, heavy and torrential rains are often due to the warm and humid air flowing in from the Eastern China Sea. Substantial precipitation, particularly during the rainy season, can lead to natural disasters⁽²⁴⁾. In this prefecture, depopulation is progressing, whereas its older population is increasing; approximately 30 % of the residents are > 65 years old at the year of that disaster⁽²⁵⁾. Owing to agricultural, forestry and fishery products, traditional dishes rooted in the local climate have been developed in each area⁽²⁶⁾. As a nutritional issue, 80 % of people in this prefecture consume excessive amounts of salt with lower consumption of vegetables than the recommended intake and the national average. Approximately 70 % of men and 40 % of women aged 40–74 years have high blood pressure or pre-hypertension⁽²⁷⁾.

Following the heavy rainfall, municipalities in Kumamoto Prefecture opened evacuation shelters and provided evacuees with food and water. In twelve evacuation shelters, administrative and/or volunteer dietitians dispatched from dietetic associations were asked to perform dietary assessments of shelter meals 16–20 d after the disaster. This period was classified as phase 2, which is not

acute, and where hot meal services or boxed meals aim to be served. In this period, dietitians need to improve the deficiency of nutrients⁽³⁾. WFR in the evacuation shelters were conducted for the prevention of lifestyle diseases due to unbalanced diets. For nutrition calculation, the prefectural government that had jurisdiction over the affected areas sent WFR sheets collected by the dietitians to the authors' institutions that voluntarily backed them up. Because two of the twelve shelters did not record the weights of food in the WFR, we used data from ten shelters for the analysis. These ten shelters had operated for a median of 100 d (74–119, 25–75 %) since July 2020's Heavy Rainfall^(28–31), so the survey was conducted in the beginning period of the shelter opening. The average number of evacuees in each of the ten shelters during that period was 64 ranging between 15 and 300. Additionally, at least 18 % of the evacuees were vulnerable individuals (the majority had hypertension, followed in order with diabetes, kidney diseases, allergies and others). However, only one generic type of meals was distributed regardless of any specific condition.

WFR was conducted as a public service under the direction of the prefectural government to improve the conditions of the shelters. However, since it was based on administrative decision, no scientific sampling was set in this study. For this study, we asked permission from the prefectural government to use WFR data collected from these shelters. Written approval was given provided that the shelters' names remain anonymous.

Weighed food record

The dietitians involved in this data collection were affiliated with various institutions, including health centres, local governments and dietetic associations within and outside the affected areas. For the WFR method, food record sheets, instructions for recorders and an example of recording were developed and sent to the prefectural nutrition officers who distributed these documents to the dietitians in charge.

During data collection, the dietitians in each shelter were instructed to weigh all food items served as breakfast, lunch and dinner. Basically, each food item was divided into individual ingredients and weighed using a digital scale so that the calculators could estimate the nutrition supply. Each meal was photographed using digital cameras or cell phones and placed on A4 paper, which was used as a scale. Data including product name, manufacturer's name and nutrition fact labelled on ready-to-eat foods or boxed meal packages were recorded as photographs.

Since April 2015, the Food Labelling Act requires to list the amount of energy, protein, fat, carbohydrate and salt on every processed food⁽³²⁾. Ready-to-eat foods and boxed meals are usually prepared by stores or companies, so nutrients other than the mandated ones, especially vitamins on RV, are generally not listed on their nutrition label and on their website. Therefore, the weight records of all the ingredients facilitates in estimation of the amount of all the nutrients on RV that are not shown on the meals nutrition label.

Nutrition calculation

Among the ten shelters, six collected three-consecutive-day WFR and the remaining four collected two-consecutive-day WFR. In total, there were 26-d WFR consisting of seventy-nine meals.

Daily energy and nutrient contents in meals were calculated using Excel Eiyō Plus (in English, Nutrition Plus), which is a widely used add-in software for nutrition calculation, commercially available from Kenpaku sha, a Japanese publisher of academic




Figure 1. Diagram of data used in the correlation analyses and paired sample tests. WFR, weighed food record. *Standard.

books. The software is based on the latest Japanese standard food composition tables published by the Ministry of Education, Culture, Sports, Science and Technology⁽³³⁾. It also provides typical recipes with the weight ratio of ingredients and seasoning for commonly consumed dishes, so simmered vegetables or fried chicken, for example, could be nutritionally calculated using these typical recipes based on their weight record.

RV, nutritional reference values for evacuation shelters set by MHLW, consist of energy, protein and vitamins B₁, B₂ and C⁽⁶⁾. A previous study granted by MHLW revealed the feasibility of the revised RV (draft), which added salt value considering hypertension⁽³⁴⁾. Because the current study examines whether 1- or 2-d WFR could be used for dietary assessment in shelters, we decided to calculate the energy and four nutrients in RV along with salt content.

For nutritional calculation of ready-to-eat foods, such as instant noodles and miso soup, we prioritised the nutritional information published in the manufacturers' nutrition label or through their website. In case of any missing data regarding the amount of nutrients, especially vitamins, we selected similar food items on '18 Food Group (Prepared and processed foods)' in Standard Tables of Food Composition in Japan 2020 (The Eighth Version)⁽³³⁾, followed by typical recipes in 'Excel Eiyō Plus' to fill the amount of these nutrients. Similarly, we referred to the nutrition label or website for nutritional information of boxed meals. Even when dishes could not be found in the Standard Tables of Food Composition, we calculated the nutrients using typical recipes in 'Excel Eiyō Plus'. Foods from hot meal services were also referred to the 18 Food Group in Standard Tables of Food Composition first, followed by typical recipes in 'Excel Eiyō Plus'.

Data analysis

Using three-consecutive-day WFR from six shelters (18-d WFR in total), pairs were formed between 1-d WFR and the average of three-consecutive-day WFR (standard) in each shelter. Correlation coefficients of the total eighteen pairs were then calculated to check whether only a 1-d WFR could identify the shelters with relatively poor diets (Fig. 1(a)). In addition, two-consecutive- and two-non-consecutive-day WFR were compared with the standard to test whether a 2-d WFR had larger correlation coefficients than a 1-d WFR (Fig. 1(b) and (c)). After conducting the Shapiro–Wilk test, Pearson's correlation coefficient was calculated for normally distributed data. Otherwise, Spearman's correlation coefficient was used.

To compare the two-consecutive and two-non-consecutive-day WFR, the average of the three-consecutive-day WFR was also used as a standard (Fig. 1(b) and (c)). When the amounts of energy and

nutrients followed a normal distribution, a paired *t* test was conducted; otherwise, the Wilcoxon signed-rank test was run.

Furthermore, the coefficients of between-day variation of energy or nutrients by meal category were calculated using all WFR from the ten shelters (26-d WFR in total). First, seventy-nine meals from 26-d WFR were grouped between two: ready-to-eat foods and boxed meals or foods from hot meal services. The former consisted of food aids mainly consisting of a carbohydrate-based diet with less food variety, whereas the latter usually included a main and/or a side dish. The CV of energy and nutrients were then calculated by meal category, in addition to the following three categories: staple, main and side dishes to identify the dietary components that contributed to the large variation. The staple dish was defined as the carbohydrate source (cereals), the main dish as the protein source (beans, fish and shellfish, meats and eggs) and the side dish as the vitamin, minerals and dietary fibres source (potatoes and starches, vegetables, fungi and algae) based on the Japanese Food Guide Spinning Top⁽³⁵⁾.

Because ready-to-eat carbohydrate-based foods were reported to be oversupplied⁽⁴⁾, we examined whether the provision quantity affected the energy and nutrients supply within the ready-to-eat foods. In this study, the number of ready-to-eat staple foods was not balanced. Therefore, they were divided ready-to-eat foods-based meals into two groups ('more than one' and 'one or less' staple foods) based on the median, and the Mann–Whitney *U* test was conducted. In addition, some meals were provided with a salty soup or beverage, so those nutritional effects were investigated by grouping 'meals with soup/beverage' or 'others'. In this study, only one type of beverage without foods was served twice. However, because they would not contribute to energy and nutrient supply compared with the meals, they were categorised into 'others'. The Mann–Whitney *U* test was used for their analysis.

All statistical analyses were performed using IBM SPSS Statistics for Windows version 28. The significance level was set at 5 %.

Results

Correlation and comparison of energy and nutrients between weighed food record for different number of days

Energy and nutrients estimated from the 1-d WFR had a strong positive correlation ($r > 0.6$)⁽³⁶⁾ with the average of three-consecutive-day WFR (Table 1). However, stronger correlations in energy and nutrients were obtained with two-consecutive- and non-consecutive-day WFR; most of them had > 0.8 , which is a very strong correlation coefficient⁽³⁶⁾. Overall, the amount of energy in the analysed food items showed a very strong positive

Table 3. Observed daily meal patterns and their frequencies

Daily meal patterns	Breakfast	Lunch	Dinner	Snack
Type 1 (2 d)*	Ready-to-eat foods –*	–*	Boxed meals or foods from hot meal services	–*
Type 2 (3 d)*	Ready-to-eat foods	Ready-to-eat foods	Boxed meals or foods from hot meal services	–*
Type 3 (3 d)*	Ready-to-eat foods	Ready-to-eat foods	Boxed meals or foods from hot meal services	Ready-to-eat foods
Type 4 (14 d)†	Ready-to-eat foods	Boxed meals or foods from hot meal services	Boxed meals or foods from hot meal services	–*
Type 5 (4 d)†	Boxed meal or foods from hot meal service	Boxed meals or foods from hot meal services	Boxed meals or foods from hot meal services	–*

WFR, weighed food record.

*Data from four shelters with two-consecutive-day WFR.

†Data from six shelters with three-consecutive-day WFR.

Table 4. Average amounts and CV of energy and nutrients by meal category

Meal category	Average amount			CV %
Ready-to-eat foods (mainly carbohydrate-based foods) (31 meals)	Energy and nutrients			
	Energy	(kcal)	465	(310–562)**
	Protein	(g)	10.1	(6.9–16.9)**
	Vitamin B ₁	(mg)	0.11	(0.08–0.19)**
	Vitamin B ₂	(mg)	0.14	(0.08–0.19)**
	Vitamin C	(mg)	4.6	(1.9–10.4)**
	Salt	(g)	2.1	(1.4–3.0)**
	Dish category			
	Staple dish	(g)	167	(75–201)**
	Main dish	(g)	0	(0–27.7)**
	Side dish	(g)	0	(0–20.6)**
Boxed meals or foods from hot meal services (48 meals)	Energy and nutrients			
	Energy	(kcal)	683	(146)*
	Protein	(g)	24.5	(6.5)*
	Vitamin B ₁	(mg)	0.27	(0.11)*
	Vitamin B ₂	(mg)	0.32	(0.13)*
	Vitamin C	(mg)	14.4	(10.5–27.1)**
	Salt	(g)	3.2	(1.4)*
	Dish category			
	Staple dish	(g)	213.6	(41.8)*
	Main dish	(g)	90.9	(34.0)*
	Side dish	(g)	40.9	(22.7–64.2)*

*Mean (SD).

**Median (interquartile range).

In six shelters with 3-d WFR, breakfast consisted of carbohydrate-based ready-to-eat foods were served for 14 d (type 4 in Table 3). All three meals (breakfast, lunch and dinner) were a combination of rice and main and/or side dishes for 4 days (type 5 in Table 3).

All meals were divided into ready-to-eat foods (*n* 31) and boxed meals or foods from hot meal services (*n* 48). As shown in Table 4, the average amounts of energy and nutrients per meal were higher in boxed meals or foods from hot meal services than in ready-to-eat foods. In contrast, CV for energy and all nutrients in

Table 5. Association between meals with 'more than one' and 'one or less' ready-to-eat staple foods in energy and nutrients' supply

		More than one*		One or less**		Mann-Whitney <i>U</i> test
		(16 meals)		(15 meals)		<i>P</i>
Energy	(kcal)	603	(248)†	292	(150–389)††	0.001
Protein	(g)	14.6	(5.0)†	7.3	(2.5–8.6)††	0.005
Vitamin B ₁	(mg)	0.14	(0.12–0.22)††	0.09	(0.05–0.11)††	0.010
Vitamin B ₂	(mg)	0.17	(0.07)†	0.09	(0.06–0.16)††	0.149
Vitamin C	(mg)	6.6	(4.3–11.9)††	2.3	(0.0–6.6)††	0.084
Salt	(g)	2.9	(2.1–4.4)††	1.2	(0.4–2.1)††	< 0.001

*Number of carbohydrate-based dishes: four (*n* 1), three (*n* 3) and two (*n* 12).

**Number of carbohydrate-based dishes: one (*n* 11) and none (*n* 4).

†Mean (sd).

††Median (interquartile range).

ready-to-eat foods were larger than those in boxed meals or foods from hot meal services. In both meals, vitamin C had the largest CV. Salt was the second (boxed meals or foods from hot meal services) or third (ready-to-eat foods) nutrient with large CV. In addition, the CV of staple dish was the smallest, followed by the main and side dishes in both meal categories.

Ready-to-eat foods containing carbohydrate-based foods, such as rice balls or bread, were almost always served. However, the number of carbohydrate-based foods supplied was different for each meal. Out of thirty-one meals, sixteen consisted of ready-to-eat foods containing more than one carbohydrate-based food such as a combination of rice and bread. Eleven meals based on ready-to-eat foods contained one carbohydrate-based food, whereas the remaining four contained none. Table 5 shows the results of the Mann-Whitney *U* test of two groups divided by the number of carbohydrate-based foods: 'more than one' and 'one or less'. Significant differences were detected in energy, protein, vitamin B₁ and salt between meals with more than one carbohydrate-based food and meals with one or less those foods.

Table 6 shows the results of the Mann-Whitney *U* test of the effect of the provision of soup or beverage on the supply of energy and nutrients in meals. In this study, soup from instant noodles (57–75 g (without water); *n* 4), instant miso soup (42–54 g (without water); *n* 2) and Japanese or Chinese style soup from hot meal services made from granule soup stock or miso (124–400 g; *n* 6) were provided with meals. Consequently, meals with soup contained significantly higher amounts of vitamin B₁ and salt. Some meals were also served with beverages such as vegetable juice (200–265 g; *n* 6), bottled green tea (500 g; *n* 5), lactic acid bacteria beverage (84 g; *n* 3), milk (206 g; *n* 1) and sports beverage (500 g; *n* 1). Table 6 shows that the amounts of vitamin C significantly increased in meals with beverages, which was about five times as meals without beverages.

Discussion

Recommended number of weighed food record

As shown in Table 1, strong positive correlations were observed between standard (average of three-consecutive-day WFR) and 1-d WFR. According to the report of the Great East Japan Earthquake, dietary surveys must be conducted efficiently, and their results

must be sent promptly to the local government to improve the nutritional status in shelters⁽³⁷⁾. Because no study has revealed the necessary number of days for WFR in emergency settings, this study was the first to show that only 1-d WFR could contribute to the relative evaluation to identify shelters with poorer nutrition supply as a substitute for 3-d WFR. This might serve as a rationale for local governments to promptly assist shelters that needed the most help. Table 1 also reveals that the energy amount had a very strong positive correlation with the standard. Therefore, a highly accurate assessment would be expected for energy supply, a necessary nutritional source for the immediate post-disaster⁽³⁾.

As shown in Table 1, the average of two-consecutive- or two-non-consecutive-day WFR showed a stronger positive correlation with a standard comparing to 1-d WFR. Although 1-d WFR can be used for assessing nutrition supply, 2-d WFR would be preferable for more accurate nutritional assessment if the dietitians in the evacuation shelters are willing to commit. No significant differences were shown in energy and most nutrients in both pairs (Table 2), but previous studies^(38,39) have revealed that non-consecutive-day data were more preferable than data collected on adjacent days because the former considered the changes in meal contents over an extended period. In addition, the dietary surveys for several days might reduce the motivation to continue consecutive-day WFR⁽⁴⁰⁾. Consequently, non-consecutive-day WFR are preferable when local governments ask dietitians to conduct 2-d WFR. If several non-consecutive-day WFR are needed in shelters, the survey period should be within the same phase, such as phase 2 (4 days to 1 month after the disaster) or phase 3 (over 1 month after the disaster)⁽³⁾. These phases focus on dealing with the nutritional issues under conditions where food availability is relatively improved⁽³⁾.

CV of energy and nutrients in meal categories and food groups

We further analysed how meals or food categories affect the number of necessary days for WFR. Compared with boxed meals or foods from hot meals services, meals consisting of ready-to-eat foods provided smaller average amounts of energy and all nutrients with larger CV (Table 4). In addition, ready-to-eat foods based meals with higher number of carbohydrate-based foods showed significantly higher levels of energy and some nutrients (Table 5). According to previous studies, rice balls, bread and biscuits^(4,41,42)

Table 6. Association between 'meals with soup/beverage' and 'others' in energy and nutrients' supply

		Meals with soup*		Others		Mann-Whitney <i>U</i> test
		(12 meals)		(67 meals)		<i>P</i>
Energy	(kcal)	686	(268)†	576	(218)†	0.144
Protein	(g)	23.4	(11.6)†	18.5	(8.4)†	0.148
Vitamin B ₁	(mg)	0.33	(0.19–0.41)††	0.20	(0.14–0.26)††	0.031
Vitamin B ₂	(mg)	0.33	(0.16)†	0.22	(0.16–0.30)††	0.074
Vitamin C	(mg)	10.8	(4.4–19.0)††	11.6	(4.4–19.0)††	0.881
Salt	(g)	5.0	(4.3–4.8)††	2.5	(2.0–3.2)††	< 0.001
		Meals with beverage**		Others		Mann-Whitney <i>U</i> test
		(16 meals)		(63 meals)		<i>P</i>
Energy	(kcal)	612	(138)†	588	(246)†	0.262
Protein	(g)	20.0	(5.2)†	19.2	(10.4–26.4)††	0.323
Vitamin B ₁	(mg)	0.24	(0.11)†	0.20	(0.13–0.30)††	0.520
Vitamin B ₂	(mg)	0.35	(0.11)†	0.21	(0.14–0.27)††	0.068
Vitamin C	(mg)	43.5	(21.8–152.2)††	9.4	(3.4–14.2)††	< 0.001
Salt	(g)	2.7	(0.9)†	2.7	(2.0–3.5)††	0.187

*Meals with soup: instant noodle (57–75 g (without water); *n* 4), instant miso soup (42–54 g (without water); *n* 2) and Japanese or Chinese style soup from hot meal services made from granule soup stock or miso (124–400 g; *n* 6).

**Meals with beverage: vegetable juice (200–265 g; *n* 6), bottled green tea (500 g; *n* 5), lactic acid bacteria beverage (84 g; *n* 3), milk (206 g; *n* 1) and sports beverage (500 g; *n* 1).

†Mean (SD).

††Median (interquartile range).

were often provided in disaster-affected areas. Although boxed meals or foods from hot meal services can increase some food categories⁽⁴³⁾, differences in the number of carbohydrate-based foods in ready-to-eat foods-based meals appear to significantly contribute to CV in terms of energy and nutrients. Because CV of ready-to-eat foods might be larger than those of other meal types, 2-d WFR, preferably non-consecutive-day WFR, should be conducted instead of 1-d WFR considering variations when ready-to-eat foods are mainly served in the surveyed shelters.

Regarding food groups, staple dishes had the smallest CV (Table 4). In Japan where rice is usually consumed in a relatively stable amount, CV of cereals are relatively small^(44,45). Therefore, our results were consistent, although the food environment in emergencies is different from that in normal situations. On the contrary, previous studies^(44,45) on side dishes have shown that fungi and algae require multiple days for estimation: > 2 years within 10 % of their true value because of their large within-individual variations. Vegetables took > 1 month within 10 % of their true value^(44,45). Similar results were observed in this study, making it difficult to assess the nutritional supply (Table 4).

Other factors affecting large CV of energy and nutrients

Soup or beverages with meals can affect CV in terms of energy and nutrients (Table 6). For salt, meals with soup had significantly higher amounts than those without soup; the difference in quantity was approximately twice as much. Previous study has shown that the salt supply in evacuation shelters was high in food aids such as instant noodles⁽⁴⁾. In addition, hypertension risk due to increased salt intake was reported after disasters⁽⁵⁾. Therefore, the salty soup might contribute to the relatively large variations (Table 6). Beverages served with meals had significantly higher levels of vitamin C, approximately five times. The main dietary factors may

be vegetable juice and bottled green tea, which are reported to be important sources of vitamin C⁽¹¹⁾. In general, the addition of vegetables in meals, which are generally vitamin C sources, is difficult because of their scarcity⁽⁴⁶⁾. Therefore, estimating the accurate vitamin C supply might be hard given the limited number of WFR when vitamin C rich beverages are served.

Implementation of nutritional assessment in evacuation shelters

Conducting nutritional surveys in disasters might reduce Japan's vulnerability. The results of this study would provide insights into the nutritional improvement in evacuation shelters. It would also be useful for understanding dietary issues during disasters in countries with very high vulnerability, such as Africa, Asia and Central America. For example, in Puerto Rico, after Hurricane María, a federal distribution centre provided snacks and sweets at a high frequency, whereas fruits, vegetables, proteins and grains accounted for only approximately 10 % of all foods provided⁽⁴⁷⁾. In the aftermath of disasters in Iran, food aid was reported to be inappropriate because it contained high amounts of Na and sugar, with limited amounts of fibre⁽⁴¹⁾. Although such dietary issues have been reported, no evidence shows that other countries have conducted dietary surveys for accurate nutritional assessment in times of disaster. Therefore, this study presents the validity of 1- or 2-d WFR and the important notice for assessment.

Strengths and limitations

This study has several limitations. First, the number of shelters included in this study was relatively small compared with those in studies conducted in normal settings. While previous studies have collected data from thirty-five volunteers⁽³⁹⁾, the present study

focused on assessing shelters rather than individuals, which led to a smaller number of facilities. In addition, the July 2020 Heavy Rainfall affected a few regions, which further limited the number of shelters that could be included. For example, while there were 2417 shelters during the Great East Japan Earthquake in 2011⁽⁴⁸⁾, only a maximum of 212 shelters were available⁽⁴⁹⁾. However, food and nutrition assistance during disasters was strengthened after the Great East Japan Earthquake such as issuing RV⁽⁶⁾ and developing food and nutritional assistance manuals for professionals⁽³⁾. Although the scale of damage was not as large as that of the Great East Japan Earthquake, the July 2020 Heavy Rainfall had the largest number of human casualties following that earthquake⁽⁵⁰⁾ and these new food and nutrition assistance system was applied at that time. Therefore, the WFR in this study would reflect the latest food and nutritional assistance, and the analysis results are useful for future disaster management.

Second, a maximum of three-consecutive-day WFR is available for this study. Although this approach is commonly used in disaster settings because of the limited time available for data collection, it may not provide the same level of accuracy as the 7-d dietary records used in a previous study⁽³⁹⁾. However, the Disaster Relief Act limits the duration of evacuation shelters to a maximum of 7 d in general, which makes it challenging to collect data over a longer period unless a rare catastrophic disaster occurs. Whenever evacuation shelters are open for a longer period in the future, more research-based WFR might be collected to estimate the necessary number of days.

Conclusions

This study showed that even 1-d WFR could substitute the 3-d WFR in identifying shelters in immediate need of nutrition support. If the dietitians agree, two-non-consecutive-day WFR are recommended as it gives better degree of accuracy to assess the nutritional supply. Considering the large CV, 2-d WFR, preferably non-consecutive-day WFR, are recommended especially when the meals included ready-to-eat foods. Salty soup or beverages, which are rich in specific nutrients, can also affect the variation. To demonstrate the robustness and generalisability of these findings, a study, particularly one that spans larger areas should be conducted after large-scale disasters in the future.

Acknowledgements. The authors thank all the investigators and the government of the Kumamoto Prefecture for providing the relevant data for this study.

This study was funded by Grant-in Aid for Scientific Research from the Ministry of Health, Labor and Welfare (grant number 20FA2001) (Principal Investigator: Noriko Sudo). However, the funding body played no role in the design of the study, collection, analysis, interpretation of data and in writing the manuscript.

N. S. was in charge of administration and supervision of this study. N. S. and N. T-K. conceptualized, designed the research (including investigation and methodology), and contributed to funding acquisition. N. S., N. T-K. and S-N-S. contributed to the data collection. H. S. and N. S. conducted data curation. H. S. analyzed data using software and visualized the results. Major contributors to the manuscripts were H. S. and N. S. All authors critically revised and commented on the previous versions of the manuscript. After revisions, the final version of the manuscript was approved by all the authors. N. S. was responsible for submitting the manuscript.

The author(s) declare no conflicts of interest.

Due to the nature of the study, the Kumamoto prefectural government did not allow us to share their data publicly. However, these data are available from the corresponding authors upon reasonable request.

The government of Kumamoto Prefecture conducted this dietary survey as part of shelter management following disasters. Because it is based on the secondary use of data collected by local governments, this study was deemed not applicable for review by the Ethical Review Committee of the Ochanomizu University Research in Humanities and Social Sciences. The authors obtained permission for the secondary use of data from the government of Kumamoto Prefecture (Permission notice: Kenzusui No. 885). Consent to participate does not apply to this study. All methods were conducted in accordance with the relevant guidelines and regulations of the institution and the Declaration of Helsinki.

References

1. Ministry of Foreign Affairs (MOFA) (2021) Disasters and Disaster Prevention in Japan. Chiyoda (JPN). <https://www.mofa.go.jp/policy/disaster/21st/2.html> (accessed February 2024).
2. Sudo N, Shimada I, Tsuboyama-Kasaoka N, et al. (2021) Revising 'Nutritional Reference Values for Feeding at Evacuation Shelters' according to nutrition assistance by public health dietitians based on past major natural disasters in Japan: a quantitative study. *Int J Environ Res Public Health* **18**, 10063.
3. National Institute of Health and Nutrition (NIHN) & The Japan Dietetic Association (JDA) (2011) Manual for Nutrition and Dietary Assistance during Emergencies. Minato (JPN). <https://www.dietician.or.jp/assets/data/learn/marterial/h23evacuation5.pdf> (accessed April 2024) (in Japanese).
4. Tsuboyama-Kasaoka N, Hoshi Y, Onodera K, et al. (2014) What factors were important for dietary improvement in emergency shelters after Great East Japan earthquake? *Asia Pac J Clin Nutr* **23**, 159–166.
5. Hoshida S, Nishizawa M, Okawara Y, et al. (2019) Salt intake and risk of disaster hypertension among evacuees in a shelter after the Great East Japan Earthquake. *Hypertension* **74**, 564–571.
6. Ministry of Health, Labour and Welfare (MHLW) (2011) Nutrition and Food Support in Large-Scale Disasters. Chiyoda (JPN). https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000089299_00005.html (accessed April 2024) (in Japanese).
7. Ito S, Sudo N, Tsuboyama-Kasaoka N, et al. (2015) An analysis of support activities by registered dietitians and dietitians dispatched by the Japan Dietetic Association after the Great East Japan Earthquake. *J Jpn Diet Assoc* **58**, 111–120. (in Japanese).
8. Baranowski T (2012) 24-hour recall and diet record methods. In *Nutritional Epidemiology*, 3rd ed., pp. 49–69 [W Willet, editor]. Oxford (GBR): Oxford University Press.
9. International Dietary Data Expansion (INDDEX) (2024) Weighed Food Record (WFR) Boston (USA). <https://inddex.nutrition.tufts.edu/data4diets/data-source/weighed-food-record-wfr> (accessed April 2024).
10. Takahashi M (2011) Meals in Emergency Shelters, Nutrient Deficiency, Vitamin C Deficiency in All Shelters, The Survey Results in Miyagi Prefecture. The Asahi Shimbun. Chuo (JPN). <https://www.asahi.com/special/10005/TKY201104250400.html> (accessed November 2024) (in Japanese).
11. Takeda T, Sudo N, Tsuboyama-Kasaoka N, et al. (2023) Meal plans for meeting the reference values using food items available in shelters. *BMC Nutr* **9**, 73.
12. AAR Japan (2024) Activity Report, Menus to Make Everyone Happy: Noto Peninsula Earthquake. Shinagawa (JPN). <https://aarjapan.gr.jp/report/13608/> (accessed September 2024) (in Japanese).
13. Moghadam MN, Amiresmailei M, Hassibi M, et al. (2017) Toward a better nutritional aiding in disasters: relying on learned during the Bam Earthquake. *Prehosp Disaster Med* **32**, 382–386.
14. Nishimura K (2011) Diet and nutrition issues in the disaster area-report on the affected areas of Tohoku region Pacific coast earthquake (Kesennuma City, Miyagi). *J Jpn Diabetes Soc* **54**, 724–726. (in Japanese).
15. Centers for Disease Control and Prevention (2024) Measuring Guides for the Dietary Recall Interview. Atlanta (USA). https://www.cdc.gov/nchs/nhanes/measuring_guides_dri/measuringguides.htm (accessed September 2024).

16. Mitsopoulou AV, Magriplis E, Michas G, *et al.* (2021) Micronutrient dietary intakes and their food sources in adults: the Hellenic National Nutrition and Health Survey (HNNHS). *J Hum Nutr Diet* **34**, 616–628.
17. Kaartinen N, Tapanainen H, Reinivuo H, *et al.* (2020) The Finnish National Dietary Survey in adults and elderly (FinDiet 2017). EFSA Support 17, EN-1914.
18. Max Rubner-Institut (MRI) (2024) The German Nutritional Nutrition Survey II. Karlsruhe (DEU). <https://www.mri.bund.de/en/institutes/nutritional-behaviour/research-projects/nvsii/> (accessed September 2024).
19. Kubo A, Kuno K, Maruyama K, *et al.* (2022) Impact of the COVID-19 pandemic on national and prefectural health and nutrition examination surveys: a report by the Monitoring Report Committee of the Japanese Society of Public Health. *Jpn J Public Health* **69**, 586–594. (in Japanese).
20. Ministry of Agriculture, Forestry and Fisheries (MAFF) (2019) Evacuation Shelters and Relief Supplies. Chiyoda (JPN). https://www.maff.go.jp/j/pr/aff/1909/spe1_04.html (accessed October 2024) (in Japanese).
21. Japanese Voluntary Organizations Active in Disaster (2024) Manual for Food and Nutrition Assistance at the Time of Disasters. Chiyoda (JPN). https://jvoad.jp/wp-content/uploads/2024/05/tabepro_reference_word_202405.pdf (accessed October 2024) (in Japanese).
22. Kumamoto nichinichi shimbun (2021) Kumamoto Heavy Rainfall 1 Year. Kumamoto (JPN). <https://kumanichi.com/theme/gouu2020/year> (accessed April 2024) (in Japanese).
23. Kumamoto Prefecture (2017) Statistics a la carte. Kumamoto (JPN). <https://www.pref.kumamoto.jp/uploaded/attachment/13888.pdf> (accessed September 2024) (in Japanese).
24. Japan Meteorological Agency (2024) The Climate in Kumamoto Prefecture. Minato (JPN). <https://www.data.jma.go.jp/kumamoto/shosai/climate.html> (accessed November 2024) (in Japanese).
25. Senior Citizens Support Division, Department of Health and Social Services & Kumamoto Prefectural Government (2022) Kumamoto Prefecture is Experiencing an Aging Population. Kumamoto (JPN). <https://www.pref.kumamoto.jp/soshiki/32/156988.html> (accessed April 2024) (in Japanese).
26. Kumamoto Prefecture (2024) The 4th Health and Dietary Habits in Kumamoto Prefecture, Prefectural Plan for the Promotion of Shokuiku. Kumamoto (JPN). <https://www.pref.kumamoto.jp/uploaded/attachment/245064.pdf> (accessed September 2024) (in Japanese).
27. Kumamoto Prefecture (2024) The Summary of 'The 2022 National Health and Nutrition Survey' Focusing on Kumamoto Prefecture. Kumamoto (JPN). <https://www.pref.kumamoto.jp/uploaded/attachment/243331.pdf> (accessed September 2024) (in Japanese).
28. Ashikita Town (2021) Ashikita Town Recovery and Reconstruction Plan. Ashikita (JPN). <https://www.pref.kumamoto.jp/uploaded/attachment/136996.pdf> (accessed September 2024) (in Japanese).
29. Kumamoto Rosai Hospital (2020) The Report of Disaster Rehabilitation or Reconstruction Rehabilitation in July 2020 Heavy Rainfall. Yatsushiro (JPN). <https://kumamotoh.johas.go.jp/info/docs/bd4c3683ca1bf8145fa67b0abe8475fec2ec4afe.pdf> (accessed September 2024) (in Japanese).
30. Murakami S (2020) Heavy Rainfall in Kumamoto Prefecture, the Closure of Largest-Scale Evacuation Shelters which had 836 Evacuees at One Time. The Asahi Shimbun. Chuo (JPN). <https://www.asahi.com/articles/ASNDX6SQNDXTIPE00H.html> (accessed September 2024) (in Japanese).
31. Kuma Village (2021) The Summary of Disaster Management. Kuma (JPN). <https://www.kumamura.com/gyousei/wp-content/uploads/2021/05/fe772831c2b0d6aa4b0d87fa12dc601.pdf> (accessed September 2024) (in Japanese).
32. Consumer Affairs Agency (CAA) (2020) Nutrition Labelling. Chiyoda (JPN). https://www.caa.go.jp/policies/policy/food_labeling/nutrient_decleration/ (accessed November 2024).
33. Ministry of Education, Culture, Sports, Science, and Technology (MEXT) (2020) Standard Tables of Food Composition in Japan-2020. 8th rev. ed. (dataset). Chiyoda (JPN). https://www.mext.go.jp/a_menu/syokuhinseibun/mext_01110.html (accessed April 2024) (in Japanese).
34. Sato H, Sudo N, Takeda T, *et al.* (2024) Revision of 'Nutritional Reference Values for Feeding at Evacuation Shelters' and model menus: a qualitative study. *J Am Nutr Assoc* **43**, 157–166.
35. Yoshiike N, Hayashi F, Takemi Y, *et al.* (2007) A new food guide in Japan: the Japanese food guide spinning top. *Nutr Rev* **65**, 149–154.
36. The BMJ (2024) Correlation and Regression. London (GBR). <https://www.bmj.com/about-bmj/resources-readers/publications/statistics-square-one/11-correlation-and-regression> (accessed April 2024).
37. Ministry of Health, Labour and Welfare (MHLW) (2013) The Response Situation (Food and Nutritional Assistance) After the Great East Japan Earthquake. Chiyoda (JPN). <https://www.mhlw.go.jp/content/10900000/000637201.pdf> (accessed 19 February 2024) (in Japanese).
38. Tarasuk V & Beaton GH (1992) Statistical estimation of dietary parameters: implications of patterns in within-subject variation—a case study of sampling strategies. *Am J Clin* **55**, 22–27.
39. Murakami Y, Oshikata R, Miyamoto N, *et al.* (2010) Validity of 1-, 2-, 3- and 4-day dietary records in comparison with 7-day weighed dietary records. *HEP* **37**, 405–413. (in Japanese).
40. Gersovitz M, Madden JP & Smiciklas-Wright H (1978) Validity of the 24-h dietary recall and seven-day record for group comparisons. *J Am Diet Assoc* **73**, 48–55.
41. Ainehvand S, Raeissi P, Ravaghi H, *et al.* (2019) Natural disasters and challenges toward achieving food security response in Iran. *J Educ Health Promot* **8**, 51.
42. Sasaki Y (2012) Food assistance and nutrition management as the administration dietitian in the shelter of the 2011 Japan Earthquake and disaster. *Bull Sendai Shirayuri Womens Coll* **16**, 103–118. (in Japanese).
43. Mihara M, Harada M, Oka J, *et al.* (2019) The effect of lunch box provision and mass feeding on energy and nutrient supply at emergency shelters after the Great East Japan Earthquake. *Jpn J Public Health* **66**, 629–637. (in Japanese).
44. Egami I, Wakai K, Kaitoh K, *et al.* (1999) Intra- and inter-individual variations in diets of the middle-aged and the elderly. *Jpn J Public Health* **46**, 828–837. (in Japanese).
45. Ogawa K, Tsubono Y, Nishino Y, *et al.* (1999) Inter- and intra-individual variation of food and nutrient consumption in a rural Japanese population. *Eur J Clin Nutr* **53**, 781–785.
46. Sudo N, Sawaguchi M & Yoshiike N (2010) Changes in food intakes and required nutrients under stress: to support disaster victims with food assistance. *J Jpn Diet Assoc* **53**, 349–355. (in Japanese).
47. Colón-Ramos U, Roess AA, Robien K, *et al.* (2019) Foods distributed during Federal Disaster Relief Response in Puerto Rico after Hurricane María did not fully meet Federal Nutrition Recommendations. *J Acad Nutr Diet* **119**, 1903–1915.
48. Reconstruction Agency (2011) Transition of Refugees and Shelters (The Comparison with Great East Japan Earthquake, Great Hanshin-Awaji Earthquake, and Chuetsu Earthquake). Chiyoda (JPN). 12 October 2011. <https://www.reconstruction.go.jp/topics/main-cat2/sub-cat2-6/index.html> (accessed April 2024) (in Japanese).
49. Kumamoto Prefecture (Health and Social Services Policy Division) (2020) Reiwa 2 (2020), Conditions of Support for Disaster Victims during Heavy Rain in July the 2nd Year of Reiwa. Kumamoto (JPN). 25 December 2020. <https://www.pref.kumamoto.jp/uploaded/attachment/125989.pdf> (accessed April 2024) (in Japanese).
50. Ministry of Internal Affairs and Communications (MIC) (2021) Disaster prevention/mitigation and ICT. In *White paper Information and Communications in Japan, Part 1*, pp. 241–271. Chiyoda (JPN). <https://www.soumu.go.jp/johotsusintokei/whitepaper/ja/r03/pdf/index.html> (accessed November 2024) (in Japanese).