

BEST POSSIBLE COMPACTNESS RESULTS OF LIONS–PEETRE TYPE

FERNANDO COBOS¹, MICHAEL CWIKEL² AND PEDRO MATOS^{1,3}

¹*Departamento de Análisis Matemático, Facultad de Matemáticas,
Universidad Complutense de Madrid, 28040 Madrid, Spain
(cobos@eucmax.sim.ucm.es)*

²*Department of Mathematics, Technion—Israel Institute of Technology,
Haifa 32000, Israel (mcwikel@math.technion.ac.il)*

³*ESTG, Instituto Politécnico de Leiria, 2400 Leiria,
Portugal (matos@estg.iplei.pt)*

(Received 12 November 1998)

Abstract If $T : A_0 \rightarrow B$ boundedly and $T : A_1 \rightarrow B$ compactly, then a result of Lions–Peetre shows that $T : A \rightarrow B$ compactly for a certain class of spaces A which are intermediate with respect to A_0 and A_1 . We investigate to what extent such results can hold for *arbitrary* intermediate spaces A . The ‘dual’ case of an operator S such that $S : X \rightarrow Y_0$ boundedly and $S : X \rightarrow Y_1$ compactly, is also considered, as well as similar questions for other closed operator ideals.

Keywords: interpolation of compact operators; rank-one interpolation spaces;
general intermediate spaces; operator ideals

AMS 2000 *Mathematics subject classification:* Primary 46B70; 47D50

1. Introduction

This paper deals with compactness results for interpolated operators in the style of those established by Lions and Peetre [14] in 1964 (see also the lecture by Gagliardo [10]). Lions and Peetre deal with two situations. One of these is the case of a linear operator T which is a bounded map of both of the spaces A_0 and A_1 of a Banach couple $\bar{A} = (A_0, A_1)$ into the same Banach space B with the additional assumption that $T : A_1 \rightarrow B$ is compact. The other is the ‘dual’ case of an operator T which maps a given Banach space A boundedly into both of the spaces of a Banach couple $\bar{B} = (B_0, B_1)$, and here it is always assumed that $T : A \rightarrow B_1$ is compact. In the first case, Lions and Peetre show that if A is a space of the class $\mathcal{C}_K(\theta, \bar{A})$ for some $\theta \in (0, 1)$, then $T : A \rightarrow B$ is also compact. In the ‘dual’ case, they prove that $T : A \rightarrow B$ is compact for each space B of the class $\mathcal{C}_J(\theta, \bar{B})$. (We shall recall the meaning of the notation used here in §2.)

Compactness results of Lions–Peetre type have interesting applications to function spaces (for example, to Sobolev spaces). Several authors have also investigated them

from other points of view. For example, a quantitative version of Lions–Peetre results in terms of entropy numbers can be found in the book by Pietsch [17, Propositions 12.1.11 and 12.1.12]. Similar results in terms of the measure of non-compactness were established by Edmunds and Teixeira [20]. More abstract versions that work for injective surjective closed operator ideals are due to Heinrich [12, Propositions 1.6 and 1.7].

Lions–Peetre type compactness results are also important tools for establishing a number of other compactness theorems in interpolation theory (see, for example, the papers by Persson [16], Cobos and Peetre [7], and Cobos, Kühn and Schonbek [5]).

Just as in [14], also in the other corresponding results mentioned above (see [17], [20] and [12]), the spaces A and B are required to satisfy $A \in \mathcal{C}_K(\theta, \bar{A})$ or $B \in \mathcal{C}_J(\theta, \bar{B})$. These are, in fact, rather strong restrictions on A and B . It has been pointed out by Mastysłó (see [15, Theorem 1]) that the Lions–Peetre results hold under weaker conditions than these, i.e. the compactness of $T : A \rightarrow B$ can also be proved when A or B , respectively, belong to other classes which generalize $\mathcal{C}_K(\theta, \bar{A})$ or $\mathcal{C}_J(\theta, \bar{B})$. The sufficient conditions formulated in [15] amount to requiring either that A is not ‘too close’ to A_0 , or that B is not ‘too close’ to B_0 , in some appropriate sense. More precisely, they can be expressed as

$$\limsup_{t \rightarrow 0} \left\{ \frac{K(t, a; \bar{A})}{\|a\|_A} : 0 \neq a \in A \right\} = 0 \quad (1.1)$$

and

$$\liminf_{t \rightarrow \infty} \left\{ \frac{J(t, b; \bar{B})}{\|b\|_B} : 0 \neq b \in B_0 \cap B_1 \right\} = \infty, \quad (1.2)$$

respectively.

There are other very simple conditions that are sufficient for $T : A \rightarrow B$ to be compact. If we also have compactness at the other ‘endpoint’, i.e. if

$$T : A_0 \rightarrow B \text{ is compact} \quad (1.3)$$

in the first case, or, analogously,

$$T : A \rightarrow B_0 \text{ is compact} \quad (1.4)$$

in the dual case, then it follows immediately that $T : A_0 + A_1 \rightarrow B$ is compact, or, in the dual case, that $T : A \rightarrow B_0 \cap B_1$ is compact. This in turn means that $T : A \rightarrow B$ is compact, in the first case, for *all* interpolation spaces A with respect to \bar{A} , and in the second case for *all* interpolation spaces B with respect to \bar{B} .

Remark 1.1. In fact, to obtain the compactness of $T : A \rightarrow B$ when T is compact at both ‘endpoints’, it is not even necessary to require A or B , respectively, to be an interpolation space. A weaker condition, namely $A \hookrightarrow A_0 + A_1$ or $B_0 \cap B_1 \hookrightarrow B$, is all that is needed.

In this paper we obtain a quantitative version of the Lions–Peetre Lemma and Theorem 1 of [15], in terms of the measure of non-compactness. We also investigate whether

it is possible to obtain the results of Lions and Peetre without invoking either of the sufficient conditions discussed above, i.e. what can be said if A is merely an interpolation space with respect to \bar{A} in the first context, or B is merely an interpolation space with respect to \bar{B} in the dual setting? As we show in § 3, these minimal assumptions lead to the conclusion that either $T : A \rightarrow B$ is compact, or a certain relation holds between A and A_0 or, respectively, between B and B_0 . Furthermore, under mild additional conditions on the couple \bar{A} or \bar{B} , respectively, the compactness of $T : A \rightarrow B$ turns out to be *equivalent* to requiring one or both of the above sufficient conditions (1.1) and (1.3) (or, respectively, (1.2) and (1.4)) to hold. This shows that in some sense these conditions are optimal.

Note that the preceding results are formulated for specific choices of the range space or, respectively, the domain space. But we also show another aspect of the optimality of conditions (1.1) and (1.2) in the context where compactness is required to hold for all possible range spaces or domain spaces, respectively. We prove (again under mild additional assumptions) that, for given A_0 , A_1 and A , condition (1.1) is necessary and sufficient for $T : A \rightarrow B$ to be compact for *all* Banach spaces B and *all* operators T that map A_0 to B boundedly and A_1 to B compactly. We obtain a corresponding dual result about the optimality of condition (1.2) without requiring any additional assumptions.

We also show, in § 4, that this sort of behaviour is not an exclusive property of compact operators: similar results hold for operators belonging to any other injective surjective closed operator ideal.

The conditions $A \in \mathcal{C}_K(\theta, \bar{A})$ and $B \in \mathcal{C}_J(\theta, \bar{B})$ do not in fact stipulate that A is an interpolation space with respect to \bar{A} nor that B is an interpolation space with respect to \bar{B} . Furthermore, Remark 1.1 provides a further hint that it is not entirely natural to require A or B to be interpolation spaces in the context of the questions being considered here. It turns out, in fact, that most of our results apply when A and B are in the larger class of *interpolation spaces with respect to rank-one operators*, or, in some cases, when they are merely intermediate spaces.

2. Notation and preliminaries

We recall several standard notions from interpolation theory (cf. [1], [2] and [4]). Let $\bar{A} = (A_0, A_1)$ be a *Banach couple*, that is, two Banach spaces A_j , $j = 0, 1$, which are continuously embedded in some Hausdorff topological vector space. For each $t > 0$ we put

$$J(t, a) = J(t, a; \bar{A}) = \max\{\|a\|_{A_0}, t\|a\|_{A_1}\}, \quad \text{for each } a \in A_0 \cap A_1,$$

and

$$\begin{aligned} K(t, a) &= K(t, a; \bar{A}) \\ &= \inf\{\|a_0\|_{A_0} + t\|a_1\|_{A_1} : a = a_0 + a_1, a_j \in A_j\}, \quad \text{for each } a \in A_0 + A_1. \end{aligned}$$

Then $\{K(t, \cdot)\}_{t>0}$ is a family of norms on $A_0 + A_1$, any two of which are equivalent. The family $\{J(t, \cdot)\}_{t>0}$ has similar properties on $A_0 \cap A_1$.

Let $\bar{B} = (B_0, B_1)$ be a second Banach couple. Then the notation $T : \bar{A} \rightarrow \bar{B}$ means that T is a linear operator from $A_0 + A_1$ into $B_0 + B_1$, whose restriction to A_j defines a bounded operator from A_j into B_j for $j = 0$ and $j = 1$. We denote the space of all such operators by $\mathcal{L}(\bar{A}, \bar{B})$, just as $\mathcal{L}(A, B)$ denotes, as usual, the space of all bounded operators mapping a Banach space A into a Banach space B .

For each $T \in \mathcal{L}(\bar{A}, \bar{B})$ we introduce the norm

$$\|T\|_{\bar{A}, \bar{B}} := \max\{\|T\|_{A_0, B_0}, \|T\|_{A_1, B_1}\}.$$

If one of the couples \bar{A} or \bar{B} reduces to a single Banach space, i.e. if $A_0 = A_1 = A$ or if $B_0 = B_1 = B$, then we write $T : A \rightarrow \bar{B}$ and $T \in \mathcal{L}(A, \bar{B})$ and denote the norm by $\|T\|_{A, \bar{B}}$ or, respectively, $T : \bar{A} \rightarrow B$ and $T \in \mathcal{L}(\bar{A}, B)$ with norm $\|T\|_{\bar{A}, B}$.

A Banach space A is said to be an *intermediate space* with respect to \bar{A} if

$$A_0 \cap A_1 \hookrightarrow A \hookrightarrow A_0 + A_1,$$

where, as usual, the notation \hookrightarrow means continuous inclusion. To each intermediate space A it is possible to associate two other intermediate spaces. The first of these, denoted by A° and termed the *clintersect* of A , is the closed subspace of A generated by $A_0 \cap A_1$. The second, denoted by A^\sim and termed the *Gagliardo completion* of A , consists of all those $a \in A_0 + A_1$ for which there exists a sequence $\{a_n\}_{n \in \mathbb{N}}$ in some bounded subset of A which converges to a in $A_0 + A_1$. It is normed by an appropriate infimum over all such sequences, i.e.

$$\|a\|_{A^\sim} = \inf_{\{a_n\}} \sup_{n \in \mathbb{N}} \{\|a_n\|_A\}$$

(see [10], [2] or [1]).

An intermediate space A is an *interpolation space* if, given any $T : \bar{A} \rightarrow \bar{A}$, the restriction of T to A defines a bounded operator from A into itself. It is a classical result of Aronszajn and Gagliardo (see [2, Theorem 2.4.2, p. 28]) that for each such space A there exists a constant $C = C(A, \bar{A})$ such that

$$\|T\|_{A, A} \leq C \|T\|_{\bar{A}, \bar{A}} \tag{2.1}$$

for all operators $T : \bar{A} \rightarrow \bar{A}$.

The most familiar examples of interpolation spaces with respect to a given couple $\bar{A} = (A_0, A_1)$ are the real interpolation spaces $(A_0, A_1)_{\theta, q}$ and the complex interpolation spaces $[A_0, A_1]_\theta$, which are defined for each $\theta \in (0, 1)$ and $q \in [1, \infty]$. Furthermore, $(A_0, A_1)_{\theta, q}$ and $[A_0, A_1]_\theta$ are examples of spaces A that are in both of the classes $\mathcal{C}_J(\theta, \bar{A})$ and $\mathcal{C}_K(\theta, \bar{A})$. In other words, for some constant C and for all $t > 0$ they satisfy each of the two conditions

$$\|a\|_A \leq C t^{-\theta} J(t, a), \quad \text{for all } a \in A_0 \cap A_1 \tag{2.2}$$

and

$$K(t, a) \leq C t^\theta \|a\|_A, \quad \text{for all } a \in A. \tag{2.3}$$

Condition (2.2) is equivalent to the continuous inclusion $(A_0, A_1)_{\theta,1} \hookrightarrow A$ and (2.3) is equivalent to $A \hookrightarrow (A_0, A_1)_{\theta,\infty}$ (see [2, § 3.5]). However, there are also other interpolation spaces that do not satisfy one or both of these properties for any value of $\theta \in (0, 1)$. Obviously, these include the ‘endpoint’ spaces A_0 and A_1 themselves. There are also two other examples of such interpolation spaces which will play special roles in our results here. The first of these is A_0^\wedge , the Gagliardo completion of A_0 . It is easy to show (see, for example, [1, Theorem 5.1.4]) that

$$\|a\|_{A_0^\wedge} = \lim_{t \rightarrow \infty} K(t, a; \bar{A}).$$

(Note that, analogously, the norm of the Gagliardo completion of A_1 satisfies $\|a\|_{A_1^\wedge} = \lim_{t \rightarrow 0} (1/t)K(t, a; \bar{A})$.)

Our second example of such an interpolation space is A_0° , the clintersect of A_0 .

As mentioned in §1, it turns out to be natural for our purposes here to work with another class of intermediate spaces that includes the interpolation spaces as a subclass. This class is defined with the help of rank-one operators $T : \bar{A} \rightarrow \bar{A}$ of the form $T = f \otimes a$, i.e. $Tx = f(x)a$, where a is a fixed element of $A_0 \cap A_1$ and f is a fixed bounded linear functional on $A_0 + A_1$. Obviously, each such T is bounded on A_0 and A_1 and on each intermediate space A with respect to \bar{A} . But its norms on each of these spaces depend on the particular choices of a and f . We shall say that an intermediate space A with respect to the couple \bar{A} is a *rank-one interpolation space* if, for some constant C depending only on A and \bar{A} , it satisfies (2.1) for *all* T of the special form $T = f \otimes a$.

Rank-one interpolation spaces have been used in a number of papers, [9] and [19] for example, where they are referred to as ‘partly interpolation spaces’.

We shall need to use two functions which in some sense measure the ‘position’ of a given intermediate space within a Banach couple. These are variants of functions which have been introduced and studied by Dmitriev and by Pustylnik (see [9], [18, pp. 333–334] and [19, p. 307]). They are defined as follows.

Definition 2.1. Let $\bar{A} = (A_0, A_1)$ be a Banach couple and let A be an intermediate space with respect to \bar{A} . For each $t > 0$ set

$$\psi(t) = \psi(t, A, \bar{A}) := \sup\{K(t, a) : \|a\|_A = 1\}$$

and

$$\rho(t) = \rho(t, A, \bar{A}) := \inf\{J(t, a) : a \in A_0 \cap A_1, \|a\|_A = 1\}.$$

Clearly, $\psi(t)$ and $\rho(t)$ are strictly positive, and they are also both non-decreasing, while $\psi(t)/t$ and $\rho(t)/t$ are non-increasing. We note that these functions are related to the ‘embedding functions’ $C(\alpha, \beta)$ and $D(\alpha, \beta)$ of [19, p. 307] by

$$C(\alpha, \beta) = \frac{1}{(1/\alpha)\rho(\alpha/\beta)} \quad \text{and} \quad D(\alpha, \beta) = \frac{1}{(1/\alpha)\psi(\alpha/\beta)}. \tag{2.4}$$

We also note that the conditions (1.1) and (1.2) stated in the introduction correspond to $\lim_{t \rightarrow 0} \psi(t, A, \bar{A}) = 0$ and $\lim_{t \rightarrow \infty} \rho(t, B, \bar{B}) = \infty$, respectively.

We refer to [19] for some examples indicating the behaviour of the functions C and D in specific cases.

It is possible to characterize rank-one interpolation spaces A by a simple condition on the functions $\psi(t, A, \bar{A})$ and $\rho(t, A, \bar{A})$. This was done by Dmitriev [9] under some additional hypotheses and, in general, by Pustylnik [19]. The following theorem is almost the same as Theorem 1 in [19, p. 309].

Theorem 2.2. *Let $\bar{A} = (A_0, A_1)$ be a Banach couple and let A be an intermediate space with respect to \bar{A} . Then A is a rank-one interpolation space if and only if there exists a constant C depending only on A and \bar{A} such that*

$$\psi(t, A, \bar{A}) \leq C\rho(t, A, \bar{A}), \quad \text{for all } t > 0. \quad (2.5)$$

Proof. In view of (2.4), the proof is a very slight and obvious modification of the proof given in [19]. \square

3. Compactness results

As pointed out by Mastyło, the proof of the generalized Lions–Peetre compactness Lemma, assuming the weaker conditions (1.1) and (1.2) instead of $A \in \mathcal{C}_K(\theta, \bar{A})$ and $B \in \mathcal{C}_J(\theta, \bar{B})$, is essentially the same as the proof of the original lemma (cf. [2, pp. 56–57]).

Here, however, we shall adopt a quantitative approach. The following two theorems describe interpolation properties of the measure of non-compactness. They can be applied in fact to any intermediate space and imply the generalized compactness lemma. Their estimates also point the way to results to be presented in § 4.

Let us recall that the (ball) measure of non-compactness $\beta(T) = \beta(T_{A,B})$ of an operator $T \in \mathcal{L}(A, B)$ is defined to be the infimum of the set of all numbers $\sigma > 0$ for which there exists a finite subset $B(\sigma) \subset B$ such that

$$T(U_A) \subset \bigcup_{b \in B(\sigma)} (b + \sigma U_B).$$

Here, U_A (respectively, U_B) denotes the closed unit ball of A (respectively, B). Of course, T is compact if and only if $\beta(T) = 0$.

Theorem 3.1. *Let $\bar{A} = (A_0, A_1)$ be a Banach couple, let A be an intermediate space with respect to \bar{A} and let B be another Banach space. Then, for each $T \in \mathcal{L}(\bar{A}, B)$:*

(a) *if $\beta(T_{A_0,B}) = 0$, then*

$$\beta(T_{A,B}) \leq \beta(T_{A_1,B}) \cdot \lim_{t \rightarrow \infty} \frac{\psi(t, A, \bar{A})}{t};$$

(b) *if $\beta(T_{A_1,B}) = 0$, then*

$$\beta(T_{A,B}) \leq \beta(T_{A_0,B}) \cdot \lim_{t \rightarrow 0} \psi(t, A, \bar{A});$$

and

(c) if $\beta(T_{A_i,B}) > 0$ for $i = 0, 1$, then

$$\beta(T_{A,B}) \leq 2\beta(T_{A_0,B})\psi\left(\frac{\beta(T_{A_1,B})}{\beta(T_{A_0,B})}, A, \bar{A}\right).$$

Proof. For each $\epsilon > 0$ and for $i = 0, 1$ there exists a finite set $B(\epsilon, i) \subset B$ such that

$$T(U_{A_i}) \subset \bigcup_{b \in B(\epsilon, i)} (b + (\epsilon + \beta(T_{A_i,B}))U_B). \tag{3.1}$$

Now, consider an arbitrary element $a \in U_A$ and fixed positive numbers t and δ . Since $K(t, a) < \delta + \psi(t)$, there exists a decomposition $a = a_0 + a_1$ with $a_i \in A_i$ for $i = 0, 1$ such that $\|a_0\|_{A_0} + t\|a_1\|_{A_1} < \delta + \psi(t)$. Thus,

$$a_i \in t^{-i}(\delta + \psi(t))U_{A_i}, \quad \text{for } i = 0, 1.$$

Consequently, by (3.1), for some $b_i \in B(\epsilon, i)$, we have

$$Ta_i \in t^{-i}(\delta + \psi(t))b_i + t^{-i}(\delta + \psi(t))(\epsilon + \beta(T_{A_i,B}))U_B, \quad \text{for } i = 0, 1,$$

which implies that

$$Ta \in (\delta + \psi(t))\left(b_0 + \frac{1}{t}b_1\right) + (\delta + \psi(t))\left(\epsilon + \beta(T_{A_0,B}) + \frac{1}{t}(\epsilon + \beta(T_{A_1,B}))\right)U_B.$$

Since the subset of B consisting of all elements of the form $(\delta + \psi(t))(b_0 + (1/t)b_1)$ for some $b_0 \in B(\epsilon, 0)$ and $b_1 \in B(\epsilon, 1)$ is of course finite, we have shown that

$$\beta(T_{A,B}) \leq (\delta + \psi(t))\left(\epsilon + \beta(T_{A_0,B}) + \frac{1}{t}(\epsilon + \beta(T_{A_1,B}))\right).$$

This in turn implies, since ϵ and δ can both be chosen arbitrarily small, that

$$\beta(T_{A,B}) \leq \psi(t)\left(\beta(T_{A_0,B}) + \frac{1}{t}\beta(T_{A_1,B})\right), \quad \text{for all } t > 0. \tag{3.2}$$

We can now immediately obtain cases (a) and (b) of the theorem by substituting $\beta(T_{A_i,B}) = 0$ in (3.2) for $i = 0$ or 1 , and using the facts that $\psi(t)$ is non-decreasing and $\psi(t)/t$ is non-increasing. Case (c) also follows from (3.2) by simply substituting $t = \beta(T_{A_1,B})/\beta(T_{A_0,B})$. \square

Now we turn to the analogous theorem for the ‘dual’ situation.

Theorem 3.2. Let $\bar{B} = (B_0, B_1)$ be a Banach couple, let B be an intermediate space with respect to \bar{B} and let A be another Banach space. Then, for each operator $T \in \mathcal{L}(A, \bar{B})$:

(a) if $\beta(T_{A,B_0}) = 0$, then

$$\beta(T_{A,B}) \leq 2\beta(T_{A,B_1}) \cdot \lim_{t \rightarrow 0} \frac{t}{\rho(t, B, \bar{B})};$$

(b) if $\beta(T_{A,B_1}) = 0$, then

$$\beta(T_{A,B}) \leq 2\beta(T_{A,B_0}) \cdot \lim_{t \rightarrow \infty} \frac{1}{\rho(t, B, \bar{B})};$$

and

(c) if $\beta(T_{A,B_i}) > 0$ for $i = 0, 1$, then

$$\beta(T_{A,B}) \leq \frac{2\beta(T_{A,B_0})}{\rho(\beta(T_{A,B_0})/\beta(T_{A,B_1}), B, \bar{B})}.$$

Proof. For each fixed $\epsilon > 0$ there exist finite subsets $B(\epsilon, 0)$ of B_0 and $B(\epsilon, 1)$ of B_1 such that

$$T(U_A) \subset \bigcup_{b \in B(\epsilon, i)} (b + (\epsilon + \beta(T_{A,B_i}))U_{B_i}), \quad \text{for } i = 0, 1. \quad (3.3)$$

Let us now construct a new set $\tilde{B}(\epsilon)$ of elements of $B_0 \cap B_1$ in the following way: For each choice of $b_0 \in B(\epsilon, 0)$ and $b_1 \in B(\epsilon, 1)$, if the set $E(b_0, b_1) = (b_0 + (\epsilon + \beta(T_{A,B_0}))U_{B_0}) \cap (b_1 + (\epsilon + \beta(T_{A,B_1}))U_{B_1})$ is non-empty, choose exactly one element $w(b_0, b_1)$ from this set, i.e. we set

$$\tilde{B}(\epsilon) = \{w(b_0, b_1) : b_0 \in B(\epsilon, 0), b_1 \in B(\epsilon, 1), E(b_0, b_1) \neq \emptyset\}.$$

Clearly, $\tilde{B}(\epsilon)$ is a non-empty finite set.

Given an arbitrary fixed element $a \in U_A$, it follows from (3.3) that there exist $b_0 \in B(\epsilon, 0)$ and $b_1 \in B(\epsilon, 1)$ such that $Ta \in (b_0 + (\epsilon + \beta(T_{A,B_0}))U_{B_0}) \cap (b_1 + (\epsilon + \beta(T_{A,B_1}))U_{B_1})$. Hence, $E(b_0, b_1) \neq \emptyset$, and so the element $w = w(b_0, b_1) \in \tilde{B}(\epsilon)$ must satisfy

$$\|Ta - w\|_{B_0} \leq \|Ta - b_0\|_{B_0} + \|b_0 - w\|_{B_0} \leq 2(\epsilon + \beta(T_{A,B_0})),$$

and also

$$\|Ta - w\|_{B_1} \leq \|Ta - b_1\|_{B_1} + \|b_1 - w\|_{B_1} \leq 2(\epsilon + \beta(T_{A,B_1})).$$

Consequently, for each $t > 0$,

$$\begin{aligned} \|Ta - w\|_B &\leq \frac{1}{\rho(t, B, \bar{B})} J(t, Ta - w; \bar{B}) \\ &\leq \frac{2}{\rho(t, B, \bar{B})} \max\{\epsilon + \beta(T_{A,B_0}), t(\epsilon + \beta(T_{A,B_1}))\}. \end{aligned}$$

It follows that

$$\beta(T_{A,B}) \leq \frac{2}{\rho(t, B, \bar{B})} \max\{\epsilon + \beta(T_{A,B_0}), t(\epsilon + \beta(T_{A,B_1}))\},$$

and so, since ϵ is arbitrary, we obtain

$$\beta(T_{A,B}) \leq \frac{2}{\rho(t, B, \bar{B})} \max\{\beta(T_{A,B_0}), t\beta(T_{A,B_1})\}. \quad (3.4)$$

To deal with cases (a) and (b) of the theorem, we simply substitute $\beta(T_{A,B_i}) = 0$ for $i = 0$ or $i = 1$ in (3.4) and use the facts that $1/\rho(t)$ is non-increasing and $t/\rho(t)$ is non-decreasing. We obtain case (c) by substituting $t = \beta(T_{A,B_0})/\beta(T_{A,B_1})$. \square

Now that we have prepared a number of tools, we can begin looking more closely at various ways in which natural generalizations of the Lions–Peetre compactness results may fail to hold.

Let us first consider the setting where we have $T : \bar{A} \rightarrow B$ and $T : A_1 \rightarrow B$ is compact. If $T : A_0 \rightarrow B$ is not compact, then there may exist intermediate spaces or even interpolation spaces A such that $T : A \rightarrow B$ is not compact. One trivial example is to take $A = A_0$. But the following example is more relevant to the general result that we shall obtain.

Example 3.3. Let $\bar{A} = (A_0, A_1) = (\ell_\infty, \ell_\infty(2^n))$, where ℓ_∞ is the usual space of scalar sequences $\xi = \{\xi_n\}_{n \in \mathbb{N}}$ satisfying $\|\xi\|_{\ell_\infty} := \sup_{n \in \mathbb{N}} |\xi_n| < \infty$ and $\ell_\infty(2^n)$ is the corresponding ‘weighted’ space of those sequences ξ for which $\|\xi\|_{\ell_\infty(2^n)} := \sup_{n \in \mathbb{N}} |2^n \xi_n| < \infty$. We choose $A = c_0$, the subspace of sequences in ℓ_∞ that converge to zero. This is an interpolation space since $A = A_0^\circ$. Then we take $B = \ell_\infty$ and let T be the identity operator $T\xi = \xi$. Clearly, $T : \bar{A} \rightarrow B$ and also $T : A_1 \rightarrow B$ is compact. Nevertheless, $T : A \rightarrow B$ is the inclusion from c_0 into ℓ_∞ which is not compact.

Remark 3.4. Note that whenever we have, as in the preceding example, that

$$A_0^\circ \hookrightarrow A, \tag{3.5}$$

and also that $T : A_0^\circ \rightarrow B$ is not compact, then of course $T : A \rightarrow B$ cannot be compact.

Let us now consider the ‘dual’ situation, where $T : A \rightarrow \bar{B}$ and $T : A \rightarrow B_1$ is compact. Analogously to before, if $T : A \rightarrow B_0$ is not compact, then $T : A \rightarrow B$ will not be compact in general for all intermediate spaces or even interpolation spaces B . Again, in addition to the obvious example $B = B_0$, we have another example that is more relevant for the results that we shall soon formulate.

Example 3.5. Let

$$\bar{B} = (B_0, B_1) = (c_0, \ell_\infty(2^{-n})), \quad \text{where } \|\xi\|_{\ell_\infty(2^{-n})} := \sup_{n \in \mathbb{N}} |2^{-n} \xi_n|,$$

and let $B = \ell_\infty$. It is easy to see that ℓ_∞ is the Gagliardo completion of $B_0 = c_0$ in $B_0 + B_1 = \ell_\infty(2^{-n})$. Thus, B is an interpolation space with respect to \bar{B} . We now choose $A = c_0$ and again let T be the identity operator. Then $T : A \rightarrow \bar{B}$ and $T : A \rightarrow B_1$ is compact, but $T : A \rightarrow B$ is not compact because (as in the previous example) it is the inclusion from c_0 into ℓ_∞ .

Remark 3.6. The condition analogous to (3.5) in this case is

$$B \hookrightarrow B_0^\sim. \tag{3.6}$$

Whenever it holds (as it does of course in Example 3.5) and $T : A \rightarrow B_0^\sim$ is not compact, then $T : A \rightarrow B$ cannot be compact. We shall soon see, in fact, that the only mechanisms

that can prevent a compactness result of Lions–Peetre type from holding for an arbitrary interpolation or rank-one interpolation space are those described in this remark and in Remark 3.4.

The next two lemmas will sometimes enable us to deduce that the conditions mentioned in Remarks 3.4 and 3.6 hold when either A or B , respectively, is a rank-one interpolation space.

Lemma 3.7. *Let $\bar{A} = (A_0, A_1)$ be a Banach couple and let A be a rank-one interpolation space with respect to \bar{A} .*

(i) *If $\lim_{t \rightarrow 0} \psi(t, A, \bar{A}) > 0$, then $A_0^\circ \hookrightarrow A$.*

(ii) *If $\lim_{t \rightarrow \infty} (\psi(t, A, \bar{A})/t) > 0$, then $A_1^\circ \hookrightarrow A$.*

Proof. Suppose $\lim_{t \rightarrow 0} \psi(t) > 0$. Then, by Theorem 2.2, we have $\lim_{t \rightarrow 0} \rho(t, A, \bar{A}) = M > 0$. Since ρ is non-decreasing, it follows that $M \leq (J(t, a)/\|a\|_A)$ for all $t > 0$ and all $a \in A_0 \cap A_1$. Therefore, $\|a\|_A \leq (1/M) \max\{\|a\|_{A_0}, t\|a\|_{A_1}\}$. Since we may choose $t = \|a\|_{A_0}/\|a\|_{A_1}$, this implies that $\|a\|_A \leq (1/M)\|a\|_{A_0}$ for all $a \in A_0 \cap A_1$, and, consequently, $A_0^\circ \hookrightarrow A$.

The proof of (ii) is similar. \square

Lemma 3.8. *Let $\bar{B} = (B_0, B_1)$ be a Banach couple and let B be a rank-one interpolation space with respect to \bar{B} .*

(i) *If $\lim_{t \rightarrow \infty} (1/\rho(t, B, \bar{B})) > 0$, then $B \hookrightarrow B_0^\sim$.*

(ii) *If $\lim_{t \rightarrow 0} (t/\rho(t, B, \bar{B})) > 0$, then $B \hookrightarrow B_1^\sim$.*

Proof. Assume that $\lim_{t \rightarrow \infty} (1/\rho(t)) > 0$. Then, by Theorem 2.2, we have that $\lim_{t \rightarrow \infty} \psi(t, B, \bar{B}) = M < \infty$. Consequently, each $b \in B$ satisfies

$$\|b\|_{B_0^\sim} = \lim_{t \rightarrow \infty} K(t, b; \bar{B}) \leq \lim_{t \rightarrow \infty} \psi(t) \|b\|_B = M \|b\|_B,$$

and so $B \hookrightarrow B_0^\sim$.

Case (ii) can be treated analogously. \square

We can now give a precise formulation of our claim made at the end of Remark 3.6. This is done in the following two theorems, which are almost immediate consequences of the preceding results.

Theorem 3.9. *Let $\bar{A} = (A_0, A_1)$ be a Banach couple, let A be a rank-one interpolation space with respect to \bar{A} and let B be another Banach space. Let $T : \bar{A} \rightarrow B$ be a linear operator such that $T : A_1 \rightarrow B$ is compact. Then at least one of the following conditions must hold.*

(i) *$T : A \rightarrow B$ is compact.*

(ii) *$A_0^\circ \hookrightarrow A$.*

Proof. Since $\beta(T_{A_1,B}) = 0$, Theorem 3.1 implies that

$$\beta(T_{A,B}) \leq \beta(T_{A_0,B}) \cdot \lim_{t \rightarrow 0} \psi(t, A, \bar{A}).$$

Consequently, either $T : A \rightarrow B$ is compact, i.e. $\beta(T_{A,B}) = 0$, or, alternatively, we have $\beta(T_{A,B}) > 0$. In this latter case we must have $\lim_{t \rightarrow 0} \psi(t, A, \bar{A}) > 0$, which, by Lemma 3.7, implies that $A_0^\circ \hookrightarrow A$. \square

Here is the analogous result for the ‘dual’ situation.

Theorem 3.10. *Let $\bar{B} = (B_0, B_1)$ be a Banach couple, let B be a rank-one interpolation space with respect to \bar{B} and let A be another Banach space. Let $T : A \rightarrow \bar{B}$ be a linear operator such that $T : A \rightarrow B_1$ is compact. Then at least one of the following conditions must hold.*

- (i) $T : A \rightarrow B$ is compact.
- (ii) $B \hookrightarrow B_0^\wedge$.

Proof. According to Theorem 3.2 we have

$$\beta(T_{A,B}) \leq 2\beta(T_{A,B_0}) \cdot \lim_{t \rightarrow \infty} \frac{1}{\rho(t, B, \bar{B})}.$$

So, if $T : A \rightarrow B$ is not compact, we must necessarily have

$$0 < \beta(T_{A,B}) \leq 2\beta(T_{A,B_0}) \cdot \lim_{t \rightarrow \infty} \frac{1}{\rho(t, B, \bar{B})},$$

and, consequently, by Lemma 3.8, $B \hookrightarrow B_0^\wedge$. \square

Now, it is easy to show that, under mild additional conditions on the Banach couples, the known sufficient conditions for $T : A \rightarrow B$ to be compact are also necessary.

Corollary 3.11. *Let $\bar{A} = (A_0, A_1)$ be a Banach couple such that $A_0^\circ = A_0$, let A be a rank-one interpolation space with respect to \bar{A} and let B be another Banach space. Let $T : \bar{A} \rightarrow B$ be a linear operator such that $T : A_1 \rightarrow B$ is compact. Then $T : A \rightarrow B$ is compact if and only if at least one of the following conditions hold.*

- (i) $\lim_{t \rightarrow 0} \psi(t, A, \bar{A}) = 0$.
- (ii) $T : A_0 \rightarrow B$ is compact.

Proof. From Theorem 3.1 (b) it is clear that either of conditions (i) and (ii) is sufficient to ensure that $T : A \rightarrow B$ is compact. Conversely, if $T : A \rightarrow B$ is compact and if (i) does not hold, then we deduce from Lemma 3.7 that $A_0^\circ \hookrightarrow A$, which in turn implies that $T : A_0^\circ \rightarrow B$ is compact. Since $A_0^\circ = A_0$ we obtain (ii), completing the proof. \square

Corollary 3.12. *Let $\bar{B} = (B_0, B_1)$ be a Banach couple such that $B_0^\wedge = B_0$, let B be a rank-one interpolation space with respect to \bar{B} and let A be another Banach space. Let $T : A \rightarrow \bar{B}$ be a linear operator such that $T : A \rightarrow B_1$ is compact. Then $T : A \rightarrow B$ is compact if and only if at least one of the following conditions hold.*

- (i) $\lim_{t \rightarrow \infty} \rho(t, B, \bar{B}) = \infty$.
(ii) $T : A \rightarrow B_0$ is compact.

Proof. The argument is analogous to the proof of the preceding corollary. By Theorem 3.2 (b) each of conditions (i) and (ii) is sufficient for the compactness of $T : A \rightarrow B$. Conversely, if $T : A \rightarrow B$ is compact and (i) does not hold, then Lemma 3.8 gives that $B \hookrightarrow B_0^\circ = B_0$, which implies (ii). \square

Remark 3.13. It is easy to see that the last two corollaries cannot be sharpened to hold without the condition $A_0^\circ = A_0$ or $B_0^\circ = B_0$, respectively. For example, suppose, in the context of Corollary 3.11, that $A_1 \hookrightarrow A_0$ and that A is also an interpolation space with respect to the couple (A_0°, A_1) . Then the behaviour of T on the space A is completely determined by the behaviour of T on A_0° , and it is completely irrelevant to ask whether or not T maps the larger space A_0 compactly into B . The following example can illustrate this more explicitly. Suppose that $\bar{E} = (E_0, E_1)$ is a Banach couple such that $E_1 \hookrightarrow E_0 = E_0^\circ$. Define a new Banach couple $\bar{A} = (A_0, A_1)$ by setting $A_0 = E_0 \oplus Y$ and $A_1 = E_1 \oplus \{0\}$, where Y is an infinite-dimensional Banach space and $\{0\}$ denotes its trivial subspace. (The norms for A_0 and A_1 may be taken, say, to be $\|(x, y)\|_{A_0} = \max\{\|x\|_{E_0}, \|y\|_Y\}$ and $\|(x, 0)\|_{A_1} = \|x\|_{E_1}$.) Each operator $S : \bar{A} \rightarrow \bar{A}$ can be decomposed into a 2×2 ‘matrix’

$$S = \begin{pmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{pmatrix},$$

i.e. $S(x, y) = (S_{11}x + S_{12}y, S_{21}x + S_{22}y)$ for all $(x, y) \in E_0 \oplus Y$. Since $S : A_0 \rightarrow A_0$ is bounded, it follows that each of the four operators $S_{11} : E_0 \rightarrow E_0$, $S_{12} : Y \rightarrow E_0$, $S_{21} : E_0 \rightarrow Y$ and $S_{22} : Y \rightarrow Y$ is bounded. Since $S : A_1 \rightarrow A_1$ is also bounded, we deduce that $S_{11} : E_1 \rightarrow E_1$, i.e. that $S_{11} : \bar{E} \rightarrow \bar{E}$, and also that $S_{21} = 0$. From this description of the operator S it follows that whenever E is an interpolation space with respect to \bar{E} , then $A = E \oplus \{0\}$ is an interpolation space with respect to \bar{A} .

Now, let B be another infinite-dimensional Banach space and consider a bounded operator $T : A_0 \rightarrow B$ such that $T : A_1 \rightarrow B$ is compact. This means that T has the form $T(x, y) = T_1x + T_2y$ for all $x \in E_0$ and $y \in Y$, where $T_1 : E_0 \rightarrow B$ is bounded, $T_1 : E_1 \rightarrow B$ is compact, and $T_2 : Y \rightarrow B$ is bounded. If A is an interpolation space of the form $A = E \oplus \{0\}$, then $T : A \rightarrow B$ is compact if and only if $T_1 : E \rightarrow B$ is compact. Consequently, we can apply Corollary 3.11 to the operator $T_1 : \bar{E} \rightarrow B$. We obtain that $T : A \rightarrow B$ is compact if and only if either $\lim_{t \rightarrow 0} \psi(t, E, \bar{E}) = 0$ or $T_1 : E_0 \rightarrow B$ is compact or both of these conditions hold. The latter condition is the same as the compactness of $T : A_0^\circ \rightarrow B$, since $A_0^\circ = E_0 \oplus \{0\}$. It is now quite clear that in this case the compactness of $T : A_0 \rightarrow B$ is irrelevant: the compactness of $T : A_0 \rightarrow B$ is equivalent to the condition that both of the operators $T_1 : E_0 \rightarrow B$ and $T_2 : Y \rightarrow B$ are compact, but the behaviour of T_2 has no effect on the compactness of $T : A \rightarrow B$.

In the last two results of this section we do not have to confine ourselves to rank-one interpolation spaces. Instead we can deal with all intermediate spaces. But, before

stating those results, we give the details of an example, kindly pointed out to us by Evgeniy Pustylnik, which shows that there is necessarily a difference between the case of rank-one interpolation spaces and general intermediate spaces. More specifically, it shows that we cannot obtain a result analogous to Theorem 3.9 if we assume that A is merely an intermediate space rather than a rank-one interpolation space.

Example 3.14. Let $\bar{A} = (A_0, A_1) = (L_p(0, 1), L_q(0, 1))$, where $1 < p < q < \infty$. Let A be the Lorentz space $L_{p,1}(0, 1)$, which is an intermediate space with respect to \bar{A} but not a rank-one space. (This latter claim can also be deduced from what is to follow.) Set

$$\lambda = 1 + \frac{1}{q} - \frac{1}{p}$$

and consider the potential operator T defined by

$$Tx(t) = \int_0^1 \frac{x(s)}{|t-s|^\lambda} ds.$$

As was shown in [11], $T : L_p \rightarrow L_q$ boundedly, and, consequently, we also have $T : L_q \rightarrow L_q$ boundedly. Moreover, by [13, Theorem 2.8.1], $T : L_q \rightarrow L_q$ is also compact. Despite these properties of T , neither of conditions (i) and (ii) in Theorem 3.9 holds for this example. First, we have $A_0^\circ = A_0 = L_p$, which is not contained in $A = L_{p,1}$. Furthermore, the sequence

$$x_k(t) = \begin{cases} k^{1/p}, & \text{if } 0 < t < (1/k), \\ 0, & \text{if } t \geq (1/k), \end{cases} \quad \text{for all } k \in \mathbb{N}$$

is bounded in $L_{p,1}$ because

$$\|x_k\|_{L_{p,1}} = \int_0^1 t^{(1/p)-1} x_k^*(t) dt = k^{1/p} \int_0^{1/k} t^{(1/p)-1} dt = p,$$

but, for any $t \in (0, 1/k)$, we have

$$\begin{aligned} Tx_k(t) &= k^{1/p} \int_0^{1/k} \frac{ds}{|t-s|^\lambda} \geq \left(\frac{1}{p} - \frac{1}{q}\right)^{-1} k^{(1/p)+\lambda-1} \\ &= \left(\frac{1}{p} - \frac{1}{q}\right)^{-1} k^{1/q}, \end{aligned}$$

and, thus,

$$\|Tx_k \chi_{(0,1/k)}\|_{L_q} \geq 1, \quad \text{for all } k \in \mathbb{N}.$$

This means that the functions Tx_k do not possess equi-absolutely continuous norms in L_q as $k \rightarrow \infty$. As was shown in [13], such a sequence of functions is not compact in L_q . Thus, $T : L_{p,1} \rightarrow L_q$ is not compact.

Our first result for the case of general intermediate spaces is an analogue of Corollary 3.11, where we simultaneously consider all possible range spaces B .

Theorem 3.15. Let $\bar{A} = (A_0, A_1)$ be a Banach couple and let A be an intermediate space with respect to \bar{A} . Suppose that either $A \cap A_1$ is dense in A , or that $A_0 \cap A_1$ is dense in A_0 , or that

$$\lim_{t \rightarrow 0} K(t, a; \bar{A}) = 0, \quad \text{for all } a \in A. \quad (3.7)$$

Then the following are equivalent.

- (i) $\lim_{t \rightarrow 0} \psi(t, A, \bar{A}) = 0$.
- (ii) For every Banach space B , if $T : \bar{A} \rightarrow B$ is a linear operator such that $T : A_1 \rightarrow B$ is compact, then $T : A \rightarrow B$ is compact.
- (iii) If $T : \bar{A} \rightarrow \ell_\infty$ is a linear operator such that $T : A_1 \rightarrow \ell_\infty$ is compact, then $T : A \rightarrow \ell_\infty$ is compact.

Remark 3.16. Since every separable Banach space can be isometrically embedded into ℓ_∞ , it is not surprising that it is sufficient to check condition (ii) just for the special case $B = \ell_\infty$. It will be clear from the proof that if $A_0^\circ = A_0$, then it is also sufficient to check for the special case $B = c_0$.

Proof. Let us first observe that the condition that we really need here is (3.7). It is easy to verify that it is a consequence of either of the two density conditions in the statement of the theorem.

Once again we use Theorem 3.1 (b) to obtain that (i) implies (ii). Of course (ii) implies (iii). (Both of these implications also hold when the additional density conditions or (3.7) are not satisfied.) To complete the proof we will show that (iii) implies (i), or, equivalently, that if (i) does not hold, then (iii) cannot hold.

Thus we assume that $\lim_{t \rightarrow 0} \psi(t, A, \bar{A}) = \delta > 0$, which implies that there exist a sequence $\{a_n\}_{n \in \mathbb{N}}$ of elements in U_A and a decreasing sequence $\{t_n\}_{n \in \mathbb{N}}$ of numbers in $(0, 1)$ such that $\lim_{n \rightarrow \infty} t_n = 0$ and $K(t_n, a_n; \bar{A}) > \delta/2$. In view of (3.7) we can also assume, by passing if necessary to subsequences of the original sequences, that

$$K(t_n, a_m; \bar{A}) < \delta/4, \quad \text{for all } m < n.$$

By the Hahn–Banach theorem, for each $n \in \mathbb{N}$, there exists a linear functional f_n on $A_0 + A_1$ such that $f_n(a_n) = K(t_n, a_n; \bar{A})$ and $|f_n(a)| \leq K(t_n, a; \bar{A})$ for each $a \in A_0 + A_1$. Now consider the linear operator $T : A_0 + A_1 \rightarrow \ell_\infty$ defined by $Ta = \{f_n(a)\}_{n \in \mathbb{N}}$. Clearly, $T : A_i \rightarrow \ell_\infty$ with norm not exceeding 1 for $i = 0, 1$. Furthermore, T maps the unit ball of A_1 into the set of all sequences $\{\lambda_n\}_{n \in \mathbb{N}}$ that satisfy $|\lambda_n| \leq t_n$. This is clearly a compact subset of c_0 and of ℓ_∞ . On the other hand, for each pair of positive integers $m < n$ we have

$$\|Ta_n - Ta_m\|_{\ell_\infty} \geq |f_n(a_n) - f_n(a_m)| \geq K(t_n, a_n) - K(t_n, a_m) > \frac{1}{4}\delta.$$

Thus, $T : A \rightarrow \ell_\infty$ is not compact, and so (iii) does not hold. \square

Finally we present a result analogous to Corollary 3.12 which is essentially a dual of the preceding theorem. This time we simultaneously consider all possible domain spaces A . In contrast to the previous theorem, we do not need to impose any additional conditions.

Theorem 3.17. *Let $\bar{B} = (B_0, B_1)$ be a Banach couple and let B be an intermediate space with respect to \bar{B} . Then the following are equivalent.*

- (i) $\lim_{t \rightarrow \infty} \rho(t, B, \bar{B}) = \infty$.
- (ii) For every Banach space A , if $T : A \rightarrow \bar{B}$ is a linear operator such that $T : A \rightarrow B_1$ is compact, then $T : A \rightarrow B$ is compact.
- (iii) If $T : \ell_1 \rightarrow \bar{B}$ is a linear operator such that $T : \ell_1 \rightarrow B_1$ is compact, then $T : \ell_1 \rightarrow B$ is compact.

Proof. We obtain (i) \Rightarrow (ii) by Theorem 3.2 (b). Obviously, (ii) \Rightarrow (iii). To show that (iii) \Rightarrow (i), let us suppose that (i) does not hold. Then $\lim_{t \rightarrow \infty} \rho(t, B, \bar{B}) = \delta \in (0, \infty)$ and there exist a sequence $\{b_n\}_{n \in \mathbb{N}}$ of elements in $B_0 \cap B_1$ with $\|b_n\|_B = 1$ and $\|b_n\|_{B_0} < 2\delta$ and a sequence $\{t_n\}_{n \in \mathbb{N}}$ of numbers in $(1, \infty)$ such that $\lim_{n \rightarrow \infty} t_n = \infty$ and $\|b_n\|_{B_1} < 2\delta/t_n$. We consider the operator $T : \ell_1 \rightarrow B_0 + B_1$ defined by

$$T(\{\lambda_n\}_{n \in \mathbb{N}}) = \sum_{n \in \mathbb{N}} \lambda_n b_n.$$

Clearly, $T : \ell_1 \rightarrow B_i$ for $i = 0, 1$ with norm not exceeding 2δ . Furthermore, $T : \ell_1 \rightarrow B_1$ is compact, as the limit, in operator norm, of a sequence of finite-rank operators. However, $T : \ell_1 \rightarrow B$ is not compact. This follows immediately from the fact that the sequence $\{b_n\}$ does not contain any subsequence that converges in B . Were $\{b_{m_n}\}$ to be such a sequence, its limit b_* would necessarily satisfy $\|b_*\|_B = 1$. But this would be a contradiction, since this same sequence converges to 0 in $B_0 + B_1$ and B is an intermediate space. Thus we deduce that in this case (iii) cannot hold. This shows that (iii) \Rightarrow (i) and completes the proof. \square

4. Some results for other operator ideals

The class of compact operators between Banach spaces is an injective surjective closed operator ideal in the sense of Pietsch. It is therefore natural to investigate whether the results of § 3 are valid for other operator ideals having similar properties.

Let us recall (cf. [8, p. 131] and [17, p. 45]) that an *operator ideal* \mathcal{I} is a method of ascribing to each pair (X, Y) of Banach spaces, a linear subspace $\mathcal{I}(X, Y)$ of $\mathcal{L}(X, Y)$ such that

- (i) $\mathcal{I}(X, Y)$ contains all rank one operators $x^* \otimes y$ for all $x^* \in X^*$ and $y \in Y$; and
- (ii) for all Banach spaces X, Y, E, F , whenever $U \in \mathcal{L}(E, X)$, $T \in \mathcal{I}(X, Y)$ and $S \in \mathcal{L}(Y, F)$, then the composed operator $STU \in \mathcal{I}(E, F)$.

The ideal \mathcal{I} is *closed* if $\mathcal{I}(X, Y)$ is a closed subspace of $\mathcal{L}(X, Y)$ with respect to the usual operator norm topology, for all Banach spaces X and Y .

For each Banach space A we let Q_A denote the canonical surjection $Q_A : \ell_1(U_A) \rightarrow A$ defined by

$$Q_A(\{\lambda_x\}_{x \in U_A}) = \sum_{x \in U_A} \lambda_x x,$$

and J_A denote the canonical isometric embedding $J_A : A \rightarrow \ell_\infty(U_{A^*})$ defined by

$$J_A a = \{\langle f, a \rangle\}_{f \in U_{A^*}}.$$

The operator ideal \mathcal{I} is said to be *injective* (respectively, *surjective*) if whenever $T \in \mathcal{L}(A, B)$ and in addition $J_B T \in \mathcal{I}(A, \ell_\infty(U_{B^*}))$ (respectively, $TQ_A \in \mathcal{I}(\ell_1(U_A), B)$), then it follows that $T \in \mathcal{I}(A, B)$ (cf. [17, pp. 70–75]).

Apart from the class of compact operators, other examples of injective surjective closed ideals are the classes of weakly compact operators, Rosenthal operators, Banach–Saks operators and dual Radon–Nikodým operators. The class of strictly singular operators is an injective closed ideal which is not surjective, while the class of strictly cosingular operators is a surjective closed ideal which is not injective. We refer to [17] and also to [8] and [12] for more details about operator ideals.

There are two functionals, introduced, respectively, by Astala and by Tylli, which can be used to measure the extent to which a given operator $T \in \mathcal{L}(A, B)$ fails to belong to a given operator ideal \mathcal{I} . We shall now briefly recall their definitions and some of their main properties. We refer to [6] for relevant references and further details.

The *outer measure* of $T \in \mathcal{L}(A, B)$ is denoted by $\gamma_{\mathcal{I}}(T)$ or $\gamma_{\mathcal{I}}(T_{A,B})$ and is the infimum of all positive numbers σ such that $T(U_A) \subset \sigma U_B + R(U_Z)$ for some Banach space Z and some operator $R \in \mathcal{I}(Z, B)$. The *inner measure* of the same operator is denoted by $\beta_{\mathcal{I}}(T)$ or $\beta_{\mathcal{I}}(T_{A,B})$ and is the infimum of all positive numbers σ such that for some Banach space Z and some operator $R \in \mathcal{I}(A, Z)$ the inequality

$$\|Tx\|_B \leq \sigma \|x\|_A + \|Rx\|_Z$$

holds for all $x \in A$.

It is known that

$$\text{if } \mathcal{I} \text{ is surjective and closed, then } \gamma_{\mathcal{I}}(T_{A,B}) = 0 \text{ if and only if } T \in \mathcal{I}(A, B), \quad (4.1)$$

and, analogously,

$$\text{if } \mathcal{I} \text{ is injective and closed, then } \beta_{\mathcal{I}}(T_{A,B}) = 0 \text{ if and only if } T \in \mathcal{I}(A, B). \quad (4.2)$$

In the particular case where $\mathcal{I} = \mathcal{K}$, the ideal of compact operators, then $\gamma_{\mathcal{K}}(T_{A,B})$ coincides with the measure of non-compactness of T and $\beta_{\mathcal{K}}(T_{A,B})$ is the infimum of all $\eta > 0$ such that there exists a subspace M of A with finite codimension such that $\|Tx\|_B \leq \eta \|x\|_A$ for all $x \in M$.

The following theorem is an abstract version of Theorem 3.1. In its proof we use some techniques that were developed in [6].

Theorem 4.1. Let $\bar{A} = (A_0, A_1)$ be a Banach couple, let A be an intermediate space with respect to \bar{A} and let B be another Banach space. Let \mathcal{I} be an operator ideal. Then, for each $T \in \mathcal{L}(\bar{A}, B)$:

(a) if $\gamma_{\mathcal{I}}(T_{A_0, B}) = 0$, then

$$\gamma_{\mathcal{I}}(T_{A, B}) \leq \gamma_{\mathcal{I}}(T_{A_1, B}) \cdot \lim_{t \rightarrow \infty} \frac{\psi(t, A, \bar{A})}{t};$$

(b) if $\gamma_{\mathcal{I}}(T_{A_1, B}) = 0$, then

$$\gamma_{\mathcal{I}}(T_{A, B}) \leq \gamma_{\mathcal{I}}(T_{A_0, B}) \cdot \lim_{t \rightarrow 0} \psi(t, A, \bar{A});$$

and

(c) if $\gamma_{\mathcal{I}}(T_{A_i, B}) > 0$ for $i = 0, 1$, then

$$\gamma_{\mathcal{I}}(T_{A, B}) \leq 2\gamma_{\mathcal{I}}(T_{A_0, B})\psi\left(\frac{\gamma_{\mathcal{I}}(T_{A_1, B})}{\gamma_{\mathcal{I}}(T_{A_0, B})}, A, \bar{A}\right).$$

Proof. By the definition of $\gamma_{\mathcal{I}}(T_{A_i, B})$, for $i = 0, 1$ and for each $\epsilon > 0$ there exist Banach spaces Z_i and operators $S_i \in \mathcal{I}(Z_i, B)$ such that

$$T(U_{A_i}) \subset (\epsilon + \gamma_{\mathcal{I}}(T_{A_i, B}))U_B + S_i(U_{Z_i}). \tag{4.3}$$

Now, for each choice of positive numbers δ and t and each $a \in U_A$, there exists a decomposition $a = a_0 + a_1$ with $a_i \in A_i$ and $a_i \in (\delta + \psi(t))t^{-i}U_{A_i}$ for $i = 0, 1$. From this and (4.3) it follows that

$$\begin{aligned} T(U_A) &\subset (\delta + \psi(t))\left[T(U_{A_0}) + \frac{1}{t}T(U_{A_1})\right] \\ &\subset (\delta + \psi(t))\left[\epsilon + \gamma_{\mathcal{I}}(T_{A_0, B}) + \frac{\epsilon}{t} + \frac{1}{t}\gamma_{\mathcal{I}}(T_{A_1, B})\right]U_B + S'_0(U_{Z_0}) + S'_1(U_{Z_1}), \end{aligned}$$

where $S'_0 = (\delta + \psi(t))S_0$ and $S'_1 = (1/t)(\delta + \psi(t))S_1$ are operators belonging to $\mathcal{I}(Z_0, B)$ and $\mathcal{I}(Z_1, B)$, respectively. Let Z be the Banach space $Z = Z_0 \oplus Z_1$ with norm $\|(x, y)\|_Z = \max\{\|x\|_{Z_0}, \|y\|_{Z_1}\}$ and define $S : Z \rightarrow B$ by $S(x, y) = S'_0x + S'_1y$. Then clearly $S(U_Z) = S'_0(U_{Z_0}) + S'_1(U_{Z_1})$, and, furthermore, using the projection operators from Z onto Z_0 and Z_1 , and the definition of operator ideals, we have that $S \in \mathcal{I}(Z, B)$. Consequently,

$$\gamma_{\mathcal{I}}(T_{A, B}) \leq \psi(t)\left[\gamma_{\mathcal{I}}(T_{A_0, B}) + \frac{1}{t}\gamma_{\mathcal{I}}(T_{A_1, B})\right], \quad \text{for all } t > 0.$$

This inequality implies the three cases of the theorem by the same immediate arguments as were used to obtain the three cases of Theorem 3.1 from equation (3.2). \square

We now consider an abstract version of Theorem 3.2. Note, however, that in the particular case $\mathcal{I} = \mathcal{K}$, it yields a result different from Theorem 3.2, since here we deal with the inner rather than outer measure. (Also the factors of 2 that appear in Theorem 3.2 do not appear here.)

Theorem 4.2. Let $\bar{B} = (B_0, B_1)$ be a Banach couple, let B be an intermediate space with respect to \bar{B} and let A be another Banach space. Let \mathcal{I} be an operator ideal. Then, for each operator $T \in \mathcal{L}(A, \bar{B})$:

(a) if $\beta_{\mathcal{I}}(T_{A, B_0}) = 0$, then

$$\beta_{\mathcal{I}}(T_{A, B}) \leq \beta_{\mathcal{I}}(T_{A, B_1}) \cdot \lim_{t \rightarrow 0} \frac{t}{\rho(t, B, \bar{B})};$$

(b) if $\beta_{\mathcal{I}}(T_{A, B_1}) = 0$, then

$$\beta_{\mathcal{I}}(T_{A, B}) \leq \beta_{\mathcal{I}}(T_{A, B_0}) \cdot \lim_{t \rightarrow \infty} \frac{1}{\rho(t, B, \bar{B})};$$

and

(c) if $\beta_{\mathcal{I}}(T_{A, B_i}) > 0$ for $i = 0, 1$, then

$$\beta_{\mathcal{I}}(T_{A, B}) \leq \frac{\beta_{\mathcal{I}}(T_{A, B_0})}{\rho(\beta_{\mathcal{I}}(T_{A, B_0})/\beta_{\mathcal{I}}(T_{A, B_1}), B, \bar{B})}.$$

Proof. In view of the definition of $\beta_{\mathcal{I}}(T_{A, B_i})$, given any $\epsilon > 0$, there are Banach spaces Z_i and operators $S_i \in \mathcal{I}(A, Z_i)$ such that

$$\|Ta\|_{B_i} \leq (\epsilon + \beta_{\mathcal{I}}(T_{A, B_i}))\|a\|_A + \|S_i a\|_{Z_i}, \quad \text{for all } a \in A, \quad i = 0, 1.$$

Since $Ta \in B_0 \cap B_1$ and $\|b\|_B \leq (1/\rho(t, B, \bar{B}))J(t, b, \bar{B})$ for all $b \in B_0 \cap B_1$ and all $t > 0$, we obtain that

$$\begin{aligned} \|Ta\|_B &\leq \frac{1}{\rho(t)} J(t, Ta) \\ &\leq \frac{1}{\rho(t)} \max\{(\epsilon + \beta_{\mathcal{I}}(T_{A, B_0})), t(\epsilon + \beta_{\mathcal{I}}(T_{A, B_1}))\} \|a\|_A + \|S'_0 a\|_{Z_0} + \|S'_1 a\|_{Z_1}, \end{aligned}$$

where $S'_0 = (1/\rho(t))S_0$ and $S'_1 = (t/\rho(t))S_1$. Let Z be the Banach space $Z = Z_0 \oplus Z_1$ with norm $\|(x, y)\|_Z = \|x\|_{Z_0} + \|y\|_{Z_1}$ and consider the operator $S : A \rightarrow Z$ defined by $Sa = (S'_0 a, S'_1 a)$. Then, using the ideal properties of \mathcal{I} and the canonical embeddings of Z_0 and Z_1 into Z , we see that $S \in \mathcal{I}(A, Z)$ and we have

$$\|Ta\|_B \leq \frac{1}{\rho(t)} \max\{(\epsilon + \beta_{\mathcal{I}}(T_{A, B_0})), t(\epsilon + \beta_{\mathcal{I}}(T_{A, B_1}))\} \|a\|_A + \|Sa\|_Z, \quad \text{for all } a \in A.$$

This implies that

$$\beta_{\mathcal{I}}(T_{A, B}) \leq \frac{1}{\rho(t)} \max\{\beta_{\mathcal{I}}(T_{A, B_0}), t\beta_{\mathcal{I}}(T_{A, B_1})\}, \quad \text{for all } t > 0,$$

and we can conclude the proof in the same way as at the end of the proof of Theorem 3.2. \square

Suppose that $T : \bar{A} \rightarrow B$ and $T \in \mathcal{I}(A_1, B)$ for some operator ideal \mathcal{I} . For which intermediate spaces or rank-one interpolation spaces A do we necessarily have that $T \in \mathcal{I}(A, B)$? Under what circumstances can it happen that $T \notin \mathcal{I}(A, B)$?

The preceding two theorems enable us to find answers to these questions and the corresponding questions for operators $T : A \rightarrow \bar{B}$ with $T \in \mathcal{I}(A, B_1)$. These answers take the form of analogues of Theorems 3.9 and 3.10 and Corollaries 3.11 and 3.12, where the ideal \mathcal{K} of compact operators is replaced by an appropriate more general operator ideal \mathcal{I} .

First we state a result that incorporates the analogues of Theorem 3.9 and Corollary 3.11. Here, we need the ideal to be surjective and closed so that we can use property (4.1). The proof of the theorem can then follow exactly the same path as its predecessor theorem and corollary, except that the inner measure $\gamma_{\mathcal{I}}$ now plays the former role of the measure of non-compactness.

Theorem 4.3. *Let $\bar{A} = (A_0, A_1)$ be a Banach couple, let A be a rank-one interpolation space with respect to \bar{A} and let B be another Banach space. Let \mathcal{I} be a surjective closed operator ideal and let $T : \bar{A} \rightarrow B$ be a linear operator such that $T \in \mathcal{I}(A_1, B)$. Then at least one of the following conditions must hold.*

- (i) $T \in \mathcal{I}(A, B)$.
- (ii) $A_0^\circ \hookrightarrow A$.

If, in addition, the couple \bar{A} satisfies $A_0^\circ = A_0$, then $T \in \mathcal{I}(A, B)$ if and only if at least one of the following conditions hold.

- (i') $\lim_{t \rightarrow 0} \psi(t, A, \bar{A}) = 0$.
- (ii') $T \in \mathcal{I}(A_0, B)$.

Now we turn to a theorem that contains the analogues of Theorem 3.10 and Corollary 3.12. Here, the ideal must be taken to be injective and closed so that the proof can use property (4.2) and the outer measure $\beta_{\mathcal{I}}$ in place of the measure of non-compactness.

Theorem 4.4. *Let $\bar{B} = (B_0, B_1)$ be a Banach couple, let B be a rank-one interpolation space with respect to \bar{B} , and let A be another Banach space. Let \mathcal{I} be an injective closed operator ideal, and suppose that $T \in \mathcal{L}(A, \bar{B})$ satisfies $T \in \mathcal{I}(A, B_1)$. Then at least one of the following conditions must hold.*

- (i) $T \in \mathcal{I}(A, B)$.
- (ii) $B \hookrightarrow B_0^\frown$.

If, in addition, we have $B_0^\frown = B_0$, then $T \in \mathcal{I}(A, B)$ if and only if at least one of the following conditions hold.

- (i') $\lim_{t \rightarrow \infty} \rho(t, B, \bar{B}) = \infty$.
- (ii') $T \in \mathcal{I}(A, B_0)$.

Theorems 4.1 and 4.2 immediately lead to stronger versions of Propositions 1.6 and 1.7 in Heinrich [12], i.e. if \mathcal{I} is closed and, respectively, surjective or injective, and if $T \in \mathcal{I}(A_1, B)$ or, respectively, $T \in \mathcal{I}(A, B_1)$, and if (1.1) or, respectively, (1.2) holds, then Theorem 4.1 or, respectively, Theorem 4.2 immediately gives us that $T \in \mathcal{I}(A, B)$. Applying these results to the ideals of strictly singular operators and strictly cosingular operators, we obtain new information about the behaviour under interpolation of these operator ideals which complements the results established by Beucher in [3, §2].

In the cases where A or B , respectively, are merely intermediate spaces rather than rank-one interpolation spaces we apparently, in general, do not have analogues of Theorems 3.15 and 3.17.

Acknowledgements. F.C. was supported in part by Dirección General de Enseñanza Superior (PB97-0254). M.C. was supported by the Fund for Promotion of Research at the Technion. P.M. was supported by Progamma Praxis XXI da Fundação para a Ciência e Tecnologia (BD-11466-97) and by Instituto Politécnico de Leiria. We are grateful to Evgeniy Pustyl'nik for drawing our attention to an interesting example and also for a number of other helpful suggestions and remarks.

References

1. C. BENNETT AND R. SHARPLEY, *Interpolation of operators* (Academic Press, New York, 1988).
2. J. BERGH AND J. LÖFSTRÖM, *Interpolation spaces. An introduction*, Grundlehren der Mathematische Wissenschaften, vol. 223 (Springer, Berlin, 1976).
3. O. J. BEUCHER, On interpolation of strictly (co-)singular linear operators, *Proc. R. Soc. Edinb. A* **112** (1989), 263–269.
4. Y. BRUDNYI AND N. KRUGLJAK, *Interpolation functors and interpolation spaces*, vol. 1 (North Holland, Amsterdam, 1991).
5. F. COBOS, T. KÜHN AND T. SCHONBEK, One-sided compactness results for Aronszajn–Gagliardo functors, *J. Funct. Analysis* **106** (1992), 274–313.
6. F. COBOS, A. MANZANO AND A. MARTÍNEZ, Interpolation theory and measures related to operator ideals, *Q. J. Math. Oxford* (2) **50** (1999), 401–416.
7. F. COBOS AND J. PEETRE, Interpolation of compactness using Aronszajn–Gagliardo functors, *Israel J. Math.* **68** (1989), 220–240.
8. J. DIESTEL, H. JARCHOW AND A. TONGE, *Absolutely summing operators*, Cambridge Studies in Advanced Mathematics, vol. 43 (Cambridge University Press, 1995).
9. A. A. DMITRIEV, The interpolation of one-dimensional operators, *Voronež Gos. Univ. Trudy Naučn.-Issled. Inst. Mat. VGU Vyp. 11 Sb. Statej Funkcional. Anal. i Priložen* **11** (1973), 31–43 (in Russian).
10. E. GAGLIARDO, A unified structure in various families of function spaces. Compactness and closure theorems, in *Proc. Int. Symp. Linear Spaces, Jerusalem 1960*, pp. 237–241 (Pergamon, Oxford, 1961).
11. G. H. HARDY AND J. E. LITTLEWOOD, Some properties of fractional integrals (I), *Math. Z.* **27** (1928), 565–606.
12. S. HEINRICH, Closed operator ideals and interpolation, *J. Funct. Analysis* **35** (1980), 397–411.
13. M. A. KRASNOSELSKII, P. P. ZABREIKO, E. I. PUSTYL'NIK AND P. E. SOBOLEVSKII, *Integral operators in spaces of summable functions* (Noordhoff, Leyden, 1976).

14. J. L. LIONS AND J. PEETRE, Sur une classe d'espaces d'interpolation, *Inst. Hautes Etudes Sci. Publ. Math.* **19** (1964), 5–68.
15. M. MASTYŁO, On interpolation of compact operators, *Funct. Approx. Comment. Math.* **26** (1998), 293–311.
16. A. PERSSON, Compact linear mappings between interpolation spaces, *Ark. Mat.* **5** (1964), 215–219.
17. A. PIETSCH, *Operator ideals* (North-Holland, Amsterdam, 1980).
18. E. I. PUSTYLNİK, On optimal interpolation and some interpolation properties of Orlicz spaces, *Soviet. Math. Dokl.* **27** (1983), 333–336. (Translation from *Dokl. Akad. Nauk SSSR* **269** (1983), 292–295.)
19. E. I. PUSTYLNİK, Embedding functions and their role in interpolation theory, *Abstract Appl. Analysis* **1** (1996), 305–325.
20. M. F. TEIXEIRA AND D. E. EDMUNDS, Interpolation theory and measures of non-compactness, *Math. Nachr.* **104** (1981), 129–135.