

## CORRESPONDENCE

The Editor,  
*Journal of Glaciology*

SIR,

### *About the origin of rock glaciers*

I do not wish to enter into a public controversy with W. Haerberli about the origin of rock glaciers; he has always been deaf to my arguments. Nevertheless, the readers of his passionate assertions (Haerberli, 1989) must be aware that he intentionally omits to quote my detailed observations in the dry Andes (Lliboutry, 1955, 1965, 1986). The concept that many rock glaciers come from old glaciers, or from "buried glacierets" (i.e. debris-rich glacierets, entirely covered at the end of the ablation season, and not nourished by ice every year) is not at all "purely speculative", as he says. If an interested glaciologist comes to Grenoble, I can show him the aerial photograph coverage of the area that I have studied, with scores of such rock glaciers.

I do not deny that many (not all) rock glaciers are below melting point at depth. In this case, they consist of permafrost, but should slide on unfrozen ground ... . Nevertheless, for the advancement of science, the essential point is not "must rock glaciers be left to scientists claiming to be permafrost specialists" but "what can we learn from the existence of rock glaciers in a given area"? I maintain that the geographical study of rock glaciers as an extreme case of glacier fluctuations, as an indicator of favourable mass balances in the past, or of past surges, would be much more rewarding than to consider them as a mere case of standard permafrost, or of creeping regolith.

[Université Joseph Fourier]  
Laboratoire de Glaciologie et Géophysique  
de l'Environnement du CNRS,  
38402 Saint-Martin-d'Hères Cedex,  
France

LOUIS LLIBOUTRY

26 October 1989

### REFERENCES

- Haerberli, W. 1989. Glacier ice-cored rock glaciers in the Yukon Territory, Canada? *J. Glaciol.*, 35(120), 294-295.  
Lliboutry, L. 1955. Origine et évolution des glaciers rocheux. *C.R. Acad. Sci. (Paris)*, 240, 1913-1915.  
Lliboutry, L. 1965. *Traité de glaciologie. Tome 2. Glaciers, variations du climat, sols gelés*. Paris, Masson.  
Lliboutry, L. 1986. Rock glaciers in the dry Andes. *Mater. Glyatsiol. Issled.*, 58, 139-144.

SIR,

### *An unusual jökulhlaup involving potholes on Black Rapids Glacier, Alaska Range, Alaska, U.S.A.*

Black Rapids Glacier is a large valley glacier in the Alaska Range which surged in 1936-37 (Hance, 1937) and was dubbed the "galloping glacier" by the popular press. The glacier surface is covered by several hundred potholes near the equilibrium line (Sturm, 1987). These depressions or cauldron-like pits, 10-100 m across, are occasionally water-filled, and are thought to be a complex surface and

englacial water system that is unique to surging glaciers. Twenty-six fields of potholes have been identified on the glaciers of Alaska and Yukon Territory; virtually all are on surging glaciers (though more than 200 surging glaciers do not have fields of potholes) (Sturm, 1987). Due to the remote location of most of these glaciers, including Black Rapids Glacier, there have been few direct observations of potholes filling or draining, and the nature of the water system connecting the potholes is poorly known.

On 16 August 1989, we observed water draining from a marginal ice-dammed lake and flowing through several potholes on Black Rapids Glacier in a manner that confirmed that the pothole-drainage system was both supra- and englacial. The jökulhlaup was coming from a lake located along the north margin of the glacier (Fig. 1-A) which has a history of being the source of jökulhlaups. A distinctive strandline on the ice surface indicated that water had recently been at least 20 m above the current lake surface (Fig. 1-B). The lake, which is over 1000 m long when full, probably fills and drains every summer. It was observed draining on 10 August 1985, and a draining event was photographed in July 1987 using an automatic 8 mm movie camera (unpublished data of T. Clarke). In both years, the discharge was englacial. In March 1984, the water level in the lake rose several meters and fractured a thick ice cover, even though below-freezing air temperatures prevailed and there was no surface run-off, suggesting that there was a significant recharge of water into the lake from an englacial source (personal communication from M. Wumkes, 1987).

In 1989, the discharge from the lake was supraglacial. Water flowed approximately 800 m through a vertical-walled canyon incised at least 20 m into the ice surface (Fig. 1-C). This canyon had several sharply angled bends in it, suggesting that fractures or weaknesses in the ice controlled its location.

The canyon ended where the water disappeared into the ice. Several hundred meters farther down-glacier the water emerged in a pothole that was approximately 50 m in diameter (Fig. 1-D). It was unmistakable that the water in the pothole was coming from the lake because both contained turbid, brown water, in contrast to the clear water seen elsewhere on the glacier surface. Water flowed over the down-glacier lip of the pothole, then through a shallow channel to a second pothole (Fig. 1-E). It cascaded into the second pothole in a waterfall which was at least 10 m high. The water in this pothole was observed to be swirling rapidly in a clockwise direction, suggesting that it was draining through a hole in the bottom of the pothole. The water carried numerous small icebergs and bits of ice. Dozens of small icebergs were stranded above the water surface along the down-glacier end of the pothole, indicating that the water level had been higher in the recent past.

The water flowed under the ice for several hundred meters and emerged in a third pothole (Fig. 1-F), where it was upwelling in a boil with significant super-elevation. Water flowed out of the third pothole and over the surface of the ice in a poorly defined, shallow channel into a fourth pothole (Fig. 1-G). The pothole is located at the extreme down-glacier end of the pothole field. Water flowed out of this pothole in a sheet which spread over the surface of the ice. During the 2 h we were in the vicinity, the sheet of water propagated several hundred meters down-glacier along the ice surface, covering a swath hundreds of meters wide. Crevasses, which are numerous in the section of glacier below the pothole field, were filled with water as the flood passed (Fig. 1-H).

Our observations verify several characteristics of potholes which we have suspected were true. First, potholes