

ITO'S FORMULA AND LÉVY'S LAPLACIAN

KIMIYAKI SAITO

§ 1. Introduction

The class of normal functionals

$$\int_{\mathbf{R}^n} \cdots \int f(x_1, \dots, x_n) : \dot{B}_{x_1}^{p_1} \cdots \dot{B}_{x_n}^{p_n} : dx_1 \cdots dx_n, \quad f \in L^1(\mathbf{R}^n),$$

$$(p_1, \dots, p_n) \in (N \cup \{0\})^n,$$

is, as is well known, adapted to the domain of Lévy's Laplacian and plays important roles in the works by P. Lévy and T. Hida (cf. [1], [2] and [8]), where \dot{B}_x denotes one-dimensional parameter white noise and $:\dot{B}_{x_1}^{p_1} \cdots \dot{B}_{x_n}^{p_n}:$ denotes the renormalization of $\dot{B}_{x_1}^{p_1} \cdots \dot{B}_{x_n}^{p_n}$.

We are interested in a generalization of this class to that of generalized functionals of two-dimensional parameter white noise $\{W(t, x); (t, x) \in \mathbf{R}^2\}$, which is a generalized stochastic process with the characteristic functional

$$C(\xi) = E(\exp \{i \langle W, \xi \rangle\}) = \exp \left\{ -\frac{1}{2} \|\xi\|^2 \right\}, \quad \xi \in S(\mathbf{R}^2).$$

As in the case [1], we are able to introduce, in Section 2, a space $(L^2)^{(-\alpha)}$ of generalized functionals and the \mathcal{S} -transform on $(L^2)^{(\alpha)}$ for every $\alpha > 0$. Then the calculus in terms of the white noise $W(t, x)$ will quickly be discussed.

The main purpose of this paper is to investigate how Lévy's Laplacian appears in Itô's formula for generalized Brownian functionals depending on t . To this end we first discuss a class of generalized Brownian functionals, often without any renormalization, having interest in its own right. For instance, a monomial $B_x(t)^p$ is sometimes more significant rather than the renormalized quantity $:B_x(t)^p:$ $\equiv : \left\{ \int_0^t W(r, x) dr \right\}^p :$ which is living in $(L^2)^{(-\alpha)}$. We are therefore led to construct a new space $[[L^2]]^{(-\alpha)}$,

Received April 11, 1986.

in which $B_x(t)^p$ lives, in Section 3. The \mathcal{S} -transform and the $W(t, x)$ -differentiation can be introduced on $[[L^2]]^{(-\alpha)}$ for every $\alpha > 0$ in a similar manner to those in [6]. The symbol $1/dx$ which has often been used by H.H. Kuo (cf. [7]) is now understood as a shift operator acting on $[[L^2]]^{(-\alpha)}$.

In Section 4, we define $B_{x_1}(\cdot)^{p_1} \cdots B_{x_n}(\cdot)^{p_n}$ by

$$B_{x_1}(\cdot)^{p_1} \cdots B_{x_n}(\cdot)^{p_n} = \left[\left[:B_{x_1}(\cdot)^{p_1} \cdots B_{x_n}(\cdot)^{p_n}: , \frac{p_1(p_1 - 1)}{2}(\cdot); B_{x_1}(\cdot)^{p_1-2} B_{x_2}(\cdot)^{p_2} \cdots B_{x_n}(\cdot)^{p_n} : + \cdots + \frac{p_n(p_n - 1)}{2}(\cdot); B_{x_1}(\cdot)^{p_1} \cdots B_{x_{n-1}}(\cdot)^{p_{n-1}} B_{x_n}(\cdot)^{p_n-2} : \right] \right],$$

for any $n \in \mathbb{N}$, $(p_1, \dots, p_n) \in (\mathbb{N} \cup \{0\})^n$, and $x_1, \dots, x_n \in \mathbb{R}$, and we introduce a class \mathcal{D}_L of generalized functionals as follows:

$$\mathcal{D}_L = LS \left\{ \int_{\mathbb{R}^n} f(x_1, \dots, x_n) B_{x_1}(\cdot)^{p_1} \cdots B_{x_n}(\cdot)^{p_n} dx_1 \cdots dx_n; f \in L^1(\mathbb{R}^n), (p_1, \dots, p_n) \in (\mathbb{N} \cup \{0\})^n, n = 0, 1, 2, \dots \right\},$$

where LS means the linear span. Then it holds that \mathcal{D}_L is contained in $\mathcal{C}([0, \infty) \rightarrow [[L^2]]^{(-\alpha)})$ for any $\alpha > 5/6$ and that for $\phi(B(\cdot))$ in \mathcal{D}_L , the $W(t, x)$ -derivative $\partial_{s,x} \phi(B(t))$ is independent of the choice of s in an open interval $(0, t)$. With these property, Itô's formula for elements in \mathcal{D}_L is proved in Theorem:

If $\phi(B(\cdot))$ is in \mathcal{D}_L , then

$$(4.7) \quad \phi(B(t)) - \phi(B(s)) = \int_{\mathbb{R}} \int_s^t \partial_x \phi(B(u)) dB_x(u) dx + \frac{1}{2} \cdot \frac{1}{dx} \cdot \int_s^t \Delta_L \phi(B(u)) du$$

holds for $0 \leq s \leq t$.

Finally, we should like to note that the Lévy's Laplacian Δ_L is involved in the Itô's formula only for generalized Brownian functionals and that Δ_L , in fact, annihilates ordinary Brownian functionals.

§ 2. Preliminaries

1°) Let $S(\mathbb{R}^2)$ be the Schwartz space on \mathbb{R}^2 and $S^*(\mathbb{R}^2)$ be the dual space of $S(\mathbb{R}^2)$. Let μ be the measure of white noise introduced on $S^*(\mathbb{R}^2)$ by the characteristic functional

$$C(\xi) = \exp \left\{ - \frac{1}{2} \|\xi\|^2 \right\}, \quad \xi \in S(\mathbb{R}^2),$$

where $\|\cdot\|$ denotes the $L(\mathbb{R}^2)$ -norm and set $(L^2) = L^2(S^*(\mathbb{R}^2), \mu)$. The Hilbert

space (L^2) admits the Wiener-Itô decomposition

$$(L^2) = \sum_{n=0}^{\infty} \oplus \mathcal{H}_n,$$

where \mathcal{H}_n is the space of n -ple Wiener integrals, i.e.

$$\mathcal{H}_n = \left\{ \int_{\mathbb{R}^{2n}} \cdots \int F(t_1, x_1, \dots, t_n, x_n) W(t_1, x_1) \cdots W(t_n, x_n) dt_1 dx_1 \cdots dt_n dx_n ; \right. \\ \left. F \in \hat{L}^2((\mathbb{R}^2)^n) \right\},$$

the space $\hat{L}^2((\mathbb{R}^2)^n)$ being the totality of symmetric $L^2((\mathbb{R}^2)^n)$ -functions.

The \mathcal{S} -transform of a Brownian functional ϕ in (L^2) is defined by

$$(\mathcal{S}\phi)(\xi) = \int_{S^*(\mathbb{R}^2)} \phi(W + \xi) d\mu(W), \quad \xi \in S(\mathbb{R}^2).$$

It can be easily checked that

$$\mathcal{S}\mathcal{H}_n = \left\{ \int_{\mathbb{R}^{2n}} \cdots \int F(t_1, x_1, \dots, t_n, x_n) \xi(t_1, x_1) \cdots \xi(t_n, x_n) dt_1 dx_1 \cdots dt_n dx_n ; \right. \\ \left. F \in \hat{L}^2((\mathbb{R}^2)^n) \right\}.$$

We denote the space $\mathcal{S}\mathcal{H}_n$ by F_n .

2°) We then come to a background in order to introduce a class of normal functionals of \mathbb{R}^2 -parameter. Take a complete orthonormal system (c.o.n.s.) in $L^2(\mathbb{R}^2)$ formed by

$$\hat{\xi}_{(j,k)} = \xi_j \otimes \xi_k, \quad \xi_j(u) = (2^j j! \sqrt{\pi})^{-1/2} \cdot H_j(u) \cdot e^{-u^2/2}, \quad j, k = 0, 1, 2, \dots,$$

where H_j denotes the Hermite polynomial of degree j . With this c.o.n.s., we introduce a Hilbertian norm $\|\cdot\|_{\alpha,n}$ by

$$\|f\|_{\alpha,n}^2 = \sum_{j_1, k_1, \dots, j_n, k_n=0}^{\infty} \left\{ \prod_{\nu=1}^n (2j_\nu + 1)(2k_\nu + 1) \right\}^\alpha \cdot (f, \hat{\xi}_{(j_1, k_1)} \otimes \cdots \otimes \hat{\xi}_{(j_n, k_n)})^2, \\ f \in L^2((\mathbb{R}^2)^n), \quad \alpha > 0,$$

where (\cdot, \cdot) denotes the $L^2((\mathbb{R}^2)^n)$ -inner product. For $\alpha > 0$ we form Hilbert spaces

$$S_\alpha((\mathbb{R}^2)^n) = \{f \in L^2((\mathbb{R}^2)^n); \|f\|_{\alpha,n} < \infty\}, \\ \hat{S}_\alpha((\mathbb{R}^2)^n) = \{f \in S_\alpha((\mathbb{R}^2)^n); f \text{ is symmetric}\}, \quad \alpha > 0.$$

Let $\hat{S}_{-\alpha}((\mathbb{R}^2)^n)$ be the dual space of $\hat{S}_\alpha((\mathbb{R}^2)^n)$ for $\alpha > 0$. The space $F_n^{(\alpha)}$ of U -functionals is introduced in the same manner as in [2],

$$F_n^{(\alpha)} = \left\{ \dots \int_{\mathbb{R}^{2n}} F(t_1, x_1, \dots, t_n, x_n) \xi(t_1, x_1) \cdots \xi(t_n, x_n) dt_1 dx_1 \cdots dt_n dx_n ; \right. \\ \left. F \in \hat{S}_\alpha((\mathbb{R}^2)^n) \right\}, \quad \alpha > 0.$$

With the help of the \mathcal{S} -transform, we can define a subspace $\mathcal{H}_n^{(\alpha)}$ by

$$\mathcal{H}_n^{(\alpha)} = \mathcal{S}^{-1} F_n^{(\alpha)}.$$

For U_i in $F_n^{(\alpha)}$ with kernel F_i , $i = 1, 2$, we have

$$(U_1, U_2)_{F_n^{(\alpha)}} = n! (F_1, F_2)_{S_\alpha((\mathbb{R}^2)^n)}.$$

This is rephrased in the form

$$(\phi_1, \phi_2)_{\mathcal{H}_n^{(\alpha)}} = (\mathcal{S}\phi_1, \mathcal{S}\phi_2)_{F_n^{(\alpha)}}, \quad \phi_1, \phi_2 \in \mathcal{H}_n^{(\alpha)}.$$

Let $\mathcal{H}_n^{(-\alpha)}$, $\alpha > 0$, be the dual space of $\mathcal{H}_n^{(\alpha)}$, and define the spaces $(L^2)^{(\alpha)} = \sum_{n=0}^\infty \oplus \mathcal{H}_n^{(\alpha)}$ and $(L^2)^{(-\alpha)} = \sum_{n=0}^\infty \oplus \mathcal{H}_n^{(-\alpha)}$ to obtain a Gel'fand triple:

$$(L^2)^{(\alpha)} \subset (L^2) \subset (L^2)^{(-\alpha)}.$$

The \mathcal{S} -transform can be extended to the space $(L^2)^{(-\alpha)}$ to have

$$\mathcal{S}\mathcal{H}_n^{(-\alpha)} = \{ \langle F, \xi^{\otimes n} \rangle ; F \in \hat{S}_{-\alpha}((\mathbb{R}^2)^n) \},$$

which is denoted by $F_n^{(-\alpha)}$.

3° The $W(t, x)$ -derivative $\partial_{t,x}\phi \equiv \partial\phi/\partial W(t, x)$ of a generalized Brownian functional ϕ is defined by

$$\partial_{t,x}\phi = \mathcal{S}^{-1} \frac{\delta}{\delta\xi(t, x)} \mathcal{S}\phi, \quad (t, x) \in \mathbb{R}^2,$$

where $(\delta/\delta\xi(t, x))\mathcal{S}\phi$ denotes the functional derivative of $\mathcal{S}\phi$. If the second variation of the \mathcal{S} -transform $\mathcal{S}\phi$ of ϕ in $(L^2)^{(-\alpha)}$ is given by a following form

$$(\delta^2 \mathcal{S}\phi)_\xi(\eta, \zeta) = \iint_{\mathbb{R}^2} U_1''(\xi; t, x) \eta(t, x) \zeta(t, x) dt dx \\ + \iiint_{\mathbb{R}^4} U_2''(\xi; t, x, s, y) \eta(t, x) \zeta(s, y) dt dx ds dy, \quad \xi, \eta, \zeta \in S(\mathbb{R}^2),$$

then the Lévy's Laplacian Δ_L is defined by

$$\Delta_L \phi = \mathcal{S}^{-1} \left\{ \iint_{\mathbb{R}^2} U_1''(\xi; t, x) dt dx \right\} \quad (\text{see [2], [7] and [8]}).$$

§ 3. The spaces of generalized functionals

In this section, we construct the various spaces of generalized functionals, on which the $W(t, x)$ -differentiation, the operator $1/dx$ and other related notations are introduced.

We introduce the spaces $(\tilde{L}^2)^{(\alpha)}$ and $\tilde{F}^{(\alpha)}$ for every $\alpha \in \mathbf{R}$:

$$\begin{aligned}
 &(\tilde{L}^2)^{(\alpha)} \\
 &= \left\{ \phi = (\phi_1, \phi_2, \dots, \phi_n, \dots); \phi_j \in (L^2)^{(\alpha)}, j = 1, 2, \dots, n, \dots, \sum_{j=1}^{\infty} \|\phi_j\|_{(L^2)^{(\alpha)}}^2 < \infty \right\}, \\
 &\tilde{F}^{(\alpha)} \\
 &= \left\{ f = (f_1, f_2, \dots, f_n, \dots); f_j \in F^{(\alpha)}, j = 1, 2, \dots, n, \dots, \sum_{j=1}^{\infty} \|f_j\|_{F^{(\alpha)}}^2 < \infty \right\}, \\
 &F^{(\alpha)} = \sum_{n=0}^{\infty} \oplus F_n^{(\alpha)}.
 \end{aligned}$$

The spaces $(\tilde{L}^2)^{(\alpha)}$ and $\tilde{F}^{(\alpha)}$ are Hilbert spaces with the inner products

$$(\phi, \psi)_{(\tilde{L}^2)^{(\alpha)}} = \sum_{j=1}^{\infty} (\phi, \psi_j)_{(L^2)^{(\alpha)}}, \quad \phi = (\phi_1, \phi_2, \dots), \quad \psi = (\psi_1, \psi_2, \dots) \in (\tilde{L}^2)^{(\alpha)}$$

and

$$(f, g)_{\tilde{F}^{(\alpha)}} = \sum_{j=1}^{\infty} (f_j, g_j)_{F^{(\alpha)}}, \quad f = (f_1, f_2, \dots), \quad g = (g_1, g_2, \dots) \in \tilde{F}^{(\alpha)}$$

respectively. We define the spaces $(\tilde{L}^2)_*^{(\alpha)}$ and $\tilde{F}_*^{(\alpha)}$ for every $\alpha \in \mathbf{R}$ as follows:

$$\begin{aligned}
 &(\tilde{L}^2)_*^{(\alpha)} = \{ \phi = (\phi_1, \phi_2, \dots) \in (\tilde{L}^2)^{(\alpha)}; \phi_1 = \phi_2 = 0 \}, \\
 &\tilde{F}_*^{(\alpha)} = \{ f = (f_1, f_2, \dots) \in \tilde{F}^{(\alpha)}; f_1 = f_2 = 0 \}.
 \end{aligned}$$

The spaces $(\tilde{L}^2)_*^{(\alpha)}$ and $\tilde{F}_*^{(\alpha)}$ are closed subspaces of $(\tilde{L}^2)^{(\alpha)}$ and $\tilde{F}^{(\alpha)}$ respectively. Set $[[L^2]]^{(\alpha)} = (\tilde{L}^2)^{(\alpha)} / (\tilde{L}^2)_*^{(\alpha)}$ and set $[[F]]^{(\alpha)} = \tilde{F}^{(\alpha)} / \tilde{F}_*^{(\alpha)}$. Both $[[L^2]]^{(\alpha)}$ and $[[F]]^{(\alpha)}$ are Hilbert spaces with the norms

$$\|\phi + (\tilde{L}^2)_*^{(\alpha)}\|_{[[L^2]]^{(\alpha)}} = \inf \{ \|\psi\|_{(L^2)^{(\alpha)}}; \psi \in \phi + (\tilde{L}^2)_*^{(\alpha)} \}, \quad \phi \in (\tilde{L}^2)^{(\alpha)},$$

and

$$\|f + \tilde{F}_*^{(\alpha)}\|_{[[F]]^{(\alpha)}} = \inf \{ \|g\|_{\tilde{F}^{(\alpha)}}; g \in f + \tilde{F}_*^{(\alpha)} \}, \quad f \in \tilde{F}^{(\alpha)},$$

respectively. The spaces $[[L^2]]^{(\alpha)}$ and $[[L^2]]^{(-\alpha)}$ are mutually dual by the canonical bilinear form

$$\begin{aligned}
 &\langle \Phi + (\tilde{L}^2)_*^{(-\alpha)}, \phi + (\tilde{L}^2)_*^{(\alpha)} \rangle_{[[L^2]]^{(-\alpha)}, [[L^2]]^{(\alpha)}} = \langle \Phi_1, \phi_1 \rangle + \langle \Phi_2, \phi_2 \rangle, \\
 &\Phi = (\Phi_1, \Phi_2, \dots) \in (\tilde{L}^2)^{(-\alpha)}, \quad \phi = (\phi_1, \phi_2, \dots) \in (\tilde{L}^2)^{(\alpha)},
 \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ denotes the canonical bilinear form connecting $(L^2)^{(-\alpha)}$ and $(L^2)^{(\alpha)}$. Any element $\phi + (\tilde{L}^2)_*^{(\alpha)}$, with $\phi = (\phi_1, \phi_2, \dots) \in (\tilde{L}^2)^{(\alpha)}$, may be represented as $[[\phi_1, \phi_2]]$. For any $\alpha > 0$, the spaces $[[L^2]]^{(-\alpha)}$ and $[[L^2]]^{(\alpha)}$ are viewed as the space of generalized functionals and the space of testing functionals respectively.

The \mathcal{S} -transform on $[[L^2]]^{(-\alpha)}$, $\alpha > 0$, is given by

$$(3.1) \quad \mathcal{S}[[\phi_1, \phi_2]] = [[\mathcal{S}\phi_1, \mathcal{S}\phi_2]], \quad [[\phi_1, \phi_2]] \in [[L^2]]^{(-\alpha)}.$$

The \mathcal{S} -transform gives an isomorphism $[[L^2]]^{(-\alpha)} \simeq [[F]]^{(-\alpha)}$. The $W(t, x)$ -differentiation $\partial_{t,x} \equiv \partial/\partial W(t, x)$ in $[[L^2]]^{(-\alpha)}$, $\alpha > 0$, is naturally defined by

$$(3.2) \quad \partial_{t,x}[[\phi_1, \phi_2]] = [[\partial_{t,x}\phi_1, \partial_{t,x}\phi_2]]$$

for every differentiable element $[[\phi_1, \phi_2]]$ in $[[L^2]]^{(-\alpha)}$. We now introduce the shift $1/dx$ on $[[L^2]]^{(-\alpha)}$ by the formula

$$(3.3) \quad \frac{1}{dx}[[\phi_1, \phi_2]] = [[0, \phi_1]], \quad [[\phi_1, \phi_2]] \in [[L^2]]^{(-\alpha)}.$$

For $\phi(B(t)) = [[\phi_1(B(t)), \phi_2(B(t))]]$ in $[[L^2]]^{(-\alpha)}$ for some $\alpha > 0$, we understand the integral $\int_s^t \phi(B(u))du$ as

$$(3.4) \quad \int_s^t \phi(B(u))du = \left[\left[\int_s^t \phi_1(B(u))du, \int_s^t \phi_2(B(u))du \right] \right].$$

Similarly, we can define the stochastic integral $\int_s^t \phi(B(u))dB_x(u)$ as

$$(3.5) \quad \int_s^t \phi(B(u))dB_x(u) = \left[\left[\int_s^t \phi_1(B(u))dB_x(u), \int_s^t \phi_2(B(u))dB_x(u) \right] \right].$$

Concerning the first component of (3.5), we can see a similarity to the stochastic integral introduced in [5].

§ 4. Itô's formula and Lévy's Laplacian

We are now in a position to define the domain of the Lévy's Laplacian. The product $B_{x_1}(\cdot)^{p_1} \cdots B_{x_n}(\cdot)^{p_n}$, which has only formal significance, will be understood to be

$$\left[\left[:B_{x_1}(\cdot)^{p_1} \cdots B_{x_n}(\cdot)^{p_n}; \sum_{j=1}^n C_1(p_j)(\cdot); \prod_{\substack{1 \leq \nu \leq n \\ \nu \neq j}} B_{x_\nu}(\cdot)^{p_\nu} B_{x_j}(\cdot)^{p_j-2} \right] \right],$$

where $C_1(p_j) = p_j(p_j - 1)/2$, $j = 1, 2, \dots, n$ and $:B_{x_1}(\cdot)^{p_1} \cdots B_{x_n}(\cdot)^{p_n}$: denotes

the renormalization of $B_{x_1}(\cdot)^{p_1} \dots B_{x_n}(\cdot)^{p_n}$. Then an integral

$$\int_{\mathbb{R}^n} \dots \int_{\mathbb{R}^n} f(x_1, \dots, x_n) B_{x_1}(\cdot)^{p_1} \dots B_{x_n}(\cdot)^{p_n} dx_1 \dots dx_n$$

is given by

$$\left[\int_{\mathbb{R}^n} \dots \int_{\mathbb{R}^n} f(x_1, \dots, x_n) : B_{x_1}(\cdot)^{p_1} \dots B_{x_n}(\cdot)^{p_n} : dx_1 \dots dx_n, \right. \\ \left. \sum_{j=1}^n C_i(p_j)(\cdot) \int_{\mathbb{R}^n} \dots \int_{\mathbb{R}^n} f(x_1, \dots, x_n) : \prod_{\substack{1 \leq \nu \leq n \\ \nu \neq j}} B_{x_\nu}(\cdot)^{p_\nu} B_{x_j}(\cdot)^{p_j-2} : dx_1 \dots dx_n \right].$$

Set

$$\mathcal{D}_L = LS \left\{ \int_{\mathbb{R}^n} \dots \int_{\mathbb{R}^n} f(x_1, \dots, x_n) B_{x_1}(\cdot)^{p_1} \dots B_{x_n}(\cdot)^{p_n} dx_1 \dots dx_n; f \in L^1(\mathbb{R}^n), \right. \\ \left. (p_1, \dots, p_n) \in (N \cup \{0\})^n, n = 0, 1, 2, \dots \right\}.$$

LEMMA 1. We have $\mathcal{D}_L \subset \mathcal{C}([0, \infty) \rightarrow \llbracket L^2 \rrbracket^{(-\alpha)})$ for any $\alpha > 5/6$.

Proof. Take

$$(4.1) \quad \phi(B(\cdot)) = \int_{\mathbb{R}^n} \dots \int_{\mathbb{R}^n} f(x_1, \dots, x_n) B_{x_1}(\cdot)^{p_1} \dots B_{x_n}(\cdot)^{p_n} dx_1 \dots dx_n, \\ f \in L^1(\mathbb{R}^n), \quad p_1 + \dots + p_n = N.$$

It is sufficient to prove $\phi(B(\cdot)) \in \mathcal{C}([0, \infty) \rightarrow \llbracket L^2 \rrbracket^{(-\alpha)})$ for any $\alpha > 5/6$. We will first prove $\phi(B(t)) \in \llbracket L^2 \rrbracket^{(-\alpha)}$ for any $\alpha > 5/6$ and $t \geq 0$. Set

$$F = \int_{\mathbb{R}^n} \dots \int_{\mathbb{R}^n} f(x_1, \dots, x_n) I_{[0,t]}^{\otimes N} \otimes \bigotimes_{\nu=1}^n \delta_{x_\nu}^{\otimes p_\nu} dx_1 \dots dx_n$$

and set

$$G = \sum_{j=1}^n C_i(p_j)t \int_{\mathbb{R}^n} \dots \int_{\mathbb{R}^n} f(x_1, \dots, x_n) I_{[0,t]}^{\otimes (N-2)} \otimes \bigotimes_{\substack{1 \leq \nu \leq n \\ \nu \neq j}} \delta_{x_\nu}^{\otimes p_\nu} \otimes \delta_{x_j}^{\otimes (p_j-2)} dx_1 \dots dx_n.$$

Then what we should prove can be reduced to show that two series

$$(4.2) \quad \sum_{j_1, k_1, \dots, j_N, k_N=0}^{\infty} \left\{ \prod_{\nu=1}^N (2j_\nu + 1)(2k_\nu + 1) \right\}^{-\alpha} \left\langle F, \frac{1}{N!} \sum_{\sigma} \xi_{\sigma(j_1, k_1)} \otimes \dots \otimes \xi_{\sigma(j_N, k_N)} \right\rangle^2$$

and

$$(4.3) \quad \sum_{j_1, k_1, \dots, j_{N-2}, k_{N-2}=0}^{\infty} \left\{ \prod_{\nu=1}^{N-2} (2j_\nu + 1)(2k_\nu + 1) \right\}^{-\alpha} \\ \times \left\langle G, \frac{1}{(N-2)!} \cdot \sum_{\tau} \xi_{\tau(j_1, k_1)} \otimes \dots \otimes \xi_{\tau(j_{N-2}, k_{N-2})} \right\rangle^2$$

converge for any $\alpha > 5/6$, where σ and τ extend over the set of all possible permutations. It is easily checked that

$$(4.4) \quad \left\langle F, \frac{1}{N!} \sum_{\sigma} \xi_{\sigma(j_1, k_1)} \otimes \cdots \otimes \xi_{\sigma(j_N, k_N)} \right\rangle^2 \leq t^{2N} \|f\|_{L^1(\mathbb{R}^n)}^2 \|\xi_{j_1}\|_{\infty}^2 \cdots \|\xi_{j_N}\|_{\infty}^2 \|\xi_{k_1}\|_{\infty}^2 \cdots \|\xi_{k_N}\|_{\infty}^2$$

and

$$(4.5) \quad \left\langle G, \frac{1}{(N-2)!} \cdot \sum_{\tau} \xi_{\tau(j_1, k_1)} \otimes \cdots \otimes \xi_{\tau(j_{N-2}, k_{N-2})} \right\rangle^2 \leq t^{2(n+N-2)} \left\{ \sum_{i,j=1}^n C_1(p_i) C_1(p_j) \right\} \times \|f\|_{L^1(\mathbb{R}^n)}^2 \|\xi_{j_1}\|_{\infty}^2 \cdots \|\xi_{j_{N-2}}\|_{\infty}^2 \|\xi_{k_1}\|_{\infty}^2 \cdots \|\xi_{k_{N-2}}\|_{\infty}^2,$$

where $\|\cdot\|_{L^1(\mathbb{R}^n)}$ is the $L^1(\mathbb{R}^n)$ -norm and $\|\cdot\|_{\infty}$ is the maximum norm. By E. Hille and R. S. Phillips [3], p 571, (21.3.3), it holds that

$$(4.6) \quad \|\xi_j\|_{\infty}^2 = O(j^{-1/6}), \quad j > 0.$$

From (4.4), (4.5) and (4.6), follows the convergence of two series (4.2) and (4.3) for any $\alpha > 5/6$. Next, we prove the continuity of $\phi(B(\cdot))$. Set $\phi(B(\cdot)) = [\phi_1(B(\cdot)), \phi_2(B(\cdot))]$. Then $\|\phi(B(t))\|_{[[L^2]]^{(-\alpha)}}^2 = \|\phi_1(B(t))\|_{(L^2)^{(-\alpha)}}^2 + \|\phi_2(B(t))\|_{(L^2)^{(-\alpha)}}^2$. It is clear that, for any $\alpha > 5/6$ and $0 \leq s \leq t$, $\|\phi_1(B(t)) - \phi_1(B(s))\|_{(L^2)^{(-\alpha)}}^2 \leq N! \sum_{j_1, k_1, \dots, j_N, k_N=0}^{\infty} \{ \prod_{\nu=1}^N (2j_{\nu} + 1)(2k_{\nu} + 1) \}^{-\alpha} (t-s) \{ \text{polynomial in } (t-s) \} \times \|f\|_{L^1(\mathbb{R}^n)}^2 \|\xi_{j_1}\|_{\infty}^2 \cdots \|\xi_{j_N}\|_{\infty}^2 \|\xi_{k_1}\|_{\infty}^2 \cdots \|\xi_{k_N}\|_{\infty}^2$. Similar evaluation can be obtained for $\phi_2(B(t)) - \phi_2(B(s))$. Thus follows the continuity of $\phi(B(\cdot))$.

Q.E.D.

LEMMA 2. For any $\phi(B(\cdot))$ in \mathcal{D}_L , the $W(t, x)$ -derivative $\partial_{s,x}\phi(B(t))$ exists and is independent of the choice of s in the interval $(0, t)$.

Proof. It is sufficient to prove this Lemma for a functional given by (4.1). Set $\Xi(t, x) = \int_0^t \xi(r, x) dr$ for $\xi \in S(\mathbb{R}^2)$. Then by Lemma 1, the \mathcal{L} -transform of $\phi(B(t))$ is given by

$$\mathcal{L}[\phi(B(t))](\xi) = \left[\left[\int_{\mathbb{R}^n} \cdots \int_{\mathbb{R}^n} f(x_1, \dots, x_n) \prod_{\nu=1}^n \Xi(t, x_{\nu})^{p_{\nu}} dx_1 \cdots dx_n, \sum_{j=1}^n C_1(p_j) t \int_{\mathbb{R}^n} \cdots \int_{\mathbb{R}^n} f(x_1, \dots, x_n) \prod_{\substack{1 \leq \nu \leq n \\ \nu \neq j}} \Xi(t, x_{\nu})^{p_{\nu}} \Xi(t, x_j)^{p_j-2} dx_1 \cdots dx_n \right] \right], \quad \xi \in S(\mathbb{R}^2).$$

Hence,

$$\begin{aligned}
 & \frac{\delta}{\delta \xi(s, x)} \mathcal{S}[\phi(B(t))](\xi) \\
 &= \left[\left[\sum_{j=1}^n p_j \Xi(t, x)^{p_j-1} \int_{R^{n-1}} \dots \int f(x_1, \dots, x_{j-1}, x, x_{j+1}, \dots, x_n) \right. \right. \\
 & \quad \times \prod_{\substack{1 \leq \nu \leq n \\ \nu \neq j}} \Xi(t, x_\nu)^{p_\nu} dx_1 \dots dx_{j-1} dx_{j+1} \dots dx_n, \\
 & \quad \sum_{k=1}^n C_1(p_k) t \sum_{\substack{1 \leq j \leq n \\ j \neq k}} p_j \Xi(t, x)^{p_j-1} \int_{R^{n-1}} \dots \int f(x_1, \dots, x_{j-1}, x, x_{j+1}, \dots, x_n) \\
 & \quad \times \prod_{\substack{1 \leq \nu \leq n \\ \nu \neq j, k}} \Xi(t, x_\nu)^{p_\nu} \Xi(t, x_k)^{p_k-2} dx_1 \dots dx_{j-1} dx_{j+1} \dots dx_n \\
 & \quad + \sum_{j=1}^n C_1(p_j) (p_j - 2) t \Xi(t, x)^{p_j-3} \int_{R^{n-1}} \dots \int f(x_1, \dots, x_{j-1}, x, x_{j+1}, \dots, x_n) \\
 & \quad \left. \left. \times \sum_{\substack{1 \leq \nu \leq n \\ \nu \neq j}} \Xi(t, x_\nu)^{p_\nu} dx_1 \dots dx_{j-1} dx_{j+1} \dots dx_n \right] \right], \quad \xi \in S(R^2).
 \end{aligned}$$

By the definition of $\partial_{s,x}\phi(B(t))$ and by the above form, we can see that $\partial_{s,x}\phi(B(t))$ is independent of the choice of s in $(0, t)$. Q.E.D.

By Lemma 2, we may denote $\partial_{s,x}$ simply by ∂_x , when it acts on \mathcal{D}_L .

THEOREM. *If $\phi(B(\cdot))$ is in \mathcal{D}_L , then*

$$(4.7) \quad \phi(B(t)) - \phi(B(s)) = \int_R \int_s^t \partial_x \phi(B(u)) dB_x(u) dx + \frac{1}{2} \cdot \frac{1}{dx} \cdot \int_s^t \Delta_L \phi(B(u)) du$$

holds for $0 \leq s \leq t$.

Proof. It suffices for us to prove (4.7) for an element $\phi(B(t))$ of the form (4.1). The \mathcal{S} -transform of $\partial_x \phi(B(t))$ is given in the proof of Lemma 2. Hence we can easily compute the \mathcal{S} -transform of $\int_R \int_s^t \partial_x \phi(B(u)) dB_x(u) dx$ by the definition of the stochastic integral. The \mathcal{S} -transform of $\frac{1}{dx} \int_s^t \Delta_L \phi(B(u)) du$ is given by

$$\begin{aligned}
 & \mathcal{S} \left[\frac{1}{dx} \int_s^t \Delta_L \phi(B(u)) du \right] (\xi) = \left[\left[0, 2 \sum_{j=1}^n C_1(p_j) \int_{R^n} \dots \int f(x_1, \dots, x_n) \right. \right. \\
 & \quad \left. \left. \times \int_s^t \prod_{\substack{1 \leq \nu \leq n \\ \nu \neq j}} \Xi(u, x_\nu)^{p_\nu} \Xi(u, x_j)^{p_j-2} du dx_1 \dots dx_n \right] \right], \quad \xi \in S(R^2).
 \end{aligned}$$

By comparing $\mathcal{S}[\phi(B(t))](\xi) - \mathcal{S}[\phi(B(s))](\xi)$ with

$$\mathcal{S} \left[\int_R \int_s^t \partial_x \phi(B(u)) dB_x(u) dx \right] (\xi) + \frac{1}{2} \mathcal{S} \left[\frac{1}{dx} \int_s^t \Delta_L \phi(B(u)) du \right] (\xi),$$

we obtain (4.7).

Q.E.D.

REFERENCES

- [1] T. Hida, *Brownian motion in Applications of Mathematics 11*, Springer-Verlag (1980).
- [2] ———, *Brownian motion and its functionals*, *Ricerche di Matematica*, Vol. XXXIV, fasc. 1° (1985).
- [3] E. Hille and R. S. Phillips, *Functional Analysis and Semi-groups*, *Colloq. Publ. Amer. Math. Soc.*, (1957).
- [4] K. Itô, *Foundations of stochastic differential equations in infinite dimensional spaces*, *CBMS-NSF Regional Conference Series in Applied Math.*, **47** (1984).
- [5] I. Kubo, *Ito formula for generalized Brownian functionals*, *Lecture Notes in Control and Information Science*, **49** (1983), 156–166.
- [6] I. Kubo and S. Takenaka, *Calculus on Gaussian white noise, I–IV* *Proc. Japan Acad.*, **56A** (1980), 376–380, 411–416, **57A** (1981), 433–437, **58A** (1982), 186–189.
- [7] H.-H. Kuo, *On Laplacian operators of generalized Brownian functionals*, *Lecture Notes in Mathematics*, **1203**, Springer-Verlag (1986), 119–128.
- [8] P. Lévy, *Problèmes concrets d'analyse fonctionnelle*, Gauthier-Villars (1951).

Department of Mathematics
Nagoya University
Chikusa-ku, Nagoya 464
Japan

Current address:
Department of Mathematics
Meijo University
Tenpaku-ku, Nagoya 468
Japan