



## COMMENTARY

# Capitalizing on natural experiments in low- to middle-income countries to explore epigenetic contributions to disease risk in migrant populations

J. Jaime Miranda<sup>1,2\*</sup>, Caren Weinhouse<sup>3</sup>, Rodrigo M. Carrillo-Larco<sup>1</sup> and Lijing L. Yan<sup>3,4</sup>

<sup>1</sup>CRONICAS Centre of Excellence in Chronic Diseases, Universidad Peruana Cayetano Heredia, Lima, Peru

<sup>2</sup>Department of Medicine, School of Medicine, Universidad Peruana Cayetano Heredia, Lima, Peru

<sup>3</sup>Duke Global Health Institute, Duke University, NC, USA

<sup>4</sup>Duke Kunshan University, Kunshan, China

*Global Health, Epidemiology and Genomics* (2016), 1, e3, page 1 of 3. doi:10.1017/ghg.2015.4

Received 3 July 2015; Revised 15 November 2015; Accepted 24 November 2015

**Key words:** Chronic disease, epigenetics, global health, migration.

Migration poses a significant and worsening public health problem. As the world becomes increasingly interdependent and the global population continues to expand, rates of both within-country and international migration are rising. Migrants tend to experience differential risks for chronic disease, including cardiovascular and metabolic diseases [1–7]. Differential health outcomes in international migrants are not limited to migrants from developing to developed countries; migrants from one developed country to another with regional differences in chronic disease risk may be impacted, as well [8].

Lifestyle factors do not fully explain increased disease risk in some migrant populations. Prior studies have suggested that increases in body mass and blood pressure in migrant populations are related to stress-induced dietary or physical activity changes. These increased risk factors may subsequently influence disease risk [3]. However, individuals that migrated from a subsistence lifestyle on Pacific atoll Tokelau to an urbanized Western lifestyle in New Zealand showed increased blood pressure in men that cannot be fully explained by concomitant dietary changes and weight gain [4]. Migrants often display cardiovascular disease

(CVD) risk intermediate to that of non-migrants in their country of origin and to host population natives [5, 9]. These outcomes suggest that setting of origin, together with initial exposures to such settings, plays a role in acquired disease even in the presence of host population lifestyle factors [5, 9]. Although lifetime risks in migrant groups may approach those of the host population over time, there is evidence for differential health outcomes in migrant populations as compared with non-migrants in studies with relatively long follow-up periods. For example, the Finnish Twins Cohort study reported CVD risk intermediate to that of the migrants' country of origin and of the host population after a 23-year follow-up [5]. Further, in cases in which lifetime risks of migrants do approach the host population over time, the intervening period of differential health is of strong public health interest.

Genetic differences do not fully explain differential disease risk, either. Genetic heterogeneity within a country may contribute to differences in health outcomes between migrants and non-migrants if migration is non-random for genetic markers [8]. However, twins that migrated from Finland to Sweden displayed a higher CVD risk than low-risk native Swedes, but a lower risk than their co-twins in high-risk Finland. These data suggest that differential health by the migration status is strongly influenced by environmental factors [5]. In addition, cardiovascular risk factors in

\* Address for correspondence: J. Jaime Miranda, MD, PhD, CRONICAS Centro de Excelencia en Enfermedades Crónicas, Universidad Peruana Cayetano Heredia, Av. Armendariz 497, Miraflores, Lima 18, Peru. (Email: [jaime.miranda@upch.pe](mailto:jaime.miranda@upch.pe))



rural-to-urban migrants in Peru were dependent on age at first migration [9]. Individuals that migrated when aged older than 12 years were at higher risk for diabetes and metabolic syndrome as compared with individuals that migrated at younger ages [9]. Individuals in both groups are likely genetically similar on the population level and have experienced similar environments, albeit at different life course stages.

Therefore, we propose that epigenetic reprogramming due to early life environment may contribute to differential chronic disease risk in migrant populations. This effect may be strongest in rural-to-urban migrants that experience significantly different environments across the life course. Current evidence supports epigenetic mediation of the link between developmental exposures and metabolic dysfunction and CVD [10], health outcomes with common differential risk in migrant populations as compared with non-migrants.

Epigenetic modifications can cause a change in phenotype with no change in underlying genotype. The epigenome, or genome-wide collection of epigenetic modifications, consists of somatically heritable gene regulatory marks, including DNA methylation, posttranslational histone tail modifications, and chromatin remodeling proteins [11]. The field of environmental epigenetics, or the study of epigenetic responsiveness to the external environment, may partly explain the developmental origins of health and disease, or inter-individual variation in health outcomes in adulthood based on environmental exposures during early life development. [11].

Low- and middle-income countries in the developing world are ideal locations for studying epigenetic contributions to migrant health due to rising urbanization and emigration. Based on our research expertise, we propose Latin America and China as study sites for epigenetic questions in migrant populations. Latin American urbanization rates have risen very quickly in recent decades. In fact, the rural population is now two-thirds smaller than it was in the 1950s [12, 13]. China has experienced similar massive internal migration over the last 30 years, with millions of people relocating from rural to urban areas, offering potentially large study populations [14]. In addition, both Latin America and China are large and topographically diverse regions, allowing for setting-specific studies of rural-to-urban migrants coming from different locations, such as the coast or highlands, and from different altitudes [9, 14]. Lastly, migrants coming to urban settings for socio-economic purposes are not identical to those who have migrated to flee earthquakes and floods or violence [9]. Latin American and China have both experienced historical political unrest and natural disasters that invite studies of multiple motivations for migration [9, 14].

Most environmental epigenetic research has focused on social, nutritional, and chemical exposures, rather than demographic shifts. To the best of our knowledge, only

one published study explores epigenetic profiles of migrants to date, specifically, within-country migrants in Italy [15]. None have been published in the developing world. Therefore, systematic studies of international and within-country migration represent a timely and important opportunity to investigate a potential role for the epigenome in altered chronic disease risk of migrants.

### Acknowledgement

The authors thank Dr Timesh Pilay for critical review of the manuscript.

R.M.C.-L., J.J.M., and the CRONICAS Center of Excellence in Chronic Diseases were supported by the National Heart, Lung, and Blood Institute Global Health Initiative under the contract Global Health Activities in Developing Countries to Combat Non-Communicable Chronic Diseases (Project Number 268200900033C-1-0-1). Support for C.W. was provided by the Inter-American Institute for Global Change Research (CRN#3036).

### Declaration of Interest

None.

### Ethical Standards

No human or animal experimentation involved.

### References

1. **McKay L, MacIntyre S, Ellaway A.** Migration and health: a review of the international literature. Occasional paper No. 12. Glasgow: Medical Research Council, Social & Public Health Sciences Unit, University of Glasgow; 2003.
2. **Dominguez K, et al.** Vital signs: leading causes of death, prevalence of diseases and risk factors, and use of health services among hispanics in the United States – 2009–2013. *Morbidity and Mortality Weekly Report* 2015; **64**: 469–478.
3. **Gadd M, et al.** Do immigrants have an increased prevalence of unhealthy behaviours and risk factors for coronary heart disease? *European Journal of Cardiovascular Prevention and Rehabilitation* 2005; **12**: 535–541.
4. **Salmond CE, Prior IA, Wessen AF.** Blood pressure patterns and migration: a 14-year cohort study of adult Tokelauans. *American Journal of Epidemiology* 1989; **130**: 37–52.
5. **Hedlund E, et al.** Migration and coronary heart disease: a study of Finnish twins living in Sweden and their co-twins residing in Finland. *Scandinavian Journal of Public Health* 2007; **35**: 468–474.
6. **Poulter NR, et al.** Migration-induced changes in blood pressure: a controlled longitudinal study. *Clinical and Experimental Pharmacology and Physiology*; **12**: 211–216.



7. **Unwin N, et al.** Rural to urban migration and changes in cardiovascular risk factors in Tanzania: a prospective cohort study. *BMC Public Health* 2010; **10**: 272.
8. **Cobbaert CM, et al.** Regional differences of HFE (C282Y, H63D) allele frequencies in the Netherlands: a model case illustrating the significance of genographics and prehistorical population migration. *Acta Clinica Belgica*; **67**: 430–435.
9. **Miranda JJ, Gilman RH, Smeeth L.** Differences in cardiovascular risk factors in rural, urban and rural-to-urban migrants in Peru. *Heart* 2011; **97**: 787–796.
10. **Kelishadi R, Poursafa P.** A review on the genetic, environmental, and lifestyle aspects of the early-life origins of cardiovascular disease. *Current Problems in Pediatric and Adolescent Health Care* 2014; **44**: 54–72.
11. **Hochberg Z, et al.** Child health, developmental plasticity, and epigenetic programming. *Endocrine Review* 2011; **32**: 159–224.
12. World Urbanization Prospects, the 2014 Revision. 2014. ([www.esa.un.org/unpd/wup](http://www.esa.un.org/unpd/wup)). Accessed on December 9, 2015.
13. **Spence M, Annez PC, Buckley RM.** Urbanization and Growth. 2009. ([www.openknowledge.worldbank.org](http://www.openknowledge.worldbank.org)). Accessed on December 9, 2015.
14. **Qiu P, et al.** Rural-to-urban migration and its implication for new cooperative medical scheme coverage and utilization in China. *BMC Public Health* 2011; **11**: 520.
15. **Campanella G, et al.** Epigenetic signatures of internal migration in Italy. *International Journal of Epidemiology* 2014; **44**: 1442–1449.