Another Criterion For The Riemann Hypothesis

Frank Vega

CopSonic, 1471 Route de Saint-Nauphary 82000 Montauban, France

Abstract

Let's define $\delta(x) = (\sum_{q \leq x} \frac{1}{q} - \log \log x - B)$, where $B \approx 0.2614972128$ is the Meissel-Mertens constant. The Robin theorem states that $\delta(x)$ changes sign infinitely often. Let's also define $S(x) = \theta(x) - x$, where $\theta(x)$ is the Chebyshev function. It is known that $S(x)$ changes sign infinitely often. We define the another function $\varpi(x) = (\sum_{q \leq x} \frac{1}{q} - \log \log \theta(x) - B)$. We prove that when the inequality $\varpi(x) \leq 0$ is satisfied for some number $x \geq 3$, then the Riemann Hypothesis should be false. The Riemann Hypothesis is also false when the inequalities $\delta(x) \leq 0$ and $S(x) \geq 0$ are satisfied for some number $x \geq 3$ or when $\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}} + \frac{\log x}{\log \theta(x)} \leq 1$ is satisfied for some number $x \geq 13.1$.

Keywords: Riemann hypothesis, Nicolas inequality, Chebyshev function, prime numbers
2000 MSC: 11M26, 11A41, 11A25

1. Introduction

In mathematics, the Riemann Hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$ [1]. Let $N_n = 2 \times 3 \times 5 \times 7 \times 11 \times \dots \times p_n$ denotes a primorial number of order n such that p_n is the n^{th} prime number. Say $\text{Nicolas}(p_n)$ holds provided

$$\prod_{q|N_n} \frac{q}{q-1} > e^\gamma \times \log \log N_n.$$

The constant $\gamma \approx 0.57721$ is the Euler-Mascheroni constant, $\log$ is the natural logarithm, and $q | N_n$ means the prime number q divides to N_n . The importance of this property is:

Theorem 1.1. [2], [3]. $\text{Nicolas}(p_n)$ holds for all prime numbers $p_n > 2$ if and only if the Riemann Hypothesis is true.

In mathematics, the Chebyshev function $\theta(x)$ is given by

$$\theta(x) = \sum_{p \leq x} \log p$$

where $p \leq x$ means all the prime numbers p that are less than or equal to x . We know this:

Theorem 1.2. [4].

$$\lim_{x \rightarrow \infty} \frac{\theta(x)}{x} = 1.$$

Let's define $S(x) = \theta(x) - x$. It is a known result that:

Theorem 1.3. [5]. $S(x)$ changes sign infinitely often.

We also know that

Theorem 1.4. [6]. If the Riemann hypothesis holds, then

$$\left(\frac{e^{-\gamma}}{\log x} \times \prod_{q \leq x} \frac{q}{q-1} - 1 \right) < \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}$$

for all numbers $x \geq 13.1$.

Let's define $H = \gamma - B$ such that $B \approx 0.2614972128$ is the Meissel-Mertens constant [7]. We know from the constant H , the following formula:

Theorem 1.5. [8].

$$\sum_q \left(\log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right) = \gamma - B = H.$$

For $x \geq 2$, the function $u(x)$ is defined as follows

$$u(x) = \sum_{q > x} \left(\log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right).$$

Nicolas showed that

Theorem 1.6. [3]. For $x \geq 2$:

$$0 < u(x) \leq \frac{1}{2 \times (x-1)}.$$

Let's define:

$$\delta(x) = \left(\sum_{q \leq x} \frac{1}{q} - \log \log x - B \right).$$

Robin theorem states the following result:

Theorem 1.7. [9]. $\delta(x)$ changes sign infinitely often.

In addition, the Mertens second theorem states that:

Theorem 1.8. [7].

$$\lim_{x \rightarrow \infty} \delta(x) = 0.$$

Besides, we use the following theorem

Theorem 1.9. [10]. For $x > -1$:

$$\frac{x}{x+1} \leq \log(1+x) \leq x.$$

We define another function:

$$\varpi(x) = \left(\sum_{q \leq x} \frac{1}{q} - \log \log \theta(x) - B \right).$$

Putting all together yields the proof that the inequality $\varpi(x) > u(x)$ is satisfied for a number $x \geq 3$ if and only if $\text{Nicolas}(p)$ holds, where p is the greatest prime number such that $p \leq x$. In this way, we introduce another criterion for the Riemann Hypothesis based on the Nicolas criterion and deduce some of its consequences.

2. Results

Theorem 2.1. *The inequality $\varpi(x) > u(x)$ is satisfied for a number $x \geq 3$ if and only if $\text{Nicolas}(p)$ holds, where p is the greatest prime number such that $p \leq x$.*

Proof. We start from the inequality:

$$\varpi(x) > u(x)$$

which is equivalent to

$$\left(\sum_{q \leq x} \frac{1}{q} - \log \log \theta(x) - B \right) > \sum_{q > x} \left(\log \left(\frac{q}{q-1} \right) - \frac{1}{q} \right).$$

Let's add the following formula to the both sides of the inequality,

$$\sum_{q \leq x} \left(\log \left(\frac{q}{q-1} \right) - \frac{1}{q} \right)$$

and due to the theorem 1.5, we obtain that

$$\sum_{q \leq x} \log \left(\frac{q}{q-1} \right) - \log \log \theta(x) - B > H$$

because of

$$H = \sum_{q \leq x} \left(\log \left(\frac{q}{q-1} \right) - \frac{1}{q} \right) + \sum_{q > x} \left(\log \left(\frac{q}{q-1} \right) - \frac{1}{q} \right)$$

and

$$\sum_{q \leq x} \log \left(\frac{q}{q-1} \right) = \sum_{q \leq x} \frac{1}{q} + \sum_{q \leq x} \left(\log \left(\frac{q}{q-1} \right) - \frac{1}{q} \right).$$

Let's distribute it and remove B from the both sides:

$$\sum_{q \leq x} \log \left(\frac{q}{q-1} \right) > \gamma + \log \log \theta(x)$$

since $H = \gamma - B$. If we apply the exponentiation to the both sides of the inequality, then we have that

$$\prod_{q \leq x} \frac{q}{q-1} > e^\gamma \times \log \theta(x)$$

which means that $\text{Nicolas}(p)$ holds, where p is the greatest prime number such that $p \leq x$. The same happens in the reverse implication. $\square$

Theorem 2.2. *The Riemann Hypothesis is true if and only if the inequality $\varpi(x) > u(x)$ is satisfied for all numbers $x \geq 3$.*

Proof. This is a direct consequence of theorems 1.1 and 2.1. $\square$

Theorem 2.3. *If the inequality $\varpi(x) \leq 0$ is satisfied for some number $x \geq 3$, then the Riemann Hypothesis should be false.*

Proof. This is an implication of theorems 1.6, 2.1 and 2.2. $\square$

Theorem 2.4. *If the inequalities $\delta(x) \leq 0$ and $S(x) \geq 0$ are satisfied for some number $x \geq 3$, then the Riemann Hypothesis should be false.*

Proof. If the inequalities $\delta(x) \leq 0$ and $S(x) \geq 0$ are satisfied for some number $x \geq 3$, then we obtain that $\varpi(x) \leq 0$ is also satisfied, which means that the Riemann Hypothesis should be false according to the theorem 2.3. $\square$

Theorem 2.5.

$$\lim_{x \rightarrow \infty} \varpi(x) = 0.$$

Proof. We know that $\lim_{x \rightarrow \infty} \varpi(x) = 0$ for the limits $\lim_{x \rightarrow \infty} \delta(x) = 0$ and $\lim_{x \rightarrow \infty} \frac{\theta(x)}{x} = 1$. In this way, this is a consequence from the theorems 1.8 and 1.2. $\square$

Theorem 2.6. *If the Riemann hypothesis holds, then*

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}} + \frac{\log x}{\log \theta(x)} > 1$$

for all numbers $x \geq 13.1$.

Proof. Under the assumption that the Riemann hypothesis is true, then we would have

$$\prod_{q \leq x} \frac{q}{q-1} < e^\gamma \times \log x \times \left(1 + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}\right)$$

after of distributing the terms based on the theorem 1.4 for all numbers $x \geq 13.1$. If we apply the logarithm to the both sides of the previous inequality, then we obtain that

$$\sum_{q \leq x} \log\left(\frac{q}{q-1}\right) < \gamma + \log \log x + \log\left(1 + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}\right).$$

That would be equivalent to

$$\sum_{q \leq x} \frac{1}{q} + \sum_{q \leq x} \left(\log\left(\frac{q}{q-1}\right) - \frac{1}{q}\right) < \gamma + \log \log x + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}$$

where we know that

$$\log\left(1 + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}\right) \leq \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}$$

according to theorem 1.9. We use the theorems 1.5 and 1.6 to show that

$$\sum_{q \leq x} \left(\log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right) = H - u(x)$$

and $\gamma = H + B$. So,

$$H - u(x) < H + B + \log \log x - \sum_{q \leq x} \frac{1}{q} + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}$$

which is the same as

$$H - u(x) < H - \delta(x) + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}.$$

We eliminate the value of H and thus,

$$-u(x) < -\delta(x) + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}$$

which is equal to

$$u(x) + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}} > \delta(x).$$

We know from the theorem 2.1 that $\varpi(x) > u(x)$ and therefore,

$$\varpi(x) + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}} > \delta(x).$$

Hence,

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}} > \log \log \theta(x) - \log \log x.$$

Suppose that $\theta(x) = \epsilon \times x$ for some constant $\epsilon > 0$. Then,

$$\begin{aligned} \log \log \theta(x) - \log \log x &= \log \log (\epsilon \times x) - \log \log x \\ &= \log (\log x + \log \epsilon) - \log \log x \\ &= \log \left(\log x \times \left(1 + \frac{\log \epsilon}{\log x} \right) \right) - \log \log x \\ &= \log \log x + \log \left(1 + \frac{\log \epsilon}{\log x} \right) - \log \log x \\ &= \log \left(1 + \frac{\log \epsilon}{\log x} \right). \end{aligned}$$

In addition, we know that

$$\log \left(1 + \frac{\log \epsilon}{\log x} \right) \geq \frac{\log \epsilon}{\log \theta(x)}$$

using the theorem 1.9. Certainly, we will have that

$$\log \left(1 + \frac{\log \epsilon}{\log x} \right) \geq \frac{\frac{\log \epsilon}{\log x}}{\frac{\log \epsilon}{\log x} + 1} = \frac{\log \epsilon}{\log \epsilon + \log x} = \frac{\log \epsilon}{\log \theta(x)}.$$

Thus,

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}} > \frac{\log \epsilon}{\log \theta(x)}.$$

If we add the following value of $\frac{\log x}{\log \theta(x)}$ to the both sides of the inequality, then

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}} + \frac{\log x}{\log \theta(x)} > \frac{\log \epsilon}{\log \theta(x)} + \frac{\log x}{\log \theta(x)} = \frac{\log \epsilon + \log x}{\log \theta(x)} = \frac{\log \theta(x)}{\log \theta(x)} = 1.$$

Therefore, the proof is done. $\square$

Theorem 2.7. *If the inequality $\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}} + \frac{\log x}{\log \theta(x)} \leq 1$ is satisfied for some number $x \geq 13.1$, then the Riemann Hypothesis should be false.*

Proof. This is a direct consequence of theorem 2.6. $\square$

References

- [1] P. B. Borwein, S. Choi, B. Rooney, A. Weirathmueller, The Riemann Hypothesis: A Resource for the Afficionado and Virtuoso Alike, Vol. 27, Springer Science & Business Media, 2008.
- [2] J.-L. Nicolas, Petites valeurs de la fonction d'Euler et hypothese de Riemann, Séminaire de Théorie des nombres DPP, Paris 82 (1981) 207–218.
- [3] J.-L. Nicolas, Petites valeurs de la fonction d'Euler, Journal of number theory 17 (3) (1983) 375–388. doi:10.1016/0022-314X(83)90055-0.
- [4] T. H. Grönwall, Some asymptotic expressions in the theory of numbers, Transactions of the American Mathematical Society 14 (1) (1913) 113–122. doi:10.2307/1988773.
- [5] D. J. Platt, T. S. Trudgian, On the first sign change of $\theta(x) - x$, Math. Comput. 85 (299) (2016) 1539–1547. doi:10.1090/mcom/3021.
- [6] J. B. Rosser, L. Schoenfeld, Sharper Bounds for the Chebyshev Functions $\theta(x)$ and $\psi(x)$, Mathematics of computation (1975) 243–269 doi:10.1090/S0025-5718-1975-0457373-7.
- [7] F. Mertens, Ein Beitrag zur analytischen Zahlentheorie., J. reine angew. Math. 1874 (78) (1874) 46–62. doi:10.1515/crll.1874.78.46. URL <https://doi.org/10.1515/crll.1874.78.46>
- [8] Y. Choie, N. Lichiardopol, P. Moree, P. Solé, On Robin's criterion for the Riemann hypothesis, Journal de Théorie des Nombres de Bordeaux 19 (2) (2007) 357–372. doi:10.5802/jtnb.591.
- [9] G. Robin, Sur l'ordre maximum de la fonction somme des diviseurs, Séminaire Delange-Pisot-Poitou Paris 82 (1981) 233–242.
- [10] L. Kozma, Useful Inequalities, http://www.lkozma.net/inequalities_cheat_sheet/ineq.pdf, accessed on 2021-10-08 (2021).