

# Another Criterion For The Riemann Hypothesis

Frank Vega

CopSonic, 1471 Route de Saint-Nauphary 82000 Montauban, France

---

## Abstract

Let's define  $\delta(x) = (\sum_{q \leq x} \frac{1}{q} - \log \log x - B)$ , where  $B \approx 0.2614972128$  is the Meissel-Mertens constant. The Robin theorem states that  $\delta(x)$  changes sign infinitely often. Let's also define  $S(x) = \theta(x) - x$ , where  $\theta(x)$  is the Chebyshev function. It is known that  $S(x)$  changes sign infinitely often. We define the another function  $\varpi(x) = (\sum_{q \leq x} \frac{1}{q} - \log \log \theta(x) - B)$ . We prove that when the inequality  $\varpi(x) \leq 0$  is satisfied for some number  $x \geq 3$ , then the Riemann hypothesis should be false. The Riemann hypothesis is also false when the inequalities  $\delta(x) \leq 0$  and  $S(x) \geq 0$  are satisfied for some number  $x \geq 3$  or when  $\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} + \frac{\log x}{\log \theta(x)} \leq 1$  is satisfied for some number  $x \geq 13.1$ .

**Keywords:** Riemann hypothesis, Nicolas inequality, Chebyshev function, prime numbers  
**2000 MSC:** 11M26, 11A41, 11A25

---

## 1. Introduction

In mathematics, the Riemann hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part  $\frac{1}{2}$  [1]. Let  $N_n = 2 \times 3 \times 5 \times 7 \times 11 \times \dots \times p_n$  denotes a primorial number of order  $n$  such that  $p_n$  is the  $n^{th}$  prime number. Say Nicolas( $p_n$ ) holds provided

$$\prod_{q|N_n} \frac{q}{q-1} > e^\gamma \times \log \log N_n.$$

The constant  $\gamma \approx 0.57721$  is the Euler-Mascheroni constant,  $\log$  is the natural logarithm, and  $q | N_n$  means the prime number  $q$  divides to  $N_n$ . The importance of this property is:

**Theorem 1.1.** [2]. Nicolas( $p_n$ ) holds for all prime numbers  $p_n > 2$  if and only if the Riemann hypothesis is true.

In mathematics, the Chebyshev function  $\theta(x)$  is given by

$$\theta(x) = \sum_{p \leq x} \log p$$

where  $p \leq x$  means all the prime numbers  $p$  that are less than or equal to  $x$ . We know this property for this function:

**Theorem 1.2.** [3].

$$\lim_{x \rightarrow \infty} \frac{\theta(x)}{x} = 1.$$

Let's define  $S(x) = \theta(x) - x$ . It is a known result that:

**Theorem 1.3.** [4].  $S(x)$  changes sign infinitely often.

We also know that

**Theorem 1.4.** [5]. If the Riemann hypothesis holds, then

$$\left( \frac{e^{-\gamma}}{\log x} \times \prod_{q \leq x} \frac{q}{q-1} - 1 \right) < \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}$$

for all numbers  $x \geq 13.1$ .

Let's define  $H = \gamma - B$  such that  $B \approx 0.2614972128$  is the Meissel-Mertens constant [6]. We know from the constant  $H$ , the following formula:

**Theorem 1.5.** [7].

$$\sum_q \left( \log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right) = \gamma - B = H.$$

For  $x \geq 2$ , the function  $u(x)$  is defined as follows

$$u(x) = \sum_{q > x} \left( \log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right).$$

Nicolas showed that

**Theorem 1.6.** [2]. For  $x \geq 2$ :

$$0 < u(x) \leq \frac{1}{2 \times (x-1)}.$$

Let's define:

$$\delta(x) = \left( \sum_{q \leq x} \frac{1}{q} - \log \log x - B \right).$$

Robin theorem states the following result:

**Theorem 1.7.** [8].  $\delta(x)$  changes sign infinitely often.

In addition, the Mertens second theorem states that:

**Theorem 1.8.** [6].

$$\lim_{x \rightarrow \infty} \delta(x) = 0.$$

Besides, we use the following theorems:

**Theorem 1.9.** [9]. For  $x > -1$ :

$$\frac{x}{x+1} \leq \frac{\log(1+x)}{2}.$$

**Theorem 1.10.** [10]. For  $x \geq 1$ :

$$\log\left(1 + \frac{1}{x}\right) < \frac{1}{x + 0.4}.$$

We define another function:

$$\varpi(x) = \left( \sum_{q \leq x} \frac{1}{q} - \log \log \theta(x) - B \right).$$

Putting all together yields the proof that the inequality  $\varpi(x) > u(x)$  is satisfied for a number  $x \geq 3$  if and only if  $\text{Nicolas}(p)$  holds, where  $p$  is the greatest prime number such that  $p \leq x$ . In this way, we introduce another criterion for the Riemann hypothesis based on the Nicolas criterion and deduce some of its consequences.

## 2. Results

**Theorem 2.1.** *The inequality  $\varpi(x) > u(x)$  is satisfied for a number  $x \geq 3$  if and only if  $\text{Nicolas}(p)$  holds, where  $p$  is the greatest prime number such that  $p \leq x$ .*

*Proof.* We start from the inequality:

$$\varpi(x) > u(x)$$

which is equivalent to

$$\left( \sum_{q \leq x} \frac{1}{q} - \log \log \theta(x) - B \right) > \sum_{q > x} \left( \log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right).$$

Let's add the following formula to the both sides of the inequality,

$$\sum_{q \leq x} \left( \log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right)$$

and due to the theorem 1.5, we obtain that

$$\sum_{q \leq x} \log\left(\frac{q}{q-1}\right) - \log \log \theta(x) - B > H$$

because of

$$H = \sum_{q \leq x} \left( \log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right) + \sum_{q > x} \left( \log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right)$$

and

$$\sum_{q \leq x} \log\left(\frac{q}{q-1}\right) = \sum_{q \leq x} \frac{1}{q} + \sum_{q \leq x} \left( \log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right).$$

Let's distribute it and remove  $B$  from the both sides:

$$\sum_{q \leq x} \log\left(\frac{q}{q-1}\right) > \gamma + \log \log \theta(x)$$

since  $H = \gamma - B$ . If we apply the exponentiation to the both sides of the inequality, then we have that

$$\prod_{q \leq x} \frac{q}{q-1} > e^\gamma \times \log \theta(x)$$

which means that  $\text{Nicolas}(p)$  holds, where  $p$  is the greatest prime number such that  $p \leq x$ . The same happens in the reverse implication.  $\square$

**Theorem 2.2.** *The Riemann hypothesis is true if and only if the inequality  $\varpi(x) > u(x)$  is satisfied for all numbers  $x \geq 3$ .*

*Proof.* This is a direct consequence of theorems 1.1 and 2.1.  $\square$

**Theorem 2.3.** *If the inequality  $\varpi(x) \leq 0$  is satisfied for some number  $x \geq 3$ , then the Riemann hypothesis should be false.*

*Proof.* This is an implication of theorems 1.6, 2.1 and 2.2.  $\square$

**Theorem 2.4.** *If the inequalities  $\delta(x) \leq 0$  and  $S(x) \geq 0$  are satisfied for some number  $x \geq 3$ , then the Riemann hypothesis should be false.*

*Proof.* If the inequalities  $\delta(x) \leq 0$  and  $S(x) \geq 0$  are satisfied for some number  $x \geq 3$ , then we obtain that  $\varpi(x) \leq 0$  is also satisfied, which means that the Riemann hypothesis should be false according to the theorem 2.3.  $\square$

**Theorem 2.5.**

$$\lim_{x \rightarrow \infty} \varpi(x) = 0.$$

*Proof.* We know that  $\lim_{x \rightarrow \infty} \varpi(x) = 0$  for the limits  $\lim_{x \rightarrow \infty} \delta(x) = 0$  and  $\lim_{x \rightarrow \infty} \frac{\theta(x)}{x} = 1$ . In this way, this is a consequence from the theorems 1.8 and 1.2.  $\square$

**Theorem 2.6.** *If the Riemann hypothesis holds, then*

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} + \frac{\log x}{\log \theta(x)} > 1$$

for all numbers  $x \geq 13.1$ .

*Proof.* Under the assumption that the Riemann hypothesis is true, then we would have

$$\prod_{q \leq x} \frac{q}{q-1} < e^\gamma \times \log x \times \left(1 + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}\right)$$

after of distributing the terms based on the theorem 1.4 for all numbers  $x \geq 13.1$ . If we apply the logarithm to the both sides of the previous inequality, then we obtain that

$$\sum_{q \leq x} \log\left(\frac{q}{q-1}\right) < \gamma + \log \log x + \log\left(1 + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}\right).$$

That would be equivalent to

$$\sum_{q \leq x} \frac{1}{q} + \sum_{q \leq x} \left(\log\left(\frac{q}{q-1}\right) - \frac{1}{q}\right) < \gamma + \log \log x + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2}$$

where we know that

$$\begin{aligned} \log\left(1 + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x}}\right) &< \frac{1}{\frac{8 \times \pi \times \sqrt{x}}{3 \times \log x + 5} + 0.4} \\ &= \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 0.4 \times (3 \times \log x + 5)} \\ &= \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} \end{aligned}$$

according to theorem 1.10 since  $\frac{8 \times \pi \times \sqrt{x}}{3 \times \log x + 5} \geq 1$  for all numbers  $x \geq 13.1$ . We use the theorems 1.5 and 1.6 to show that

$$\sum_{q \leq x} \left( \log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right) = H - u(x)$$

and  $\gamma = H + B$ . So,

$$H - u(x) < H + B + \log \log x - \sum_{q \leq x} \frac{1}{q} + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2}$$

which is the same as

$$H - u(x) < H - \delta(x) + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2}.$$

We eliminate the value of  $H$  and thus,

$$-u(x) < -\delta(x) + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2}$$

which is equal to

$$u(x) + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} > \delta(x).$$

We know from the theorem 2.1 that  $\varpi(x) > u(x)$  for all numbers  $x \geq 13.1$  and therefore,

$$\varpi(x) + \frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} > \delta(x).$$

Hence,

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} > \log \log \theta(x) - \log \log x.$$

Suppose that  $\theta(x) = \epsilon \times x$  for some constant  $\epsilon > 1$ . Then,

$$\begin{aligned} \log \log \theta(x) - \log \log x &= \log \log(\epsilon \times x) - \log \log x \\ &= \log(\log x + \log \epsilon) - \log \log x \\ &= \log\left(\log x \times \left(1 + \frac{\log \epsilon}{\log x}\right)\right) - \log \log x \\ &= \log \log x + \log\left(1 + \frac{\log \epsilon}{\log x}\right) - \log \log x \\ &= \log\left(1 + \frac{\log \epsilon}{\log x}\right). \end{aligned}$$

In addition, we know that

$$\log\left(1 + \frac{\log \epsilon}{\log x}\right) \geq \frac{\log \epsilon}{\log \theta(x)}$$

using the theorem 1.9 since  $\frac{\log \epsilon}{\log x} > -1$  when  $\epsilon > 1$ . Certainly, we will have that

$$\log\left(1 + \frac{\log \epsilon}{\log x}\right) \geq \frac{\frac{\log \epsilon}{\log x}}{\frac{\log \epsilon}{\log x} + 1} = \frac{\log \epsilon}{\log \epsilon + \log x} = \frac{\log \epsilon}{\log \theta(x)}.$$

Thus,

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} > \frac{\log \epsilon}{\log \theta(x)}.$$

If we add the following value of  $\frac{\log x}{\log \theta(x)}$  to the both sides of the inequality, then

$$\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} + \frac{\log x}{\log \theta(x)} > \frac{\log \epsilon}{\log \theta(x)} + \frac{\log x}{\log \theta(x)} = \frac{\log \epsilon + \log x}{\log \theta(x)} = \frac{\log \theta(x)}{\log \theta(x)} = 1.$$

We know this inequality is satisfied when  $0 < \epsilon \leq 1$  since we would obtain that  $\frac{\log x}{\log \theta(x)} \geq 1$ . Therefore, the proof is done.  $\square$

**Theorem 2.7.** *If the inequality  $\frac{3 \times \log x + 5}{8 \times \pi \times \sqrt{x} + 1.2 \times \log x + 2} + \frac{\log x}{\log \theta(x)} \leq 1$  is satisfied for some number  $x \geq 13.1$ , then the Riemann hypothesis should be false.*

*Proof.* This is a direct consequence of theorem 2.6.  $\square$

## References

- [1] P. B. Borwein, S. Choi, B. Rooney, A. Weirathmueller, The Riemann Hypothesis: A Resource for the Afficionado and Virtuoso Alike, Vol. 27, Springer Science & Business Media, 2008.
- [2] J.-L. Nicolas, Petites valeurs de la fonction d'Euler, Journal of number theory 17 (3) (1983) 375–388. doi:10.1016/0022-314X(83)90055-0.
- [3] T. H. Grönwall, Some asymptotic expressions in the theory of numbers, Transactions of the American Mathematical Society 14 (1) (1913) 113–122. doi:10.2307/1988773.
- [4] D. J. Platt, T. S. Trudgian, On the first sign change of  $\theta(x) - x$ , Math. Comput. 85 (299) (2016) 1539–1547. doi:10.1090/mcom/3021.
- [5] J. B. Rosser, L. Schoenfeld, Sharper Bounds for the Chebyshev Functions  $\theta(x)$  and  $\psi(x)$ , Mathematics of computation (1975) 243–269 doi:10.1090/S0025-5718-1975-0457373-7.
- [6] F. Mertens, Ein Beitrag zur analytischen Zahlentheorie., J. reine angew. Math. 1874 (78) (1874) 46–62. doi:10.1515/crll.1874.78.46. URL <https://doi.org/10.1515/crll.1874.78.46>
- [7] Y. Choie, N. Lichiardopol, P. Moree, P. Solé, On Robin's criterion for the Riemann hypothesis, Journal de Théorie des Nombres de Bordeaux 19 (2) (2007) 357–372. doi:10.5802/jtnb.591.
- [8] G. Robin, Sur l'ordre maximum de la fonction somme des diviseurs, Séminaire Delange-Pisot-Poitou Paris 82 (1981) 233–242.
- [9] L. Kozma, Useful Inequalities, [http://www.lkozma.net/inequalities\\_cheat\\_sheet/ineq.pdf](http://www.lkozma.net/inequalities_cheat_sheet/ineq.pdf), accessed on 2021-10-13 (2021).
- [10] A. Ghosh, An Asymptotic Formula for the Chebyshev Theta Function, arXiv preprint arXiv:1902.09231.