

# Slide Presentation on the Riemann Hypothesis

## Submitted For Short Communications Satellite 2022

Frank Vega, CopSonic France

vega.frank@gmail.com

June 21, 2022

## Abstract

The Riemann Hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part  $\frac{1}{2}$ . In 2011, Patrick Solé and Michel Planat stated a new criterion for the Riemann Hypothesis. We prove **the Riemann Hypothesis is true** using this criterion.

## Keywords

Riemann Hypothesis, Prime numbers, Chebyshev function, Riemann zeta function.

The Chebyshev function  $\theta(x)$  is given by

$$\theta(x) = \sum_{p \leq x} \log p$$

with the sum extending over all prime numbers  $p$  that are less than or equal to  $x$ , where  $\log$  is the natural logarithm. We provide a proof for the Riemann Hypothesis using the properties of the Chebyshev function.

Say  $\text{Dedekinds}(q_n)$  holds provided

$$\prod_{q \leq q_n} \left(1 + \frac{1}{q}\right) > \frac{e^\gamma}{\zeta(2)} \times \log \theta(q_n)$$

where the constant  $\gamma \approx 0.57721$  is the Euler-Mascheroni constant,  $q_n$  is the  $n$ th prime number and  $\zeta(x)$  is the Riemann zeta function.

## Theorem 1

*Dedekinds( $q_n$ ) holds for all prime numbers  $q_n > 3$  if and only if the Riemann Hypothesis is true (Solé and Planat, 2011).*

# What if the Riemann Hypothesis were false?

## Theorem 2

*If the Riemann Hypothesis is false, then there are infinitely many prime numbers  $q_n$  for which  $\text{Dedekinds}(q_n)$  does not hold.*

The Riemann Hypothesis is false, if there exists some natural number  $x_0 \geq 5$  such that  $g(x_0) > 1$  or equivalent  $\log g(x_0) > 0$  (Solé and Planat, 2011):

$$g(x) = \frac{e^\gamma}{\zeta(2)} \times \log \theta(x) \times \prod_{q \leq x} \left(1 + \frac{1}{q}\right)^{-1}.$$

We know the bound (Solé and Planat, 2011):

$$\log g(x) \geq \log f(x) - \frac{2}{x}$$

where  $f$  is introduced in the Nicolas paper (Nicolas, 1983):

$$f(x) = e^\gamma \times \log \theta(x) \times \prod_{q \leq x} \left(1 - \frac{1}{q}\right).$$

## Remark

When the Riemann Hypothesis is false, then there exists a real number  $b < \frac{1}{2}$  for which there are infinitely many natural numbers  $x$  such that  $\log f(x) = \Omega_+(x^{-b})$  (Nicolas, 1983).

According to the Hardy and Littlewood definition, this would mean that

$$\exists k > 0, \forall y_0 \in \mathbb{N}, \exists y \in \mathbb{N} > y_0: \log f(y) \geq k \times y^{-b}.$$

That inequality is equivalent to  $\log f(y) \geq (k \times y^{-b} \times \sqrt{y}) \times \frac{1}{\sqrt{y}}$ , but we note that

$$\lim_{y \rightarrow \infty} (k \times y^{-b} \times \sqrt{y}) = \infty$$

for every possible positive value of  $k$  when  $b < \frac{1}{2}$ .



In this way, this implies that

$$\exists k > 0, \forall y_0 \in \mathbb{N}, \exists y \in \mathbb{N} > y_0: \log f(y) \geq \frac{1}{\sqrt{y}}.$$

Hence, if the Riemann Hypothesis is false, then there are infinitely many natural numbers  $x$  such that  $\log f(x) \geq \frac{1}{\sqrt{x}}$ . Since  $\frac{2}{x} = o(\frac{1}{\sqrt{x}})$ , then it would be infinitely many natural numbers  $x_0$  such that  $\log g(x_0) > 0$  (Solé and Planat, 2011).

In addition, if  $\log g(x_0) > 0$  for some natural number  $x_0 \geq 5$ , then  $\log g(x_0) = \log g(q_n)$  where  $q_n$  is the greatest prime number such that  $q_n \leq x_0$ . In fact,

$$\prod_{q \leq x_0} \left(1 + \frac{1}{q}\right)^{-1} = \prod_{q \leq q_n} \left(1 + \frac{1}{q}\right)^{-1}$$

and

$$\theta(x_0) = \theta(q_n)$$

according to the definition of the Chebyshev function. ■

We define  $H = \gamma - B$  such that  $B \approx 0.2614972128$  is the Meissel-Mertens constant (Mertens, 1874).

## Theorem 3

We have that (Choie et al., 2007):

$$\sum_{k=1}^{\infty} \left( \log\left(\frac{q_k}{q_k - 1}\right) - \frac{1}{q_k} \right) = \gamma - B = H.$$

## Theorem 4

*It is known that (Edwards, 2001):*

$$\zeta(2) = \prod_{k=1}^{\infty} \frac{q_k^2}{q_k^2 - 1} = \frac{\pi^2}{6}.$$

## Theorem 5

$$\sum_{k=1}^{\infty} \left( \frac{1}{q_k} - \log\left(1 + \frac{1}{q_k}\right) \right) = \log(\zeta(2)) - H.$$

We obtain that

$$\begin{aligned}\log(\zeta(2)) - H &= \log\left(\prod_{k=1}^{\infty} \frac{q_k^2}{q_k^2 - 1}\right) - H \\&= \sum_{k=1}^{\infty} \left( \log\left(\frac{q_k^2}{(q_k^2 - 1)}\right) \right) - H \\&= \sum_{k=1}^{\infty} \left( \log\left(\frac{q_k^2}{(q_k - 1) \times (q_k + 1)}\right) \right) - H \\&= \sum_{k=1}^{\infty} \left( \log\left(\frac{q_k}{q_k - 1}\right) + \log\left(\frac{q_k}{q_k + 1}\right) \right) - H\end{aligned}$$

$$\begin{aligned}
&= \sum_{k=1}^{\infty} \left( \log\left(\frac{q_k}{q_k-1}\right) - \log\left(\frac{q_k+1}{q_k}\right) \right) - H \\
&= \sum_{k=1}^{\infty} \left( \log\left(\frac{q_k}{q_k-1}\right) - \log\left(1 + \frac{1}{q_k}\right) \right) - \sum_{k=1}^{\infty} \left( \log\left(\frac{q_k}{q_k-1}\right) - \frac{1}{q_k} \right) \\
&= \sum_{k=1}^{\infty} \left( \log\left(\frac{q_k}{q_k-1}\right) - \log\left(1 + \frac{1}{q_k}\right) - \log\left(\frac{q_k}{q_k-1}\right) + \frac{1}{q_k} \right) \\
&= \sum_{k=1}^{\infty} \left( \frac{1}{q_k} - \log\left(1 + \frac{1}{q_k}\right) \right)
\end{aligned}$$

and the proof is done. ■

## Theorem 6

*Dedekinds( $q_n$ ) holds if and only if the inequality*

$$\sum_{k=1}^{\infty} \left( \frac{1}{q_k} - (\chi_{\{x: x > q_n\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) > B + \log \log \theta(q_n)$$

*is satisfied for the prime number  $q_n$ , where the set  $S = \{x : x > q_n\}$  contains all the real numbers greater than  $q_n$  and  $\chi_S$  is the characteristic function of the set  $S$  (This is the function defined by  $\chi_S(x) = 1$  when  $x \in S$  and  $\chi_S(x) = 0$  otherwise).*



When  $\text{Dedekinds}(q_n)$  holds, we apply the logarithm to the both sides of the inequality:

$$\log(\zeta(2)) + \sum_{q \leq q_n} \log\left(1 + \frac{1}{q}\right) > \gamma + \log \log \theta(q_n)$$

$$\log(\zeta(2)) - H + \sum_{q \leq q_n} \log\left(1 + \frac{1}{q}\right) > B + \log \log \theta(q_n)$$

$$\sum_{k=1}^{\infty} \left( \frac{1}{q_k} - \log\left(1 + \frac{1}{q_k}\right) \right) + \sum_{q \leq q_n} \log\left(1 + \frac{1}{q}\right) > B + \log \log \theta(q_n).$$

Let's distribute the elements of the inequality to obtain that

$$\sum_{k=1}^{\infty} \left( \frac{1}{q_k} - (\chi_{\{x: x > q_n\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) > B + \log \log \theta(q_n)$$

when  $\text{Dedekinds}(q_n)$  holds. The same happens in the reverse implication. ■

## Theorem 7

*The Riemann Hypothesis is true if the inequality*

$$\theta(q_n)^{1+\frac{1}{q_n}} \geq \theta(q_{n+1})$$

*is satisfied for all sufficiently large prime numbers  $q_n$ .*

The inequality

$$\sum_{k=1}^{\infty} \left( \frac{1}{q_k} - (\chi_{\{x: x > q_n\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) > B + \log \log \theta(q_n)$$

is satisfied when

$$\sum_{k=1}^{\infty} \left( \frac{1}{q_k} - (\chi_{\{x: x \geq q_n\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) > B + \log \log \theta(q_n)$$

is also satisfied, where the set  $S = \{x : x \geq q_n\}$  contains all the real numbers greater than or equal to  $q_n$ .

In the inequality

$$\sum_{k=1}^{\infty} \left( \frac{1}{q_k} - (\chi_{\{x: x \geq q_n\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) > B + \log \log \theta(q_n)$$

only change the values of

$$\log\left(1 + \frac{1}{q_n}\right) + \log \log \theta(q_n)$$

and

$$\log \log \theta(q_{n+1})$$

between the consecutive primes  $q_n$  and  $q_{n+1}$ . It is enough to show that

$$\log\left(1 + \frac{1}{q_n}\right) + \log \log \theta(q_n) \geq \log \log \theta(q_{n+1})$$

for all sufficiently large prime numbers  $q_n$ .

Indeed, the inequality

$$\sum_{k=1}^{\infty} \left( \frac{1}{q_k} - (\chi_{\{x: x \geq q_n\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) > B + \log \log \theta(q_n)$$

is the same as

$$\begin{aligned} & \sum_{k=1}^{\infty} \left( \frac{1}{q_k} - (\chi_{\{x: x \geq q_{n+1}\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) \\ & > B + \log \log \theta(q_{n+1}) + \log\left(1 + \frac{1}{q_n}\right) + \log \log \theta(q_n) - \log \log \theta(q_{n+1}) \end{aligned}$$

where  $q_n$  and  $q_{n+1}$  are consecutive primes.

If the Riemann Hypothesis is false, then

$$\log\left(1 + \frac{1}{q_n}\right) + \log \log \theta(q_n) \geq \log \log \theta(q_{n+1})$$

must be violated for infinitely many  $n$ 's, since  $\text{Dedekinds}(q_{n+1})$  will not hold for infinitely many  $q_{n+1}$ 's. By contraposition, the Riemann Hypothesis should be true when the previous inequality is satisfied for all sufficiently large prime numbers  $q_n$ .

This is

$$\log \left( \left( 1 + \frac{1}{q_n} \right) \times \log \theta(q_n) \right) \geq \log \log \theta(q_{n+1}).$$

That is equivalent to

$$\log \log \theta(q_n)^{1+\frac{1}{q_n}} \geq \log \log \theta(q_{n+1}).$$

To sum up, the Riemann Hypothesis is true when

$$\theta(q_n)^{1+\frac{1}{q_n}} \geq \theta(q_{n+1})$$

is satisfied for all sufficiently large prime numbers  $q_n$ . ■

## Theorem 8

For all  $n \geq 2$  (Ghosh, 2019):

$$\frac{\theta(q_n)}{\log q_{n+1}} \geq n \times \left(1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}\right).$$

## Theorem 9

For  $n \geq 8602$  (Ghosh, 2019):

$$q_n \leq n \times (\log n + \log \log n - 0.9385).$$



## Theorem 10

*The Riemann Hypothesis is true.*

The Riemann Hypothesis is true when

$$\theta(q_n)^{1+\frac{1}{q_n}} \geq \theta(q_{n+1})$$

is satisfied for all sufficiently large prime numbers  $q_n$ . That is the same as

$$\theta(q_n)^{1+\frac{1}{q_n}} \geq \theta(q_n) + \log(q_{n+1})$$

$$\theta(q_n)^{\frac{1}{q_n}} \geq 1 + \frac{\log(q_{n+1})}{\theta(q_n)}$$

after dividing the both sides of the inequality by  $\theta(q_n)$ .

We would only need to prove that

$$1 + \frac{\log \theta(q_n)}{q_n} \geq 1 + \frac{1}{n \times \left(1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}\right)}$$

because of

$$\frac{\theta(q_n)}{\log q_{n+1}} \geq n \times \left(1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}\right).$$

$$\theta(q_n)^{\frac{1}{q_n}} = e^{\frac{\log \theta(q_n)}{q_n}} \geq 1 + \frac{\log \theta(q_n)}{q_n}$$

using the known results. That is equivalent to

$$\left(n \times \left(1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}\right)\right) \times \log \theta(q_n) \geq q_n.$$

Therefore,

$$\left( n \times \left( 1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n} \right) \right) \times \log \theta(q_n) \geq n \times (\log n + \log \log n - 0.9385)$$

which is

$$\left( 1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n} \right) \times \log \theta(q_n) + 0.9385 \geq \log n + \log \log n$$

$$\theta(q_n)^{1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}} \times e^{0.9385} \geq n \times \log n$$

$$e^{0.9385} \geq \frac{\theta(q_n)^{\frac{1}{\log n}} \times n \times \log n}{\theta(q_n)^{1 + \frac{\log \log n}{4 \times \log^2 n}}}$$

under the assumption of the known results. Finally, we conclude this last inequality is satisfied for all sufficiently large prime numbers  $q_n$ . ■

- Aditya Ghosh. An asymptotic formula for the Chebyshev theta function. *Notes on Number Theory and Discrete Mathematics*, 25(4):1–7, 2019. URL <https://nntdm.net/papers/nntdm-25/NTDM-25-4-001-007.pdf>.
- Franz Mertens. Ein Beitrag zur analytischen Zahlentheorie. *J. reine angew. Math.*, 1874 (78):46–62, 1874. URL <https://doi.org/10.1515/crll.1874.78.46>.
- Harold M. Edwards. *Riemann's Zeta Function*. Dover Publications, 2001. URL <https://www.book-info.com/isbn/0-486-41740-9.htm>.

- YoungJu Choie, Nicolas Lichiardopol, Pieter Moree, and Patrick Solé. On Robin's criterion for the Riemann hypothesis. *Journal de Théorie des Nombres de Bordeaux*, 19 (2):357–372, 2007. URL <https://jtnb.centre-mersenne.org/item/10.5802/jtnb.591.pdf>.
- Jean-Louis Nicolas. Petites valeurs de la fonction d'Euler. *Journal of number theory*, 17 (3):375–388, 1983. URL [https://doi.org/10.1016/0022-314X\(83\)90055-0](https://doi.org/10.1016/0022-314X(83)90055-0).
- Patrick Solé and Michel Planat. Extreme values of the Dedekind  $\psi$  function. *Journal of Combinatorics and Number Theory*, 3(1):33–38, 2011. arXiv URL <https://arxiv.org/pdf/1011.1825.pdf>.