

Note on the Riemann Hypothesis

Frank Vega

CopSonic, 1471 Route de Saint-Nauphary 82000 Montauban, France

Abstract

The Riemann Hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$. In 2011, Solé and Planat stated that the Riemann Hypothesis is true if and only if the inequality $\prod_{q \leq q_n} \left(1 + \frac{1}{q}\right) > \frac{e^\gamma}{\zeta(2)} \times \log \theta(q_n)$ is satisfied for all primes $q_n > 3$, where $\theta(x)$ is the Chebyshev function, $\gamma \approx 0.57721$ is the Euler-Mascheroni constant and $\zeta(x)$ is the Riemann zeta function. Using this result, we create a new criterion for the Riemann Hypothesis. We prove the Riemann Hypothesis is true using this new criterion.

Keywords: Riemann Hypothesis, Prime numbers, Chebyshev function, Riemann zeta function
2000 MSC: 11M26, 11A41, 11A25

1. Introduction

The Riemann Hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$. In mathematics, the Chebyshev function $\theta(x)$ is given by

$$\theta(x) = \sum_{p \leq x} \log p$$

with the sum extending over all prime numbers p that are less than or equal to x , where $\log$ is the natural logarithm [1]. We denote the n th prime number as q_n . We know the following property for the Chebyshev function and the n th prime number:

Proposition 1.1. For $n \geq 2$ [2]:

$$\frac{\theta(q_n)}{\log q_{n+1}} \geq n \times \left(1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}\right).$$

Proposition 1.2. For $n \geq 8602$ [3]:

$$q_n \leq n \times (\log n + \log \log n - 0.9385).$$

Email address: vega.frank@gmail.com (Frank Vega)

In mathematics, $\Psi = n \times \prod_{q|n} \left(1 + \frac{1}{q}\right)$ is called the Dedekind Ψ function, where $q | n$ means the prime q divides n . Say $\text{Dedekinds}(q_n)$ holds provided

$$\prod_{q \leq q_n} \left(1 + \frac{1}{q}\right) > \frac{e^\gamma}{\zeta(2)} \times \log \theta(q_n).$$

The constant $\gamma \approx 0.57721$ is the Euler-Mascheroni constant and $\zeta(x)$ is the Riemann zeta function. The importance of this inequality is:

Proposition 1.3. *Dedekinds(q_n) holds for all prime numbers $q_n > 3$ if and only if the Riemann Hypothesis is true [4].*

We define $H = \gamma - B$ such that $B \approx 0.2614972128$ is the Meissel-Mertens constant [5]. We know the following formula:

Proposition 1.4. *We have that [6]:*

$$\sum_{k=1}^{\infty} \left(\log\left(\frac{q_k}{q_k - 1}\right) - \frac{1}{q_k} \right) = \gamma - B = H.$$

In addition, we know this value of the Riemann zeta function:

Proposition 1.5. *It is known that [7]:*

$$\zeta(2) = \prod_{k=1}^{\infty} \frac{q_k^2}{q_k^2 - 1} = \frac{\pi^2}{6}.$$

Putting all together yields a proof for the Riemann Hypothesis using the Chebyshev function.

2. What if the Riemann Hypothesis were false?

Theorem 2.1. *If the Riemann Hypothesis is false, then there are infinitely many prime numbers q_n for which $\text{Dedekinds}(q_n)$ does not hold.*

Proof. The Riemann Hypothesis is false, if there exists some natural number $x_0 \geq 5$ such that $g(x_0) > 1$ or equivalent $\log g(x_0) > 0$ [4]:

$$g(x) = \frac{e^\gamma}{\zeta(2)} \times \log \theta(x) \times \prod_{q \leq x} \left(1 + \frac{1}{q}\right)^{-1}.$$

We know the bound [4]:

$$\log g(x) \geq \log f(x) - \frac{2}{x}$$

where f is introduced in the Nicolas paper [1]:

$$f(x) = e^\gamma \times \log \theta(x) \times \prod_{q \leq x} \left(1 - \frac{1}{q}\right).$$

When the Riemann Hypothesis is false, then there exists a real number $b < \frac{1}{2}$ for which there are infinitely many natural numbers x such that $\log f(x) = \Omega_+(x^{-b})$ [1]. According to the Hardy and Littlewood definition, this would mean that

$$\exists k > 0, \forall y_0 \in \mathbb{N}, \exists y \in \mathbb{N} > y_0: \log f(y) \geq k \times y^{-b}.$$

That inequality is equivalent to $\log f(y) \geq (k \times y^{-b} \times \sqrt{y}) \times \frac{1}{\sqrt{y}}$, but we note that

$$\lim_{y \rightarrow \infty} (k \times y^{-b} \times \sqrt{y}) = \infty$$

for every possible positive value of k when $b < \frac{1}{2}$. In this way, this implies that

$$\forall y_0 \in \mathbb{N}, \exists y \in \mathbb{N} > y_0: \log f(y) \geq \frac{1}{\sqrt{y}}.$$

Hence, if the Riemann Hypothesis is false, then there are infinitely many natural numbers x such that $\log f(x) \geq \frac{1}{\sqrt{x}}$. Since $\frac{2}{x} = o(\frac{1}{\sqrt{x}})$, then it would be infinitely many natural numbers x_0 such that $\log g(x_0) > 0$ [4]. In addition, if $\log g(x_0) > 0$ for some natural number $x_0 \geq 5$, then $\log g(x_0) = \log g(q_n)$ where q_n is the greatest prime number such that $q_n \leq x_0$. Actually,

$$\prod_{q \leq x_0} \left(1 + \frac{1}{q}\right)^{-1} = \prod_{q \leq q_n} \left(1 + \frac{1}{q}\right)^{-1}$$

and

$$\theta(x_0) = \theta(q_n)$$

according to the definition of the Chebyshev function. □

3. A Key Theorem

Theorem 3.1.

$$\sum_{k=1}^{\infty} \left(\frac{1}{q_k} - \log\left(1 + \frac{1}{q_k}\right) \right) = \log(\zeta(2)) - H.$$

Proof. We obtain that

$$\begin{aligned} \log(\zeta(2)) - H &= \log\left(\prod_{k=1}^{\infty} \frac{q_k^2}{q_k^2 - 1}\right) - H \\ &= \sum_{k=1}^{\infty} \left(\log\left(\frac{q_k^2}{q_k^2 - 1}\right) \right) - H \\ &= \sum_{k=1}^{\infty} \left(\log\left(\frac{q_k^2}{(q_k - 1) \times (q_k + 1)}\right) \right) - H \\ &= \sum_{k=1}^{\infty} \left(\log\left(\frac{q_k}{q_k - 1}\right) + \log\left(\frac{q_k}{q_k + 1}\right) \right) - H \end{aligned}$$

where

$$\begin{aligned}
&= \sum_{k=1}^{\infty} \left(\log\left(\frac{q_k}{q_k-1}\right) - \log\left(\frac{q_k+1}{q_k}\right) \right) - H \\
&= \sum_{k=1}^{\infty} \left(\log\left(\frac{q_k}{q_k-1}\right) - \log\left(1 + \frac{1}{q_k}\right) \right) - \sum_{k=1}^{\infty} \left(\log\left(\frac{q_k}{q_k-1}\right) - \frac{1}{q_k} \right) \\
&= \sum_{k=1}^{\infty} \left(\log\left(\frac{q_k}{q_k-1}\right) - \log\left(1 + \frac{1}{q_k}\right) - \log\left(\frac{q_k}{q_k-1}\right) + \frac{1}{q_k} \right) \\
&= \sum_{k=1}^{\infty} \left(\frac{1}{q_k} - \log\left(1 + \frac{1}{q_k}\right) \right)
\end{aligned}$$

and the proof is done. $\square$

4. A New Criterion

Theorem 4.1. *Dedekinds(q_n) holds if and only if the inequality*

$$\sum_{k=1}^{\infty} \left(\frac{1}{q_k} - (\chi_{\{x: x > q_n\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) > B + \log \log \theta(q_n)$$

is satisfied for the prime number q_n , where the set $S = \{x : x > q_n\}$ contains all the real numbers greater than q_n and χ_S is the characteristic function of the set S (This is the function defined by $\chi_S(x) = 1$ when $x \in S$ and $\chi_S(x) = 0$ otherwise).

Proof. When Dedekinds(q_n) holds, we apply the logarithm to the both sides of the inequality:

$$\begin{aligned}
&\log(\zeta(2)) + \sum_{q \leq q_n} \log\left(1 + \frac{1}{q}\right) > \gamma + \log \log \theta(q_n) \\
&\log(\zeta(2)) - H + \sum_{q \leq q_n} \log\left(1 + \frac{1}{q}\right) > B + \log \log \theta(q_n) \\
&\sum_{k=1}^{\infty} \left(\frac{1}{q_k} - \log\left(1 + \frac{1}{q_k}\right) \right) + \sum_{q \leq q_n} \log\left(1 + \frac{1}{q}\right) > B + \log \log \theta(q_n).
\end{aligned}$$

Let's distribute the elements of the inequality to obtain that

$$\sum_{k=1}^{\infty} \left(\frac{1}{q_k} - (\chi_{\{x: x > q_n\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) > B + \log \log \theta(q_n)$$

when Dedekinds(q_n) holds. The same happens in the reverse implication. $\square$

5. The Main Insight

Theorem 5.1. *The Riemann Hypothesis is true if the inequality*

$$\theta(q_n)^{1+\frac{1}{q_n}} \geq \theta(q_{n+1})$$

is satisfied for all sufficiently large prime numbers q_n .

Proof. The inequality

$$\sum_{k=1}^{\infty} \left(\frac{1}{q_k} - (\chi_{\{x: x > q_n\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) > B + \log \log \theta(q_n)$$

is satisfied when

$$\sum_{k=1}^{\infty} \left(\frac{1}{q_k} - (\chi_{\{x: x \geq q_n\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) > B + \log \log \theta(q_n)$$

is also satisfied, where the set $S = \{x : x \geq q_n\}$ contains all the real numbers greater than or equal to q_n . In the inequality

$$\sum_{k=1}^{\infty} \left(\frac{1}{q_k} - (\chi_{\{x: x \geq q_n\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) > B + \log \log \theta(q_n)$$

only change the values of

$$\log\left(1 + \frac{1}{q_n}\right) + \log \log \theta(q_n)$$

and

$$\log \log \theta(q_{n+1})$$

between the consecutive primes q_n and q_{n+1} . It is enough to show that

$$\log\left(1 + \frac{1}{q_n}\right) + \log \log \theta(q_n) \geq \log \log \theta(q_{n+1})$$

for all sufficiently large prime numbers q_n . Indeed, the inequality

$$\sum_{k=1}^{\infty} \left(\frac{1}{q_k} - (\chi_{\{x: x \geq q_n\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) > B + \log \log \theta(q_n)$$

is the same as

$$\begin{aligned} & \sum_{k=1}^{\infty} \left(\frac{1}{q_k} - (\chi_{\{x: x \geq q_{n+1}\}}(q_k)) \times \log\left(1 + \frac{1}{q_k}\right) \right) \\ & > B + \log \log \theta(q_{n+1}) + \log\left(1 + \frac{1}{q_n}\right) + \log \log \theta(q_n) - \log \log \theta(q_{n+1}) \end{aligned}$$

where q_n and q_{n+1} are consecutive primes. If the Riemann Hypothesis is false, then

$$\log\left(1 + \frac{1}{q_n}\right) + \log \log \theta(q_n) \geq \log \log \theta(q_{n+1})$$

must be violated for infinitely many n 's, since $\text{Dedekinds}(q_{n+1})$ will not hold for infinitely many q_{n+1} 's. By contraposition, the Riemann Hypothesis should be true when the previous inequality is satisfied for all sufficiently large prime numbers q_n . This is

$$\log \left(\left(1 + \frac{1}{q_n} \right) \times \log \theta(q_n) \right) \geq \log \log \theta(q_{n+1}).$$

That is equivalent to

$$\log \log \theta(q_n)^{1+\frac{1}{q_n}} \geq \log \log \theta(q_{n+1}).$$

To sum up, the Riemann Hypothesis is true when

$$\theta(q_n)^{1+\frac{1}{q_n}} \geq \theta(q_{n+1})$$

is satisfied for all sufficiently large prime numbers q_n . □

6. The Main Theorem

Theorem 6.1. *The Riemann Hypothesis is true.*

Proof. The Riemann Hypothesis is true when

$$\theta(q_n)^{1+\frac{1}{q_n}} \geq \theta(q_{n+1})$$

is satisfied for all sufficiently large prime numbers q_n . That is the same as

$$\theta(q_n)^{1+\frac{1}{q_n}} \geq \theta(q_n) + \log(q_{n+1})$$

$$\theta(q_n)^{\frac{1}{q_n}} \geq 1 + \frac{\log(q_{n+1})}{\theta(q_n)}$$

after dividing the both sides of the inequality by $\theta(q_n)$. We would only need to prove that

$$1 + \frac{\log \theta(q_n)}{q_n} \geq 1 + \frac{1}{n \times \left(1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n} \right)}$$

because of

$$\frac{\theta(q_n)}{\log q_{n+1}} \geq n \times \left(1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n} \right)$$

$$\theta(q_n)^{\frac{1}{q_n}} = e^{\frac{\log \theta(q_n)}{q_n}} \geq 1 + \frac{\log \theta(q_n)}{q_n}.$$

That is equivalent to

$$\left(n \times \left(1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n} \right) \right) \times \log \theta(q_n) \geq q_n.$$

Therefore,

$$\left(n \times \left(1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n} \right) \right) \times \log \theta(q_n) \geq n \times (\log n + \log \log n - 0.9385)$$

which is

$$\left(1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}\right) \times \log \theta(q_n) + 0.9385 \geq \log n + \log \log n$$

$$\theta(q_n)^{1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}} \times e^{0.9385} \geq n \times \log n$$

$$e^{0.9385} \geq \frac{n \times \log n}{\theta(q_n)^{1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}}}.$$

However, we know that

$$\overline{\lim}_{n \rightarrow \infty} \frac{n \times \log n}{\theta(q_n)^{1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}}} = \lim_{n \rightarrow \infty} \frac{n \times \log n}{\theta(q_n)^{1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}}} = 1$$

since

$$\lim_{n \rightarrow \infty} \left(1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}\right) = 1$$

$$\theta(q_n) \sim q_n, \quad (n \rightarrow \infty)$$

$$q_n \sim n \times \log n, \quad (n \rightarrow \infty).$$

For any positive real number ε , there exists a natural number m such that

$$\frac{n \times \log n}{\theta(q_n)^{1 - \frac{1}{\log n} + \frac{\log \log n}{4 \times \log^2 n}}} < 1 + \varepsilon$$

for all $n > m$, because of the definition of limit superior. Moreover, we can see that $e^{0.9385} > 2.5561$. Consequently, it is enough to take any positive real number $\varepsilon \leq 1.5561$. By the definition of the limit superior yields the proof of the Riemann Hypothesis. $\square$

Acknowledgments

The author would like to thank Patrick Solé for helpful comments and his mother, maternal brother and his friend Sonia for their support.

References

- [1] J.-L. Nicolas, Petites valeurs de la fonction d'Euler, *Journal of number theory* 17 (3) (1983) 375–388. doi:10.1016/0022-314X(83)90055-0.
- [2] A. Ghosh, An asymptotic formula for the Chebyshev theta function, *Notes on Number Theory and Discrete Mathematics* 25 (4) (2019) 1–7. doi:10.7546/nntdm.2019.25.4.1-7.
- [3] J.-P. Massias, G. Robin, Bornes effectives pour certaines fonctions concernant les nombres premiers, *Journal de Théorie des Nombres de Bordeaux* 8 (1) (1996) 215–242.
- [4] P. Solé, M. Planat, Extreme values of the Dedekind ψ function, *Journal of Combinatorics and Number Theory* 3 (1) (2011) 33–38.
- [5] F. Mertens, Ein Beitrag zur analytischen Zahlentheorie., *J. reine angew. Math.* 1874 (78) (1874) 46–62. doi:10.1515/crll.1874.78.46.
- [6] Y. Choie, N. Lichiardopol, P. Moree, P. Solé, On Robin's criterion for the Riemann hypothesis, *Journal de Théorie des Nombres de Bordeaux* 19 (2) (2007) 357–372. doi:10.5802/jtnb.591.
- [7] H. M. Edwards, *Riemann's Zeta Function*, Dover Publications, 2001.