

# Novel Geometric Series for Application of Computational Science

Chinnaraji Annamalai

School of Management, Indian Institute of Technology, Kharagpur, India

Email: [anna@iitkgp.ac.in](mailto:anna@iitkgp.ac.in)

<https://orcid.org/0000-0002-0992-2584>

**Abstract:** Computational science is a rapidly growing multi-and inter-disciplinary area where science, engineering, information technology, management and its collaboration use advance computing capabilities to understand and solve the most complex real life problems. In this article, a new geometric series is constituted for application of computational science and engineering.

**MSC Classification codes:** 40A05 (65B10)

**Keywords:** computation, theorem, geometric series

## 1. Introduction

Geometric series played a vital role in differential and integral calculus at the earlier stage of development and still continues as an important part of the study in science, engineering, management and its applications. In this article, a new geometric series [1-6] is constructed for application [7] of computational science and engineering.

## 2. Novel Geometric Series

**Theorem :**  $\sum_{j=k}^n x^{pj} = \frac{x^{p(n+1)} - x^{pk}}{x^p - 1}$ , where p is any number in the systems of numbers.

*Proof.* Let us prove this theorem by the equality given below:

$$y^{n+1} = y^{n+1} \Rightarrow y^{n+1} = (y-1)y^n + y^n \Rightarrow y^{n+1} = (y-1)y^n + (y-1)y^{n-1} + y^{n-1}.$$

If we continue this expansion again and again, we can obtain the following series.

$$y^{n+1} = (y-1)y^n + (y-1)y^{n-1} + (y-1)y^{n-2} + (y-1)y^{n-3} + \dots + (y-1)y^k + y^k.$$

$$\text{i. e., } y^{n+1} - y^k = (y-1)(y^n + y^{n-1} + y^{n-2} + y^{n-3} + \dots + y^k) \quad (1)$$

$$\text{From the expansion (1), we obtain the series } \sum_{j=k}^n y^j = \frac{y^{n+1} - y^k}{y-1} \quad (2)$$

Now, let us prove the theorem shown below:

$$1 + x^p + x^{2p} + x^{3p} + \dots + x^{kp} = \frac{x^{p(n+1)} - x^{pk}}{x^p - 1}.$$

Let y be  $x^p$  and substitute it in the series (2).

$$\text{Then, } \sum_{j=k}^n (x^p)^j = \frac{(x^p)^{n+1} - (x^p)^k}{x^p - 1} \quad (3)$$

From the Equation (3), we conclude that

$$\sum_{j=k}^n x^{pj} = \frac{x^{p(n+1)} - x^{pk}}{x^p - 1}, \text{ where } p \text{ is any number in the defined sets of numbers.}$$

Hence, theorem is proved.

Note that the novel geometric series is: 
$$\sum_{j=0}^n x^{pj} = \frac{x^{p(n+1)} - 1}{x^p - 1}.$$

For example,

$$\begin{aligned} & 1 + x^{0.25} + x^{0.5} + x^{0.75} + x^{1.0} + x^{1.25} + x^{1.5}. \\ &= 1 + x^{1(0.25)} + x^{2(0.25)} + x^{3(0.25)} + x^{4(0.25)} + x^{5(0.25)} + x^{6(0.25)} = \frac{x^{0.25(6+1)} - 1}{x^{0.25} - 1}. \\ &\therefore 1 + x^{0.25} + x^{0.5} + x^{0.75} + x^{1.0} + x^{1.25} + x^{1.5} = \frac{x^{1.75} - 1}{x^{0.25} - 1}. \end{aligned}$$

$$\text{Also, } x^{0.5} + x^{0.75} + x^{1.0} + x^{1.25} + x^{1.5} = \frac{x^{1.75} - x^{0.5}}{x^{0.25} - 1}.$$

**Corollary 1:** 
$$\sum_{j=k}^n (ax)^{pj} = \frac{(ax)^{p(n+1)} - (ax)^{pk}}{(ax)^p - 1}.$$

**Corollary 2:** 
$$\sum_{j=0}^n (ax)^{pj} = \frac{(ax)^{p(n+1)} - 1}{(ax)^p - 1}.$$

**Corollary 3:** 
$$\sum_{j=-k}^n (ax)^{pj} = \frac{(ax)^{p(n+1)} - (ax)^{-pk}}{(ax)^p - 1}.$$

### 3. Conclusion

Nowadays, the growing complexity of mathematical and computational modelling demands the simplicity of mathematical and combinatorial equations for solving today's scientific problems and challenges. This article provides a novel geometric series and its theorem. This idea can enable the scientific researchers for further involvement in research and development.

### References

- [1] Annamalai, C. (2018) Annamalai's Computing Model for Algorithmic Geometric Series and Its Mathematical Structures. *Journal of Mathematics and Computer Science*, 3(1),1-6 <https://doi.org/10.11648/j.mcs.20180301.11>.
- [2] Annamalai, C. (2018) Algorithmic Computation of Annamalai's Geometric Series and Summability. *Journal of Mathematics and Computer Science*, 3(5),100-101. <https://doi.org/10.11648/j.mcs.20180305.11>.
- [3] Annamalai, C. (2017) Analysis and Modelling of Annamalai Computing Geometric Series and Summability. *Mathematical Journal of Interdisciplinary Sciences*, 6(1), 11-15. <https://doi.org/10.15415/mjis.2017.61002>.

- [4] Annamalai, C. (2018) Novel Computation of Algorithmic Geometric Series and Summability. *Journal of Algorithms and Computation*, 50(1), 151-153.  
<https://www.doi.org/10.22059/JAC.2018.68866>.
- [5] Annamalai, C. (2017) Computational modelling for the formation of geometric series using Annamalai computing method. *Jñānābha*, 47(2), 327-330.  
<https://zbmath.org/?q=an%3A1391.65005>.
- [6] Annamalai, C. (2018) Computing for Development of A New Summability on Multiple Geometric Series. *International Journal of Mathematics, Game Theory and Algebra*, 27(4), 511-513.
- [7] Annamalai C (2010) “Application of Exponential Decay and Geometric Series in Effective Medicine”, *Advances in Bioscience and Biotechnology*, Vol. 1(1), pp 51-54.  
<https://doi.org/10.4236/abb.2010.11008>.