







## Article

# Psychometric Properties of the Connor-Davidson Resilience Scale (CD-RISC) in Spanish Adolescents

Francisco Javier López-Fernández<sup>1</sup> , Paula Morales-Hidalgo<sup>2,3</sup> , Josefa Canals<sup>2</sup> , Juan Carlos Marzo<sup>4</sup> ,  
Luis Joaquín García-López<sup>5</sup>  and José Antonio Piqueras<sup>4</sup> 

<sup>1</sup>Universitat Jaume I (Spain); <sup>2</sup>Universitat Rovira i Virgili (Spain); <sup>3</sup>Universitat Oberta de Catalunya (Spain); <sup>4</sup>Universidad Miguel Hernández de Elche (Spain); <sup>5</sup>Universidad de Jaén (Spain)

## Abstract

Few studies have examined the psychometric properties of the Connor-Davidson Resilience Scale (CD-RISC) in a large adolescent community sample, finding a significant disparity. This study explores the psychometric properties of the CD-RISC among Spanish adolescents by means of exploratory factor analysis (EFA), Rasch analysis, and measurement invariance (MI) across sex, as well as internal consistency and criterion validity. The sample was comprised of 463 adolescents (231 girls), aged 12 to 18 years, who completed the CD-RISC and other measures on emotional status and quality of life. The EFA suggested that the CD-RISC structure presented a unidimensional model. Consequently, shorter unidimensional CD-RISC models observed in the literature were explored. Thus, the Campbell-Sills and Stein CD-RISC-10 showed the soundest psychometric properties, providing an adequate item fit and supporting MI and non-differential item functioning across sex. Item difficulty levels were biased toward low levels of resilience. Some items showed malfunctioning in lower response categories. With regard to reliability, categorical omega was .82. Strong associations with health-related quality of life, major depressive disorder symptoms, and emotional symptoms were observed. A weak association was found between resilience and the male sex. Campbell-Sills and Stein's CD-RISC-10 model emerges as the best to assess resilience among Spanish adolescents, as already reported in adults. Thus, independently of the developmental stage, the core of resilience may reside in the aspects of hardiness and persistence.

**Keywords:** adolescents; CD-RISC; factor analysis; measurement invariance; Rasch analysis

(Received: 21 February 2023; revised: 22 December 2023; accepted: 09 January 2024)

The construct of resilience, which has received increasing attention in the last decades, is conceptualized as personal characteristics that promote positive adaptation in the face of adversity (Fletcher & Sarkar, 2013). Resilient individuals actively cope with stress and trauma (Li & Nishikawa, 2012), through adaptive psychological and physiological responses (Feder et al., 2009). A vast number of studies have demonstrated that resilience emerges as a key protective factor against negative indicators of mental health, including depression, anxiety, and negative affect (see Hu et al., 2015, for a meta-analysis). It has also been related to positive mental health outcomes, such as life satisfaction and positive affect, with moderate to large effect sizes (Hu et al., 2015). Regarding intervention, significant improvements have been observed in resilience after applying a combination of cognitive behavior and pharmacological treatments, whereas post-traumatic stress symptoms are reduced (Connor & Davidson, 2003; Davidson et al., 2005). In this line, resilience interventions improve the levels of resilience and reduce

depressive and stress symptoms (see Ang et al., 2022, for a meta-analysis).

The Connor-Davidson Resilience Scale (CD-RISC; Connor & Davidson, 2003) is one of the most employed instruments to measure resilience and has shown good psychometric properties (Windle et al., 2011). Based on exploratory factor analysis (EFA), the 25 items of the original scale converge into 5 factors: Personal competence, high standards, and tenacity; trust in one's instincts, tolerance of negative affect, and strengthening effects of stress; positive acceptance of change and secure relationships; control; and spiritual influences (Connor & Davidson, 2003). However, the original factor structure usually fails to be replicated across independent samples of adolescents and adults from the same or different cultural settings, including Spanish adults (García-León et al., 2019; Pulido-Martos et al., 2020). In response to these structural problems, Campbell-Sills and Stein developed a brief 10-item version of the CD-RISC, which assesses a general factor of resilience focused on hardiness and persistence (Campbell-Sills & Stein, 2007). The CD-RISC-10 structure has been more successfully supported across research, including Spanish adults (Notario-Pacheco et al., 2011; Pulido-Martos et al., 2020) and recently adolescent populations (Chen et al., 2022). In this sense, just a few studies have examined the factor structure of the original 25-item CD-RISC in large samples of adolescents across cultural settings, whose results show a significant disparity in the structure. Even two studies carried out among Chinese adolescents who

**Corresponding author:** Correspondence concerning this article should be addressed to José Antonio Piqueras. Universidad Miguel Hernández de Elche. Área de Personalidad, Evaluación y Tratamiento Psicológico. Departamento de Psicología de la Salud. Avda. de la Universidad, s/n, Edificio Altamira. 03202 Elche (Spain). E-mail: [jpiqueras@umh.es](mailto:jpiqueras@umh.es) Phone: +34-966658343.

**Cite this article:** López-Fernández FJ, Morales-Hidalgo P, Canals J, Marzo JC, García-López LJ and Piqueras JA (2024). Psychometric Properties of the Connor-Davidson Resilience Scale (CD-RISC) in Spanish Adolescents. *The Spanish Journal of Psychology* 27, e3, 1–10. <https://doi.org/10.1017/SJP.2024.3>

suffered the 2008 Sichuan earthquake obtained different factor structures in the scale. Whereas one study, through an EFA, showed a 2-factor model consisting of rational thinking and self-awareness dimensions (Fu *et al.*, 2014), the other work replicated the original 5-factor model through confirmatory factor analysis, CFA, (Yu *et al.*, 2011). Moreover, the original factor model was not replicated in a sample of 701 South African adolescents, and an EFA suggested a possible 3-factor structure (Jorgensen & Seedat, 2008). Finally, a single-factor model of the original 25-item CD-RISC was found in a sample of 988 Colombian adolescents, and a 10-item CD-RISC version was obtained, presenting a good model fit (Guarnizo Guzmán *et al.*, 2019) but retaining different items from the Campbell-Sills and Stein CD-RISC-10 (Campbell-Sills & Stein, 2007).

Most of the psychometric studies on CD-RISC do not explore sex differences. Yet, sex has emerged as a key factor moderating the relationship between resilience and mental health (Hu *et al.*, 2015). Two meta-analyses exploring sex-based differences in resilience found mixed results: Whereas one displayed equal resilience levels between women and men (Ari & Çarkit, 2020), the other reported higher levels of resilience in favor of men (Yılmaz Koğar & Gök, 2021). In this sense, when considered a unidimensional CD-RISC factor structure, partial invariance across sex was reported over the lifespan, from young to older adults, but Items 2, 3, and 9 had to be removed because of their low loadings (Liu *et al.*, 2015). Additionally, 10-item CD-RISC versions have supported measurement invariance (MI) based on sex at metric and scalar levels in adults (Gonzalez *et al.*, 2016; Pulido-Martos *et al.*, 2020) and adolescents (Chen *et al.*, 2022; Guarnizo Guzmán *et al.*, 2019). Due to the few CD-RISC studies on sex MI, especially in adolescents, we must ensure that the resilience construct assessed by CD-RISC is psychometrically equivalent for male and female adolescents (Putnick & Bornstein, 2016) or make the necessary adjustments if this is not the case.

Rasch analyses, which provide information about item fit, have been conducted on CD-RISC validation studies among adults exclusively. When the fit of the original CD-RISC-25 has been explored, versions of 21 and 22 items with acceptable fit have also been obtained (Arias González *et al.*, 2015; Papini *et al.*, 2021). Regarding Campbell-Sills and Stein's CD-RISC-10, one study found that all the items confirmed the unifactorial structure of resilience (Pulido-Martos *et al.*, 2020), whereas others reported the misfit of some items and proposed abridged versions of 7 and 8 items (Ehrich *et al.*, 2017; Heritage *et al.*, 2021). The test for differential item functioning (DIF) according to sex suggested non-substantial differences in item responses between males and females (Arias González *et al.*, 2015; Ehrich *et al.*, 2017; Heritage *et al.*, 2021; Papini *et al.*, 2021; Pulido-Martos *et al.*, 2020), just a study found a differential item functioning in Item 2 "Close and secure relationships", more difficult for men (Arias González *et al.*, 2015). Furthermore, the exploration of item difficulty is essential to check if items effectively discriminate between individuals with different levels of resilience. In this manner, the item difficulty levels of CD-RISC seem to be biased toward low levels of resilience (Arias González *et al.*, 2015; Heritage *et al.*, 2021; Papini *et al.*, 2021). As for item response categories, some studies conducted on the original 25-item CD-RISC (Arias González *et al.*, 2015; Papini *et al.*, 2021) and the 10-item Campbell-Sills and Stein CD-RISC (Heritage *et al.*, 2021; Wadimba *et al.*, 2022) observed non-redundant response categories, with thresholds ascending ordered.

To our knowledge, CD-RISC psychometric properties (i.e., factor and Rasch analyses, and MI across sex) are still unexplored among Spanish adolescents. Validating a sound measure such as CD-RISC at these ages is important to assess resilience rigorously, which, therefore, may constitute a potential treatment outcome measure (Brownlee *et al.*, 2013). Consequently, the current study aimed to examine the psychometric properties of a CD-RISC version for Spanish adolescents.

Due to its inconsistent factor structure across research, the CD-RISC-25 was submitted to an EFA instead of a CFA (Fabrigar *et al.*, 1999). Once the factor structure is checked, we will conduct Rasch analyses, and test MI across sex. The CD-RISC scores will be associated with an extensive nomological network of psychopathological problems and health-related quality of life. It is hypothesized that the resulting CD-RISC model will exhibit positive correlations with positive indicators of mental health and negative ones with negative outcomes (Hu *et al.*, 2015). Specifically, we expect to find moderate to large associations with emotional/depression symptoms and anxiety problems (García-León *et al.*, 2019; Kuiper *et al.*, 2019; Wadimba *et al.*, 2022; Wollny & Jacobs, 2021; Yu *et al.*, 2011), as well as with health-related quality of life (Wollny & Jacobs, 2021). Finally, internal consistency for the final CD-RISC model will be properly estimated.

## Method

### Participants

Participants were recruited from different high schools located in urban areas of Alicante and Tarragona (eastern Spain). The sample consisted of 463 adolescents (49.9% females), ages 12 to 18 years ( $M = 14.81$ ,  $SD = 1.48$ ). Adolescents under 16 years of age, together with their parents/legal guardians, gave written informed consent following the Declaration of Helsinki, whereas adolescents above the age of 16 gave the informed content exclusively.

### Instruments

The *Connor-Davison Resilience Scale* (CD-RISC; Connor & Davidson, 2003). The 25-item scale assesses resilience on a 5-point Likert scale ranging from 0 (*not true at all*) to 4 (*true all the time*). The participants answered each item considering the last month. The Spanish version was employed, provided under official approval by the authors of the original version (Bobes *et al.*, 2001).

The *KIDSCREEN-10* (Ravens-Sieberer *et al.*, 2010). Participants rated the 10-item questionnaire on 5-point Likert scale. The items explore the adolescent's physical activity and fitness, depressive moods, leisure time, relationships with parents and peers, and perception of school performance and cognitive capacity. Higher scores indicate greater well-being and health-related quality of life. Categorical omega was .87.

The *Strength and Difficulties Questionnaire* (SDQ; Goodman, 2001); Spanish self-reported version (Gómez-Beneyto *et al.*, 2013). The 25-item scale assesses 5 factors: Emotional symptoms, conduct problems, hyperactivity, peer problems, and prosocial behaviors in children and adolescents on a 3-point Likert scale (0 = *not true*, 1 = *somewhat true*, and 2 = *certainly true*). In the current study, categorical omega of the factors was the following: .76 for emotional symptoms, .61 for conduct problems, .69 for hyperactivity, .62 for peer problems, and .66 for prosocial behaviors.

The *Revised Child Anxiety and Depression Scale–30 (RCADS–30)*; Sandín et al., 2010); a reduced Spanish version of the original scale (Chorpita et al., 2000). It consists of 5-item subscales corresponding to 6 factors assessing anxiety and depression symptoms: Separation anxiety disorder (SAD), social phobia (SP), generalized anxiety disorder (GAD), panic disorder (PD), obsessive-compulsive disorder (OCD), and major depressive disorder (MDD). Items are rated on a 4-point Likert scale ranging from 0 (*never*) to 3 (*always*). Categorical omegas were: .67 for SAD, .85 for SP, .82 for GAD, .82 for PD, .77 for OCD, and .82 for MDD.

### Procedure

The study was conducted within the framework of the following projects: PSI2017–88280–R, RTI2018–097124–B–100, and PID2019–111138RB–I00/AEI/10.13039/501100011033, funded respectively by the Ministerio de Economía, Industria y Competitividad, the Ministerio de Ciencia, Innovación y Universidades, and the Agencia Estatal de Investigación of Spain.

This research was approved by the university's Ethical Committees and authorized by the Valencian and Catalan regional education authorities.

### Data Analyses

Mplus, version 8 (Muthén & Muthén, 2017), was employed to perform EFAs on the 25 items of the original CD-RISC scale and the subsequent CD-RISC models. Following the recommendations for EFAs suitable for our data (Fabrigar et al., 1999; Goretzko et al., 2021; Izquierdo et al., 2014), we employed the weighted least squares means and variance adjusted (WLSMV) estimator, using polychoric correlations, and oblimin rotation. WLSMV is particularly employed to handle ordinal data which is non-normally distributed. The number of factors was determined by optimal implementation of parallel analysis (Timmerman & Lorenzo-Seva, 2011) with polychoric correlations, as suggested, more appropriate for the skewness of ordinal data than Pearson correlations (Garrido et al., 2013), using the software FACTOR (Lorenzo-Seva & Ferrando, 2006). Furthermore, multiple-group CFAs in Mplus were conducted to assess invariance of the CD-RISC model extracted across sex. Hence, a series of increasingly stringent multiple-group models was conducted (configural, ensuring that the construct has the same pattern of free and fixed loading; metric, testing for the equivalence of the item loadings on the factor; and scalar, examining that mean differences in the latent construct represent all the mean differences in the shared variance of the items; Putnick & Bornstein, 2016).

Once the CD-RISC unidimensionality was confirmed, tau-equivalence of items (i.e., equal discrimination parameters) was estimated (Viladrich et al., 2017) in Mplus<sup>1</sup>. Accordingly, Rasch analyses were conducted using the Jamovi (2.3) software (The Jamovi project, 2022), through the module snowIRT: Item Response Theory (Seol, 2022). The Polytomous Rasch Model was used to examine the quality of the individual items in the CD-RISC factor model obtained. We employed in fit and outfit statistics, which indicate the information-weighted mean square residuals between observed and expected responses (Linacre, 2002).

A Wright map (person-item map), which shows each participant's resilience level and item difficulties ordered from the highest to the lowest levels, was depicted. The goodness of item response categories, based on how much more likely an individual is to choose a higher response category compared to the previous one, given their level of proficiency; was estimated through delta-tau parameterization of the partial credit model (Masters & Wright, 1997). Delta parameters represent the differences between adjacent threshold values or category boundaries of an item, whereas tau parameters represent the threshold values themselves. The threshold values indicate the points along the latent trait continuum where the likelihood of choosing a higher response category increases. In this sense, response option characteristic curves of items were also examined, exhibiting response category probabilities for the five response category options in the scale. In addition, DIF for ordinal data analysis (Hladká & Martinková, 2020) was performed to test MI across sex at item level, that is, whether the items of the CD-RISC model functioned differently for males and females (Myers et al., 2006).

Spearman correlation coefficients between the CD-RISC final model scores and the rest of the target variables were performed with SPSS, version 28. Categorical omega (Flora, 2020) was calculated through R, version 3.5.0. We used this reliability index, appropriated for categorical data, for the rest of the study scales solely since ordinal alpha requires the assumption of tau-equivalence of items (Flora, 2020; Viladrich et al., 2017). The online Supplementary material Table S1 displays the descriptive statistics of the study variables for the total sample and divided by sex and levels of resilience.

Goodness-of-fit for the factor analyses was assessed with various indices (West et al., 2012). The chi-square ( $\chi^2$ ) was estimated; a nonsignificant chi-square implies a well-fitting model. As this test is highly sensitive to large sample sizes, other fit indices were also considered. Values of .95 or above for the comparative fit index (CFI) and the Tucker-Lewis index (TLI), and of .06 or below for the root mean square error of approximation (RMSEA) suggest a good fit (Hu & Bentler, 1999). For the RMSEA 90% CI, values below .05 for the lower boundary and below .08 for the upper boundary are considered acceptable fit (MacCallum et al., 1996). Sex MI adequacy was tested through multiple-group CFAs. The Satorra-Bentler scaled chi-square test,  $\chi^2$ S-B, was estimated to assess the change in model fit tests (Satorra, 2000). However, due to the test's sensitivity to sample size, other indices were considered to examine the invariance between more and less constrained models. Therefore, the following criteria were used to consider a model to be invariant:  $\Delta CFI \leq 0.01$  (Cheung & Rensvold, 2002) and  $\Delta RMSEA \leq 0.015$  (Chen, 2007).

For the Polytomous Rasch Model, infit and outfit values in the range of 0.60–1.40 suggest a good fit (Linacre, 1999), whereas a value below the range indicates an item is redundant, and a value above the range expresses an item being out-of-concept (Khan et al., 2013). As for sex DIF, items are considered to present a different performance between groups if they show statistical significance,  $p < .05$  (Dorans et al., 1992).

### Results

The EFA and the parallel analysis were performed on the CD-RISC–25 and suggested that the CD-RISC items measured a unidimensional construct. Then, the EFAs and the parallel analyses were conducted on the unidimensional 10-item CD-RISC versions (Campbell-Sills & Stein, 2007; Guarnizo Guzmán et al., 2019). The

<sup>1</sup><https://ddd.uab.cat/record/205870>

**Table 1.** Loading Matrix, Item Statistics, and Tau-Parameters for the CD-RISC-10

Item	Factor loading	Measure	SE Measure	Infit	Outfit	tau parameters			
						1	2	3	4
1	.572	-0.999	0.0536	0.935	0.983	-0.584	-1.152	0.334	1.403
4	.682	-0.948	0.0530	0.708	0.719	-0.790	-1.435	0.518	1.708
6	.575	-0.724	0.0509	1.001	0.990	-1.070	-0.699	0.666	1.104
7	.635	-0.351	0.0485	0.960	0.980	-1.074	-0.557	0.590	1.042
8	.555	-1.347	0.0581	1.377	1.255	-0.003	-0.478	-0.165	0.646
11	.714	-1.206	0.0561	0.807	0.806	-1.167	-0.912	0.670	1.410
14	.502	-0.211	0.0480	1.232	1.235	-0.665	-0.629	0.279	1.016
16	.528	-0.356	0.0485	1.196	1.225	-0.697	-0.661	0.447	0.912
17	.771	-1.043	0.0541	0.919	0.870	-0.267	-1.045	0.365	0.947
19	.554	-0.558	0.0497	0.957	1.003	-1.893	-0.339	0.897	1.336

Note. Factor loading = Estimator is WLSMV and Rotation is oblimin; Measure = item difficulty; SE Measure = Standard Error measure; Infit = Information-weighted mean square statistic; Outfit = Outlier-sensitive means square statistic.

1-factor model of the Campbell-Sills and Stein CD-RISC-10 was confirmed by parallel analysis, whereas the Eigenvalues from the EFA overlapped with those from the parallel analysis in the Guarnizo Guzmán et al. (2019) CD-RISC model. Eigenvalues of the EFAs and parallel analyses for the CD-RISC models are displayed at the online Supplementary material Table S2. The EFA for the Campbell-Sills and Stein CD-RISC-10 (see its factor loading in Table 1) provided a good fit,  $\chi^2 = 83.429$  (35),  $p < .001$ ; CFI = .979, TLI = .973, and RMSEA = .055, 90% CI [.040, .070]. After supporting the CD-RISC-25 unidimensionality, Guarnizo Guzmán et al. (2019) developed a new 10-item CD-RISC version removing from the original CD-RISC-25 those items with lower item-total correlations and factor loadings, as well as those reported by adolescents as being difficult to understand and more conceptually similar. However, the goodness of fit of Campbell-Sills and Stein's CD-RISC-10 version, supported among adults cross-culturally (Pulido-Martos et al., 2020) and Chinese adolescents (Chen et al., 2022), was not tested in the sample of Colombian adolescents. Given these findings, Campbell-Sills and Stein's CD-RISC-10 was chosen for the subsequent analyses.

Campbell-Sills and Stein's CD-RISC-10 items were shown to be essentially tau-equivalent with acceptable fit indices, CFI and TLI > 0.90 (Marsh et al., 2004), and RMSEA < 0.10 (Weston & Gore, 2006):  $\chi^2 = 192.100$  (44),  $p < .001$ ; CFI = .935, TLI = .934, and RMSEA = .085, 90% CI [.073, .098]. Thus, Rasch analyses were suitable. Table 1 shows the Rasch analysis for Campbell-Sills and Stein's CD-RISC-10, including item statistics and tau-parameters. All the infit and outfit values fell in the adequate range (0.708–1.377), showing an adequate model fit to the data. Figure 1 represents the Wright map of the Campbell-Sills and Stein's CD-RISC-10. It indicates that participant's resilience level presented wide distribution, albeit the item difficulty levels showed biased toward low levels of resilience, and high participant's resilience levels were not covered. The thresholds between response categories generally followed a similar trend for all the items (i.e., individuals with higher latent trait levels are more likely to choose higher response categories). However, Item 8 presented low discriminant power at the first 3 tau parameters; whereas Items 1, 4 and 17 did so at the first 2 tau parameters. In this line, Figure 2, via

response option characteristic curves, illustrates this overlap among the corresponding response categories of these items and supports the adequacy of the other item's response category distributions.


The results of the MI across sex of the CD-RISC-10 are presented in Table 2. The fit indices of the CFAs for male and female groups were adequate. Thus, a multigroup CFA was conducted. Results indicated good fit indices and minimal changes, fulfilling the criteria recommended to consider a model invariant when metric and scalar invariance are examined. Additionally, DIF analyses revealed nonsignificant differences in item responses as a function of sex, with all the adjusted  $p$ -values above .05 (see Table 3).

To explore criterion-related validity of the CD-RISC-10, the scores of the scale were associated with sex, well-being and health-related quality of life, emotional and conduct problems, and specific symptoms for anxiety, depression, and OCD (see Table 4). Large associations ( $r > .50$ ) were observed with health-related quality of life, MDD symptoms, and emotional symptoms. Additionally, moderate-to-large effect sizes ( $r = .30$ – $.50$ ) were found between resilience and SP, OCD, PD symptoms, and peer and conduct problems. Finally, resilience showed small-to-moderate associations ( $r = .10$ – $.30$ ) between the male sex and subscales measuring hyperactivity, prosocial behavior and symptomatology of SAD and GAD. Furthermore, the Campbell-Sills and Stein CD-RISC-10 presented good internal consistency (Categorical omega = .82).

## Discussion

The current study explored the psychometric properties of the CD-RISC in Spanish adolescents. Factor structure, Rasch analyses, MI by sex, and criterion validity regarding mental health outcomes were tested on the best CD-RISC model extracted.

The EFA conducted on the CD-RISC-25 suggests that resilience assessed by this scale is a unitary construct, in line with previous research (Burns & Anstey, 2010; Campbell-Sills & Stein, 2007; García-León et al., 2019; Guarnizo Guzmán et al., 2019). By contrast, other studies propose the existence of more than 1 factor, including the original CD-RISC study (e.g., Connor & Davidson, 2003; Fu et al., 2014; Jorgensen & Seedat, 2008; Yu & Zhang, 2007).



**Figure 1.** Wright Map (Person-Item Map).

Note. The number of each item corresponds to the CD-RISC-10.


Notwithstanding, these studies present some methodological flaws in the exploratory factor analyses, ignoring the recommendations for good practice (Fabrigar et al., 1999; Goretzko et al., 2021; Izquierdo et al., 2014). These shortcomings include:

1. The use of Principal Component Analysis as extraction method (Jorgensen & Seedat, 2008; Fu et al., 2014; Yu & Zhang, 2007), which is considered less sensitive than other methods.
2. The misestimation in the number of factors using just the eigenvalue as a criterion without conducting complementary analyses such as parallel analyses (Connor & Davidson, 2003; Jorgensen & Seedat, 2008; Yu & Zhang, 2007).
3. The employment of orthogonal rotation (e.g., Orthomax, Varimax, etc.) impeding factor intercorrelations (Connor & Davidson, 2003; Jorgensen & Seedat, 2008; Yu & Zhang, 2007).
4. And finally, the arbitrariness to justify the identification of factors composed by items conceptually different (Connor & Davidson, 2003; Fu et al., 2014; Jorgensen & Seedat, 2008; Yu & Zhang, 2007).

Accordingly, the 10-item unidimensional CD-RISC of Campbell-Sills and Stein emerged as the most appropriate model, in agreement with analyses conducted in Spanish adult population (Pulido-Martos et al., 2020). This model has also been previously supported among adolescent samples (Chen et al., 2022). When the psychometric properties of the different CD-RISC versions are compared (Burns & Anstey, 2010; Gonzalez et al., 2016; Peng et al., 2014; Kuiper et al., 2019; Pulido-Martos et al., 2020), Campbell-Sills and Stein's CD-RISC-10 model emerges as psychometrically superior. Furthermore, this model has been successfully validated across different cultural settings (Pulido-Martos et al.,

2020). However, the CD-RISC research focused on adolescents did not explore the goodness of fit of Campbell-Sills and Stein's CD-RISC-10 version, although the original 25-item CD-RISC version (Connor & Davidson, 2003) could not be replicated. Consequently, other versions have been developed (Fu et al., 2014; Guarnizo Guzmán et al., 2019; Jorgensen & Seedat, 2008). In this sense, the Colombian CD-RISC-10 version for adolescents (Guarnizo Guzmán et al., 2019), which shares just 3 items with the Campbell-Sills and Stein's CD-RISC-10, was developed when the CD-RISC-25 emerged as a 1-factor construct. Yet, the Campbell-Sills and Stein's CD-RISC-10, which followed a conceptual approach to coherently define the resilience factor, was not tested on such adolescent sample. In short, the findings of the present study are in line with this trend, suggesting that the Campbell-Sills and Stein CD-RISC-10 model is also the best to assess resilience in adolescents.

Rasch analyses confirmed the goodness of fit of all the items, in line with some of the previous psychometric Campbell-Sills and Stein CD-RISC-10 studies (Pulido-Martos et al., 2020) but not with all of them (Ehrich et al., 2017; Heritage et al., 2021). Our data showed that the CD-RISC-10 presented a balanced distribution of participants according to levels of resilience. However, item difficulty levels scarcely discriminated in high levels of resilience, a finding frequently observed in previous CD-RISC studies (Arias González et al., 2015; Heritage et al., 2021; Papini et al., 2021). The inclusion of new difficult items may be considered in the CD-RISC-10 to discriminate better among highly resilient participants. Our findings also indicate that some CD-RISC-10 items present response category misfits, especially on Items 8 (to a larger extent), 1, 4, and 17 in their first categories, unlike adult samples (Heritage et al., 2021; Waddimba et al., 2022).



**Figure 2.** Response Option Characteristic Curves for the Items of the CD-RISC-10.

Regarding sex and resilience, the present study shows MI across sex of Campbell-Sills and Stein's CD-RISC-10 in Spanish adolescents, replicating findings from adults (Gonzalez *et al.*, 2016; Pulido-Martos *et al.*, 2020) and other adolescent samples (Chen

*et al.*, 2022). Despite our sex balanced sample, independent CFAs for sex indicated that girls presented a slightly worse, albeit still good, fit; in line with previous research (Chen *et al.*, 2022; Pulido-Martos *et al.*, 2020). The absence of DIF across sex was also

**Table 2.** Model fit and Measurement Invariance of the CD-RISC-10 Across Sex

	$\chi^2_{S-B} (df)$	CFI	TLI	RMSEA	90% CI	Ref	$\Delta CFI$	$\Delta RMSEA$
1-factor model with 10 items								
M1. Males	55.748(35)*	.977	.971	.051	[.023, .075]	–	–	–
M2. Females	78.536(35)**	.965	.954	.073	[.052, .095]	–	–	–
Invariance testing across sex								
M3. Configural	132.421(70)**	.971	.963	.062	[.046, .078]	–	–	–
M4. Metric	131.392(79)**	.976	.972	.054	[.037, .069]	M3	.005	–.008
M5. Scalar	169.452(108)**	.972	.976	.050	[.035, .064]	M4	–.004	–.004

Note.  $\chi^2_{S-B}$  = Satorra-Bentler scaled chi-square; *df* = degrees of freedom; CFI = Comparative Fit Index; RMSEA = Root Mean Square Error of Approximation; 90% CI = 90% confidence interval of the RMSEA; MX. = number of reference model; Ref = reference model;  $\Delta CFI$  = change in CFI;  $\Delta RMSEA$  = change in RMSEA.

\**p* < .05. \*\**p* < .001.

**Table 3.** Likelihood Ratio Chi-square Statistics of the CD-RISC-10 Items

Item	Statistic	<i>p</i>	<i>Adj.p</i>
1	1.469	.480	.762
4	4.299	.117	.388
6	4.685	.096	.388
7	0.544	.762	.762
8	3.263	.196	.489
11	5.903	.052	.388
14	0.828	.661	.762
16	2.374	.305	.610
17	0.550	.760	.762
19	0.614	.736	.762

Note. *Adj.p* = The adjusted *p*-values by likelihood ratio test using multiple comparison.

supported in the CD-RISC-10, in line with other studies (Ehrich et al., 2017; Heritage et al., 2021; Pulido-Martos et al., 2020). As reported for Spanish adults (Pulido-Martos et al., 2020), adolescent males presented significantly more resilience than adolescent females in our study. In this sense, two meta-analyses based on sex differences in resilience presented mixed results: One showed no sex differences (Ari & Çarkit, 2020), and the other displayed higher levels of resilience in favor of males (Yılmaz Koğar & Gök, 2021). These findings may be substantially explained by the use of different instruments to assess resilience across studies.

Therefore, the core of resilience, independently from the socio-cultural setting and developmental stage, appears to encompass the hardiness and persistence aspects assessed by the CD-RISC-10 of Campbell-Sills and Stein. In the present study, high positive correlations were observed between the CD-RISC-10 and health-related quality of life, whereas an inverse pattern was shown with MDD and emotional symptoms; in line with the literature (Kuiper et al., 2019; Waddimba et al., 2022; Wollny & Jacobs, 2021). Accordingly, stronger relationships were found with psychopathological variables for the CD-RISC-10 compared to the CD-RISC-25 (Kuiper et al., 2019). Thus, the resilience trait may be primarily explained by emotional stability and self-confidence. In this sense, resilience showed the highest association with the personality trait of neuroticism, with an estimated average coefficient of -.46

**Table 4.** Correlation Coefficients of the Study Variables related to the CD-RISC-10

	CD-RISC-10
Sex	–.24
HRQoL	–.57
SDQ	
Emotional Symptoms	–.52
Conduct Problems	–.30
Hyperactivity	–.23
Peer Problems	–.34
Prosocial Behaviors	.23
RCADS-30	
MDD	–.57
PD	–.41
SF	–.44
SAD	–.19
GAD	–.20
OCD	–.34

Note. Sex (Male = 1, Female = 2); HRQoL = Health-related quality of life; SDQ = Strength and Difficulties Questionnaire; RCADS-30 = Revised Child Anxiety and Depression Scale-30; MDD = Major Depressive Disorder; PD = Panic Disorder; SF = Social Phobia; SAD = Separation Anxiety Disorder; GAD = Generalized Anxiety Disorder; OCD = Obsessive-Compulsive Disorder. All correlation values are significant at *p* < .001. Cohen’s *d*-values 0.10, 0.30, and 0.50 correspond to small, medium, and large effects, respectively (Cohen, 1992).

(see Oshio et al., 2018, for a meta-analysis focused on the relationship between resilience and the Big Five personality traits). Hence, when resilience is treated, internalizing symptoms consequently improve (Ang et al., 2022; Connor & Davidson, 2003; Davidson et al., 2005).

The current study is not without limitations. Firstly, the participants comprise a convenience sample obtained from different high schools in Spain. In this sense, the use of larger samples of adolescents would imply better statistical power of the findings. In addition, the data was self-reported and, therefore, subjected to well-known biases such as social desirability. Apart from that, minor differences were found in the CD-RISC items’ drafting among Spanish translations, which might hinder the comparison of findings (García-León et al., 2019; Notario-Pacheco et al., 2011;

Pulido-Martos et al., 2020). And finally, the CD-RISC-10 items present problems to discriminate among adolescents with high levels of resilience. Despite these limitations, our research provides meaningful data, examining the best CD-RISC model to assess adolescent resilience, something scarcely explored in the literature (Fu et al., 2014; Guarnizo Guzmán et al., 2019; Jorgensen & Seedat, 2008; Yu et al., 2011).

In conclusion, this study thoroughly examined psychometric properties of the CD-RISC in Spanish adolescents. The unidimensional CD-RISC-10 of Campbell-Sills and Stein emerges as the best CD-RISC model to assess resilience among Spanish adolescents. The data highlight the need to cross-culturally examine the psychometric properties of Campbell-Sills and Stein's CD-RISC-10 among adolescents to test its universality, initially suggested among this age group (Chen et al., 2022) and highly supported in adult populations (Pulido-Martos et al., 2020).

**Supplementary material.** The supplementary material for this article can be found at <http://doi.org/10.1017/SJP.2024.3>.

**Funding statement.** The Ministerio de Economía, Industria y Competitividad of the Government of Spain (Grant Number PSI2017-88280-R) and the Ministerio de Ciencia, Innovación y Universidades of Spain (Grant Number RTI2018-097124-B-100) funded this research. Additionally, it was supported by the Agencia Estatal de Investigación (Grant Number: PID2019-111138RB-I00/AEI/10.13039/501100011033) and partly supported by the Network for the PROMotion of EMotional Health and Wellbeing of Adolescents (PROEMO) and the Research Network on Chronicity, Primary Care and Prevention and Health Promotion (RICAPPS) [Grant Number RED2022-134247-T]; and also by the individual funding of the European Union – NextGenerationEU (FLF, Grant Number MGS/2021/02).

**Competing interest.** None.

**Authorship credit.** Francisco J. López-Fernández: Conceptualization; Methodology; Formal analysis; Writing - Original draft preparation. Paula Morales-Hidalgo: Writing - review & editing. Josefa Canals: Funding acquisition; Project administration; Writing - review & editing. Juan C. Marzo: Funding acquisition; Project administration; Writing - review & editing. Luis Joaquín García-López: Funding acquisition; Project administration; Writing - review & editing. José A. Piqueras: Conceptualization; Supervision; Funding acquisition; Project administration; Writing - review & editing.

**Data sharing.** The data is openly available in Open Science Framework (OSF) at [https://osf.io/l7rp3/?view\\_only=1cdcc49bc31d46f4a9f6ebdc4bdfb4d6](https://osf.io/l7rp3/?view_only=1cdcc49bc31d46f4a9f6ebdc4bdfb4d6).

## References

- Ari, F. A., & Çarkit, E. (2020). Investigation of resilience in terms of gender: A meta-analysis study. *Research on Education and Psychology*, *4*, 34–52.
- Ang, W. H. D., Lau, S. T., Cheng, L. J., Chew, H. S. J., Tan, J. H., Shorey, S., & Lau, Y. (2022). Effectiveness of resilience interventions for higher education students: A meta-analysis and meta-regression. *Journal of Educational Psychology*, *114*, 1670–1694. <https://doi.org/10.1037/edu0000719>
- Arias González, V. B., Crespo Sierra, M. T., Arias Martínez, B., Martínez-Molina, A., & Ponce, F. P. (2015). An in-depth psychometric analysis of the Connor-Davidson Resilience Scale: Calibration with Rasch-Andrich model. *Health and Quality of Life Outcomes*, *13*, Article 154. <https://doi.org/10.1186/s12955-015-0345-y>
- Bobes, J., Bascaran, M. T., García-Portilla, M. P., Bousoño, M., Sáiz, P. A., & Wallace, D. H. (2001). *Banco de instrumentos básicos de psiquiatría clínica* [Bank of basic instruments of clinical psychiatry]. Psiquiatría ED.
- Brownlee, K., Rawana, J., Franks, J., Harper, J., Bajwa, J., O'Brien, E., & Clarkson, A. (2013). A systematic review of strengths and resilience outcome literature relevant to children and adolescents. *Child and Adolescent Social Work Journal*, *30*, 435–459. <https://doi.org/10.1007/s10560-013-0301-9>
- Burns, R. A., & Anstey, K. J. (2010). The Connor-Davidson Resilience Scale (CD-RISC): Testing the invariance of a uni-dimensional resilience measure that is independent of positive and negative affect. *Personality and Individual Differences*, *48*, 527–531. <https://doi.org/10.1016/j.paid.2009.11.026>
- Campbell-Sills, L., & Stein, M. B. (2007). Psychometric analysis and refinement of the Connor-Davidson Resilience Scale (CD-RISC): Validation of a 10-item measure of resilience. *Journal of Traumatic Stress*, *20*, 1019–1028. <https://doi.org/10.1002/jts.20271>
- Chen, F. F. (2007). Sensitivity of goodness of fit indexes to lack of measurement invariance. *Structural Equation Modeling: A Multidisciplinary Journal*, *14*, 464–504. <http://doi.org/10.1080/10705510701301834>
- Chen, W., Liang, Y., Yang, T., Gao, R., & Zhang, G. (2022). Validity and longitudinal invariance of the 10-Item Connor-Davidson Resilience Scale (CD-RISC-10) in Chinese left-behind and non-left-behind children. *Psychological Reports*, *125*, 2274–2291. <https://doi.org/10.1177/00332941211013531>
- Cheung, G. W., & Rensvold, R. B. (2002). Evaluating goodness-of-fit indexes for testing measurement invariance. *Structural Equation Modeling: A Multidisciplinary Journal*, *9*, 233–255. [http://doi.org/10.1207/s15328007sem0902\\_5](http://doi.org/10.1207/s15328007sem0902_5)
- Chorpita, B. F., Yim, L., Moffitt, C., Umemoto, L. A., & Francis, S. E. (2000). Assessment of symptoms of DSM-IV anxiety and depression in children: A revised Child Anxiety and Depression scale. *Behaviour Research and Therapy*, *38*, 835–855. [https://doi.org/10.1016/s0005-7967\(99\)00130-8](https://doi.org/10.1016/s0005-7967(99)00130-8)
- Cohen, J. (1992). A power primer. *Psychological Bulletin*, *112*, 155–159. <https://doi.org/10.1037/0033-2909.112.1.155>
- Connor, K. M., & Davidson, J. R. T. (2003). Development of a new resilience scale: The Connor-Davidson Resilience Scale (CD-RISC). *Depression & Anxiety*, *18*, 76–82. <https://doi.org/10.1002/da.10113>
- Davidson, J. R. T., Payne, V. M., Connor, K. M., Foa, E. B., Rothbaum, B. O., Hertzberg, M. A., & Weisler, R. H. (2005). Trauma, resilience and salivary cortisol: Effects of treatment in post-traumatic stress disorder. *International Clinical Psychopharmacology*, *20*, 43–48. <https://doi.org/10.1097/00004850-200501000-00009>
- Dorans, N. J., Schmitt, A. P., & Bleistein, C. A. (1992). The standardization approach to assessing comprehensive differential item functioning. *Journal of Educational Measurement*, *29*, 309–319. <https://doi.org/10.1111/j.1745-3984.1992.tb00379.x>
- Ehrlich, J., Mornane, A., & Powern, T. (2017). Psychometric validation of the 10-item Connor-Davidson Resilience Scale. *Journal of Applied Measurement*, *18*, 122–136.
- Fabrigar, L. R., Wegener, D. T., MacCallum, R. C., & Strahan, E. J. (1999). Evaluating the use of exploratory factor analysis in psychological research. *Psychological Methods*, *4*, 272–299. <https://doi.org/10.1037/1082-989x.4.3.272>
- Feder, A., Nestler, E. J., & Charney, D. S. (2009). Psychobiology and molecular genetics of resilience. *Nature Reviews Neuroscience*, *10*, 446–457. <https://doi.org/10.1038/nrn2649>
- Fletcher, D., & Sarkar, M. (2013). Psychological resilience: A review and critique of definitions, concepts and theory. *European Psychologist*, *18*, 12–23. <https://doi.org/10.1027/1016-9040/a000124>
- Flora, D. B. (2020). Your coefficient alpha is probably wrong, but which coefficient omega is right? A tutorial on using R to obtain better reliability estimates. *Advances in Methods and Practices in Psychological Science*, *3*, 484–501. <https://doi.org/10.1177/2515245920951747>
- Fu, C., Leoutsakos, J.-M., & Underwood, C. (2014). An examination of resilience cross-culturally in child and adolescent survivors of the 2008 China earthquake using the Connor-Davidson Resilience Scale (CD-RISC). *Journal of Affective Disorders*, *155*, 149–153. <https://doi.org/10.1016/j.jad.2013.10.041>
- García-León, M. Á., González-Gomez, A., Robles-Ortega, H., Padilla, J. L., & Peralta-Ramirez, I. (2019). Propiedades psicométricas de la Escala de Resiliencia de Connor y Davidson (CD-RISC) en población española [Psychometric properties of the Connor-Davidson Resilience Scale, CD-RISC, in Spanish population]. *Anales de Psicología*, *35*, 33–40. <https://doi.org/10.6018/analesps.35.1.314111>
- Garrido, L. E., Abad, F. J., & Ponsoda, V. (2013). A new look at Horn's parallel analysis with ordinal variables. *Psychological Methods*, *18*, 454–474. <https://doi.org/10.1037/a0030005>
- Gómez-Beneyto, M., Nolasco, A., Moncho, J., Pereyra-Zamora, P., Tamayo-Fonseca, N., Munarriz, M., Salazar, J., Tabarés-Seisdedos, R., & Girón, M.



- (2013). Psychometric behaviour of the Strengths and Difficulties Questionnaire (SDQ) in the Spanish National Health Survey 2006. *BMC Psychiatry*, *13*, Article 95. <http://doi.org/10.1186/1471-244x-13-95>
- Gonzalez, S. P., Moore, E. W. G., Newton, M., & Galli, N. A.** (2016). Validity and reliability of the Connor-Davidson Resilience Scale (CD-RISC) in competitive sport. *Psychology of Sport and Exercise*, *23*, 31–39. <https://doi.org/10.1016/j.psychsport.2015.10.005>
- Goodman, R.** (2001). Psychometric properties of the Strengths and Difficulties Questionnaire. *Journal of the American Academy of Child & Adolescent Psychiatry*, *40*, 1337–1345. <http://doi.org/10.1097/00004583-200111000-00015>
- Goretzko, D., Pham, T. T. H., & Bühner, M.** (2021). Exploratory factor analysis: Current use, methodological developments and recommendations for good practice. *Current Psychology*, *40*, 3510–3521. <https://doi.org/10.1007/s12144-019-00300-2>
- Guarnizo Guzmán, C. P., García Martín, M. B., Suárez Falcón, J. C. & Sierra, M. A.** (2019). Psychometric properties of the Connor-Davidson Resilience Scale (CD-RISC) on vulnerable Colombian adolescents. *International Journal of Psychology and Psychological Therapy*, *19*, 277–289.
- Heritage, B., Al Asadi, A. A., & Hegney, D. G.** (2021). Examination of the Connor-Davidson Resilience Scale 10 (CD-RISC-10) using the polytomous Rasch model. *Psychological Assessment*, *33*, 672–684. <https://doi.org/10.1037/pas0001011>
- Hladká, A., & Martinková, P.** (2020). difNLR: Generalized logistic regression models for DIF and DDF Detection. *The R Journal*, *12*, 300–323. <https://doi.org/10.32614/rj-2020-014>
- Hu, L.-t., & Bentler, P. M.** (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal*, *6*, 1–55. <https://doi.org/10.1080/10705519909540118>
- Hu, T., Zhang, D., & Wang, J.** (2015). A meta-analysis of the trait resilience and mental health. *Personality and Individual Differences*, *76*, 18–27. <https://doi.org/10.1016/j.paid.2014.11.039>
- Izquierdo, I., Olea, J., & Abad, F. J.** (2014). Exploratory factor analysis in validation studies: Uses and recommendations. *Psicothema*, *26*, 395–400. <https://doi.org/10.7334/psicothema2013.349>
- Jorgensen, I. E., & Seedat, S.** (2008). Factor structure of the Connor-Davidson Resilience Scale in South African adolescents. *International Journal of Adolescent Medicine and Health*, *20*, 23–32. <https://doi.org/10.1515/ijamh.2008.20.1.23>
- Khan, A., Chien, C.-W., & Brauer, S. G.** (2013). Rasch-based scoring offered more precision in differentiating patient groups in measuring upper limb function. *Journal of Clinical Epidemiology*, *66*, 681–687. <https://doi.org/10.1016/j.jclinepi.2012.12.014>
- Kuiper, H., van Leeuwen, C. C. M., Stolwijk-Swüste, J. M., & Post, M. W. M.** (2019). Measuring resilience with the Connor-Davidson Resilience Scale (CD-RISC): Which version to choose? *Spinal Cord*, *57*, 360–366. <https://doi.org/10.1038/s41393-019-0240-1>
- Li, M. H., & Nishikawa, T.** (2012). The relationship between active coping and trait resilience across US and Taiwanese college student samples. *Journal of College Counseling*, *15*, 157–171. <https://doi.org/10.1002/j.2161-1882.2012.00013.x>
- Linacre, J. M.** (1999). Investigating rating scale category utility. *Journal of Outcome Measurement*, *3*, 103–122.
- Linacre, J. M.** (2002). What do infit and outfit, mean-square and standardized mean. *Rasch Measurement Transactions*, *16*, 878.
- Liu, D. W. Y., Fairweather-Schmidt, A. K., Burns, R. A., & Roberts, R. M.** (2015). The Connor-Davidson Resilience Scale: Establishing invariance between gender across the lifespan in a large community-based study. *Journal of Psychopathology and Behavioral Assessment*, *37*, 340–348. <https://doi.org/10.1007/s10862-014-9452-z>
- Lorenzo-Seva, U., & Ferrando, P. J.** (2006). FACTOR: A computer program to fit the exploratory factor analysis model. *Behavior Research Methods*, *38*, 88–91. <http://doi.org/10.3758/BF03192753>
- MacCallum, R. C., Browne, M. W., & Sugawara, H. M.** (1996). Power analysis and determination of sample size for covariance structure modeling. *Psychological Methods*, *1*, 130–149. <http://doi.org/10.1037/1082-989x.1.2.130>
- Marsh, H. W., Hau, K.-T., & Wen, Z.** (2004). In search of golden rules: Comment on hypothesis-testing approaches to setting cutoff values for fit indexes and dangers in overgeneralizing Hu and Bentler's (1999) findings. *Structural Equation Modeling: A Multidisciplinary Journal*, *11*, 320–341. [https://doi.org/10.1207/s15328007sem1103\\_2](https://doi.org/10.1207/s15328007sem1103_2)
- Masters, G. N., & Wright, B. D.** (1997). The Partial Credit Model. In W. J. van der Linden & R. K. Hambleton (Eds.), *Handbook of Modern Item Response Theory* (pp. 101–121). Springer. [https://doi.org/10.1007/978-1-4757-2691-6\\_6](https://doi.org/10.1007/978-1-4757-2691-6_6)
- Muthén, L. K., & Muthén, B. O.** (1997–2017). *Mplus user's guide* (8<sup>th</sup> Ed.). Muthén & Muthén. [https://www.statmodel.com/download/usersguide/MplusUserGuideVer\\_8.pdf](https://www.statmodel.com/download/usersguide/MplusUserGuideVer_8.pdf)
- Myers, N. D., Wolfe, E. W., Feltz, D. L., & Penfield, R. D.** (2006). Identifying differential item functioning of rating scale items with the Rasch model: An introduction and an application. *Measurement in Physical Education and Exercise Science*, *10*, 215–240. [https://doi.org/10.1207/s15327841mpee1004\\_1](https://doi.org/10.1207/s15327841mpee1004_1)
- Notario-Pacheco, B., Solera-Martínez, M., Serrano-Parra, M. D., Bartolomé-Gutiérrez, R., García-Campayo, J., & Martínez-Vizcaino, V.** (2011). Reliability and validity of the Spanish version of the 10-item Connor-Davidson Resilience Scale (10-item CD-RISC) in young adults. *Health and Quality of Life Outcomes*, *9*, Article 63. <https://doi.org/10.1186/1477-7525-9-63>
- Oshio, A., Taku, K., Hirano, M., & Saeed, G.** (2018). Resilience and Big Five personality traits: A meta-analysis. *Personality and Individual Differences*, *127*, 54–60. <https://doi.org/10.1016/j.paid.2018.01.048>
- Papini, N., Kang, M., Ryu, S., Griese, E., Wingert, T., & Herrmann, S.** (2021). Rasch calibration of the 25-item Connor-Davidson resilience scale. *Journal of Health Psychology*, *26*, 1976–1987. <https://doi.org/10.1177/1359105320904769>
- Peng, L., Zhang, J., Chen, H., Zhang, Y., Li, M., Yu, Y., & Liu, B.** (2014). Comparison among different versions of Connor-Davidson Resilience Scale (CD-RISC) in rehabilitation patients after unintentional injury. *Journal of Psychiatry*, *17*, Article 1000153. <http://doi.org/10.4172/2378-5756.1000153>
- Pulido-Martos, M., Fernández-Sánchez, M. D., & Lopez-Zafra, E.** (2020). Measurement invariance across gender and age in the Connor-Davidson Resilience Scale (CD-RISC) in a Spanish general population. *Quality of Life Research*, *29*, 1373–1384. <https://doi.org/10.1007/s11136-019-02389-1>
- Putnick, D. L., & Bornstein, M. H.** (2016). Measurement invariance conventions and reporting: The state of the art and future directions for psychological research. *Developmental Review*, *41*, 71–90. <https://doi.org/10.1016/j.dr.2016.06.004>
- Ravens-Sieberer, U., Erhart, M., Rajmil, L., Herdman, M., Auquier, P., Bruil, J., Power, M., Duer, W., Abel, T., Czemy, L., Mazur, J., Czimbalmos, A., Tountas, Y., Hagquist, C., Kilroe, J., & the European KIDSCREEN Group.** (2010). Reliability, construct and criterion validity of the KIDSCREEN-10 score: A short measure for children and adolescents' well-being and health-related quality of life. *Quality of Life Research*, *19*, 1487–1500. <https://doi.org/10.1007/s11136-010-9706-5>
- Sandín, B., Chorot, P., Valiente, R. M., & Chorpita, B. F.** (2010). Development of a 30-Item version of the revised Child Anxiety and Depression Scale. *Revista de Psicopatología y Psicología Clínica*, *15*, 165–178. <https://doi.org/10.5944/rppc.vol.15.num.3.2010.4095>
- Satorra, A.** (2000). Scaled and adjusted restricted tests in multisample analysis of moment structures. In R. D. H. Heijmans, D. S. G. Pollock, & A. Satorra (Eds.), *Innovations in multivariate statistical analysis: Advanced studies in theoretical and applied econometrics* (Vol. 36, pp. 233–247). Springer. [http://doi.org/10.1007/978-1-4615-4603-0\\_17](http://doi.org/10.1007/978-1-4615-4603-0_17)
- Seol, H.** (2022). *SnowIRT: Item Response Theory for jamovi* (jamovi module) [Computer software]. Jamovi. <https://github.com/hyunsooseol/snowIRT>
- The jamovi project** (2022). *Jamovi*. (Version 2.3) [Computer software]. <https://www.jamovi.org>
- Timmerman, M. E., & Lorenzo-Seva, U.** (2011). Dimensionality assessment of ordered polytomous items with parallel analysis. *Psychological Methods*, *16*, 209–220. <https://doi.org/10.1037/a0023353>
- Viladrich, C., Angulo-Brunet, A., & Doval, E.** (2017). A journey around alpha and omega to estimate internal consistency reliability. *Anales de Psicología*, *33*, 755–782. <https://doi.org/10.6018/analesps.33.3.268401>
- Waddimba, A. C., Baker, B. M., Pogue, J. R., McAuliffe, M. P., Bennett, M. M., Baxter, R. D., Mohr, D. C., & Warren, A. M.** (2022). Psychometric validity and reliability of the 10- and 2-item Connor-Davidson resilience scales among a national sample of Americans responding to the Covid-19 pandemic: An item response theory analysis. *Quality of Life Research*, *31*, 2819–2836. <https://doi.org/10.1007/s11136-022-03125-y>

- West, S. G., Taylor, A. B., & Wu, W. (2012). Model fit and model selection in structural equation modeling. In R. H. Hoyle (Ed.), *Handbook of structural equation modeling* (pp. 209–231). The Guilford Press.
- Weston, R., & Gore, P. A., Jr. (2006). A brief guide to structural equation modeling. *The Counseling Psychologist*, *34*, 719–751. <https://doi.org/10.1177/0011000006286345>
- Windle, G., Bennett, K. M., & Noyes, J. (2011). A methodological review of resilience measurement scales. *Health and Quality of Life Outcomes*, *9*, Article 8. <https://doi.org/10.1186/1477-7525-9-8>
- Wollny, A. I., & Jacobs, I. (2021). Validity and reliability of the German versions of the CD–RISC–10 and CD–RISC–2. *Current Psychology*, *42*, 3437–3448. <https://doi.org/10.1007/s12144-021-01670-2>
- Yılmaz Koğar, E., & Gök, A. (2021). A meta-analysis study on gender differences in psychological resilience levels. *Kıbrıs Türk Psikiyatri ve Psikoloji Dergisi*, *3*, 132–143. <https://doi.org/10.35365/ctjpp.21.2.15>
- Yu, X.-n., Lau, J. T. F., Mak, W. W. S., Zhang, J., Lui, W. W. S., Zhang, J. (2011). Factor structure and psychometric properties of the Connor-Davidson Resilience Scale among Chinese adolescents. *Comprehensive Psychiatry*, *52*, 218–224. <https://doi.org/10.1016/j.comppsy.2010.05.010>
- Yu, X., & Zhang, J. (2007). Factor analysis and psychometric evaluation of the Connor-Davidson Resilience Scale (CD-RISC) with Chinese people. *Social Behavior and Personality: An International Journal*, *35*, 19–30. <https://doi.org/10.2224/sbp.2007.35.1.19>